



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|---|--|
| (51) International Patent Classification⁶ : C07K 14/47, 14/52, C12N 15/12, 15/19, 15/63, A61K 38/16, 38/19, 48/00 | A1 | (11) International Publication Number: WO 99/29728 (43) International Publication Date: 17 June 1999 (17.06.99) |
| (21) International Application Number: PCT/US98/26291 (22) International Filing Date: 11 December 1998 (11.12.98) (30) Priority Data: 60/069,281 11 December 1997 (11.12.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/069,281 (CON) Filed on 11 December 1997 (11.12.97) (71) Applicant (for all designated States except US): UNIVERSITY OF MARYLAND BIOTECHNOLOGY INSTITUTE [US/US]; 4321 Hartwick Road, College Park, MD 20740 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GALLO, Robert, C. [US/US]; 8513 Thornden Terrace, Bethesda, MD 02817 (US). DEVICO, Anthony, L. [US/US]; 4533 Peacock Avenue, Alexandria, VA 22304 (US). GARZINO-DEMO, Alfredo [IT/US]; 601 North Eutaw Street, Baltimore, MD 21201 (US). | (74) Agent: BARRETT, William, A.; Intellectual Property/Technology Law, P.O. Box 14329, Research Triangle Park, NC 27709 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> | |
| (54) Title: METHOD AND COMPOSITION TO ENHANCE THE EFFICACY OF A VACCINE USING CHEMOKINES | | |
| (57) Abstract The present invention relates to a method to enhance the efficacy of a vaccine in a subject treated with the vaccine comprising administering to the subject in combination with the vaccine a one or more chemokines. The present invention also relates to compositions of vaccines containing chemokines. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

METHOD AND COMPOSITION TO ENHANCE THE EFFICACY OF A VACCINE USING CHEMOKINES

1. CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Application Serial No. 60/069,281 filed December 11, 1997.

2. BACKGROUND OF THE INVENTION

The present invention relates to a method to enhance the efficacy of a vaccine by administration of a chemokine, such as macrophage derived chemokine (MDC), in conjunction with the vaccine. The present invention also relates to compositions useful in the method.

2.1. GENERATION OF AN IMMUNE RESPONSE

The introduction of a foreign antigen into an individual elicits an immune response consisting of two major components, the cellular and humoral immune responses, mediated by two functionally distinct populations of lymphocytes known as T and B cells, respectively (see generally Coutinho, 1991, *Immune System, Encyclopedia of Human Biology*, Vol. 4, Ed. Dulbecco, Academic Press, Inc.). A subset of T cells responds to antigen stimulation by producing lymphokines which "help" or activate various other cell types in the immune system.

Another T cell subset is capable of developing into antigen-specific cytotoxic effector cells, which can directly kill antigen-positive target cells. On the other hand, the B cell response is primarily carried out by secretory proteins, antibodies, which directly bind and neutralize antigens.

Helper T cells (TH) can be distinguished from classical cytotoxic T lymphocytes (CTL) and B cells by their cell surface expression of the glycoprotein marker CD4. Although the mechanism by which CD4⁺ TH function has not been fully elucidated, the existence of functionally distinct subsets within the CD4⁺ T cell compartment has been reported (Mosmann and Coffman, 1989, *Ann. Rev. Immunol.*

Z:145-173). In the mouse, type 1 helper T cells (TH1) produce interleukin-2 (IL-2) and τ -interferon (τ -IFN) upon activation, while type 2 helper T cells (TH2) produce IL-4 and IL-5. Based on the profile of lymphokine production, TH1 appear to be involved in promoting the activation and proliferation of other T cell subsets including CTL, whereas TH2 specifically regulate B cell proliferation and differentiation, antibody synthesis, and antibody class switching.

A second T cell subpopulation is the classical CTL which express the CD8 surface marker. Unlike most TH, these cells display cytolytic activity upon direct contact with target cells, rather than through the production of lymphokines. *In vivo*, CTL function is particularly important in situations where an antibody response alone is inadequate. Significant experimental evidence indicates that CTL rather than B cells and their antibody products play a principal role in the defense against viral infections and cancer.

A salient feature of both T and B cell responses is their exquisite specificity for the immunizing antigen; however, the mechanisms for antigen recognition differ between these two cell types. B cells recognize antigens by antibodies, either acting as cell surface receptors or as secreted proteins, which bind directly to antigens on a solid surface or in solution, whereas T cells only recognize antigens that have been processed or degraded into small fragments and presented on a solid phase such as the surface of antigen-presenting cells (APC). Additionally, antigenic fragments must be presented to T cells in association with major histocompatibility complex (MHC)-encoded class I or class II molecules. The MHC refers to a cluster of genes that encode proteins with diverse immunological functions. In man, the MHC is known as HLA. Class I gene products are found on all somatic cells, and they were originally discovered as targets of major transplantation rejection responses. Class II gene products are mostly expressed on cells of various hematopoietic lineages, and they are involved in cell-cell interactions in the immune system. Most importantly, MHC-encoded proteins have been shown to function as receptors for processed antigenic fragments on the surface of APC (Bjorkman et al., 1987, *Nature* 329:506-512).

Another level of complexity in the interaction between a T cell and an antigenic fragment is that it occurs only if the MHC molecules involved are the same on the APC and the responding T cells. In other words, a T cell specific for a particular antigenic epitope expresses a receptor having low affinity for self MHC

proteins, which when such MHC proteins on APC are occupied by the epitope, engage the T cell in a stronger interaction leading to antigen-specific T cell activation. The phenomenon of a T cell reacting with a processed antigen only when presented by cells expressing a matching MHC is known as MHC-restriction.

The specificity of T cell immune responses for antigens is a function of the unique receptors expressed by these cells. The T cell receptor (TCR) is structurally homologous to an antibody; it is a heterodimer composed of disulfide-linked glycoproteins. Four TCR polypeptide chains known as α , β , τ , and δ have been identified, although the vast majority of functional T cells express the $\alpha\beta$ heterodimeric TCR. Transfer of α and β genes alone into recipient cells was shown to be both necessary and sufficient to confer antigen specificity and MHC-restriction (Dembic et al., 1986, *Nature* 320:232-238). Thus, the $\alpha\beta$ TCR appears to be responsible for recognizing a combination of antigenic fragment and MHC determinants.

The apparent basis of MHC restriction is that $CD4^+$ T cells express $\alpha\beta$ TCR which recognize antigenic fragments physically associated with MHC class II proteins, while the TCR on $CD8^+$ CTL recognize MHC class I-associated fragments. Thus, $CD4^+$ T cells can recognize only a restricted class of APC that are class II⁺, whereas $CD8^+$ CTL can interact with virtually any antigen-positive cells, since all cells express class I molecules. $CD4^+$ CTL have been identified, and they are MHC class II restricted, and lyse target cells only if the latter express self-MHC class II determinants associated with specific antigenic fragments. Both CD4 and CD8 molecules also contribute to this interaction by binding to monotypic determinants on the MHC class II and I molecules, respectively.

A second type of TCR composed of $\tau\delta$ heterodimers is expressed by a small percentage of T cells, but the involvement of $\tau\delta$ T cells in antigen-specific recognition is still poorly understood. Some studies have shown that functionally active $\tau\delta$ T cells can be cytolytic in a MHC non-restricted manner.

In summary, the generation of an immune response begins with the sensitization of $CD4^+$ and $CD8^+$ T cell subsets through their interaction with APC that express MHC-class I or class II molecules associated with antigenic fragments. The sensitized or primed $CD4^+$ T cells produce lymphokines that participate in the activation of B cells as well as various T cell subsets. The sensitized $CD8^+$ T cells increase in numbers in response to lymphokines and are capable of destroying any

cells that express the specific antigenic fragments associated with matching MHC-encoded class I molecules. For example, in the course of a viral infection, CTL eradicate virally-infected cells, thereby limiting the progression of virus spread and disease development.

2.2. ANTIGEN PRESENTING CELLS

The presentation of antigens to T cells is carried out by specialized cell populations referred to as antigen presenting cells (APC). Typically, APC include macrophages/monocytes, B cells, and bone marrow derived dendritic cells (DC). APC are capable of internalizing exogenous antigens, cleaving them into smaller fragments in enzyme-rich vesicles, and coupling the fragments to MHC-encoded products for expression on the cell surface (Goldberg and Rock, 1992, *Nature* 357:375-379). Since APC express both MHC-encoded class I and class II glycoproteins, they can present antigenic fragments to both CD4⁺ and CD8⁺ T cells for the initiation of an immune response.

By definition, APC not only can present antigens to T cells with antigen-specific receptors, but can provide all the signals necessary for T cell activation. Such signals are incompletely defined, but probably involve a variety of cell surface molecules as well as cytokines or growth factors. Further, the factors necessary for the activation of naive or unprimed T cells may be different from those required for the re-activation of previously primed memory T cells. The ability of APC to both present antigens and deliver signals for T cell activation is commonly referred to as an accessory cell function. Although monocytes and B cells have been shown to be competent APC, their antigen presenting capacities *in vitro* appear to be limited to the re-activation of previously sensitized T cells. Hence, they are not capable of directly activating functionally naive or unprimed T cell populations.

Although it had been known for a long time that APC process and present antigens to T cells, it was not shown until relatively recently that small antigenic peptides could directly bind to MHC-encoded molecules (Babbitt et al., 1985, *Nature* 317:359; Townsend et al., 1986, *Cell* 44:959). However, it is believed that, normally, complex antigens are proteolytically processed into fragments inside the APC, and become physically associated with the MHC-encoded proteins intracellularly prior to

trafficking to the cell surface as complexes. Two distinct pathways for antigen presentation have been proposed (Braciale et al., 1987, *Immunol. Rev.* 98:95-114). It was thought that exogenous antigens were taken up by APC, processed and presented by the exogenous pathway to class II restricted CD4⁺ T cells, while the endogenous pathway processed intracellularly synthesized proteins, such as products of viral genes in virally-infected cells, for association with MHC class I proteins and presentation to CD8⁺ CTL. However, although the two pathways in antigen processing and presentation may still be correct in some respects, the distinction is blurred in light of recent findings that exogenously added antigens may also be presented to class I-restricted CTL (Moore et al., 1988, *Cell* 54:777).

The term "dendritic cells" (DC) refers to a diverse population of morphologically similar cell types found in a variety of lymphoid and non-lymphoid tissues (Steinman, 1991, *Ann. Rev. Immunol.* 9:271-296). These cells include lymphoid DC of the spleen, Langerhans cells of the epidermis, and veiled cells in the blood circulation. Although they are collectively classified as a group based on their morphology, high levels of surface MHC-class II expression, and absence of certain other surface markers expressed on T cells, B cells, monocytes, and natural killer cells, it is presently not known whether they derive from a common precursor or can all function as APC in the same manner. Further, since the vast majority of published reports have utilized DC isolated from the mouse spleen, results from these studies may not necessarily correlate with the function of DC obtained from other tissue types. (Inaba et al., 1997, *J. Exp. Med.* 166:182-194; Hengel et al., 1987, *J. Immunol.*, 139:4196-4202; Kaut et al., 1988, *J. Immunol.*, 140:3186-3193; Romani et al., 1989, *J. Exp. Med.* 169:1169-1178; Macatonia et al., 1989, *J. Exp. Med.* 169:1255-1264; Inaba et al., 1990, *J. Exp. Med.* 172:631-6640). For example, despite high levels of MHC-class II expression, mouse epidermal Langerhans cells, unlike splenic DC, are not active APC in mixed leucocyte reaction (MLR), unless cultured with granulocyte-macrophage colony stimulating factor (GM-CSF) (Witmer-Pock et al., 1987, *J. Exp. Med.* 166:1484-1498; Heufler et al., 1988, *J. Exp. Med.* 167:700-705). Most human Langerhans cells express the CD1 and CD4 markers, while blood DC do not. Additionally, it has not been established the extent to which the functional characteristics observed with mouse DC are applicable to human DC, especially the DC obtained from non-splenic tissues; in part, due to inherent differences between the

human and murine immune systems.

Recently, a few studies have described the isolation of human DC from the peripheral blood, which involves the use of sheep red blood cells and/or fetal calf serum (Young and Steinman, 1990, *J. Exp. Med.* 171:1315-1332; Freudenthal and Steinman, 1990, *Proc. Natl. Acad. Sci. USA* 87:7698-7702; Macatonia et al., 1989 *Immunol.* 67:285-289; Markowicz and Engleman, 1990, *J. Clin. Invest.* 85:955-961). Engleman et al. described a partial purification procedure of DC from human blood, which does not involve the use of sheep red blood cells and/or fetal calf serum, and showed that the partially purified human DC can, in fact, present exogenous antigens to naive T cells (PCT Publication WO 94/02156 dated February 3, 1994 at page 9, lines 5-32).

Recent studies have indicated that DCs are superior APCs as compared to other APCs such as macrophages and monocytes. First, the potent accessory cell function of DCs provides for an antigen presentation system for virtually any antigenic epitopes which T and B cells are capable of recognizing through their specific receptors. For example, Engleman et al. demonstrate that human DCs can present both complex protein antigens and small peptides to CD4⁺ T cells as well as to CD8⁺ CTL (PCT Publication WO 94/02156 dated February 3, 1994, Example 7, from page 29, line 10 to page 34, line 16). Engleman et al. also show that the *in vitro* priming effect of DCs does not require the addition of exogenous lymphokines, indicating that DCs produce all of the necessary signals in antigen presentation leading to the activation of T cells (PCT Publication WO 94/02156 dated February 3, 1994, from page 32, line 36 to page 33, line 2). More importantly, DCs can induce a primary CD4⁺ T cell-mediated proliferative response when similarly prepared monocytes can not induce such a response (PCT Publication WO 94/02156 dated February 3, 1994 at page 31, lines 23-30). Similarly, when DCs and monocytes were compared for their ability to present antigens for re-activating secondary T cell response, it was observed that DCs were capable of stimulating a stronger response than monocytes (PCT Publication WO 94/02156 dated February 3, 1994 at page 32, lines 12-16).

2.3. CHEMOKINES

Chemokines, or chemoattractant cytokines, are a subgroup of immune factors

that have been shown to mediate chemotactic and other pro-inflammatory phenomena (see, Schall, 1991, *Cytokine* 3:165-183). Chemokines are small molecules of approximately 70-80 residues in length and can generally be divided into two subgroups, α which have two N-terminal cysteines separated by a single amino acid (CxC) and β which have two adjacent cysteines at the N terminus (CC). RANTES, MIP-1 α and MIP-1 β are members of the β subgroup (reviewed by Horuk, R., 1994, *Trends Pharmacol. Sci.* 15:159-165; Murphy, P.M., 1994, *Annu. Rev. Immunol.* 12:593-633; Baggiolini et al. *Annu. Rev. Immunol.* 1997, 15:675-705).

MCP-1 has been shown to attract monocytes but not neutrophils. MCP-1, MCP-2, and MCP-3 share a pyroglutamate proline NH₂-terminal motif and are structurally closely related to each other and to eotaxin (56% to 71% amino acid sequence identity). MCP-1, MCP-2, and MCP-3 attract monocytes, CD4⁺ and CD8⁺ T lymphocytes (Loetscher et al. *FAESB J.* 1994, 8:1055-60), as well as basophil leukocytes. MCP-2, MCP-3, and MCP-4 (but not MCP-1) attracts eosinophil leukocytes. All four MCPs attract activated T lymphocytes, natural killer (NK) cells, and dendritic cells (see Baggiolini et al. *Annu. Rev. Immunol.* 1997, 15:675-705).

Eotaxin acts on eosinophils and is inactive on neutrophils and monocytes, but has weak-to-moderate chemotactic activity toward IL-2-conditioned T lymphocytes (see Baggiolini et al. *Annu. Rev. Immunol.* 1997, 15:675-705). Due to its preferential, powerful action on eosinophils and its occurrence in different species, eotaxin is considered to be an important chemokine in the pathophysiology of allergic conditions and asthma (See Baggiolini et al. *Annu. Rev. Immunol.* 1997, 15:675-705).

IP10 is a CXC chemokine attracts human monocytes, T lymphocytes, and NK cells, and Mig attracts tumor-infiltrating T lymphocytes. It has been suggested that IP10 and Mig may also be involved in the regulation of lymphocyte recruitment and the formation of the lymphoid infiltrates observed in autoimmune inflammatory lesions, delayed-type hypersensitivity, some viral infections, and certain tumors (Baggiolini et al. *Annu. Rev. Immunol.* 1997, 15:675-705).

SDF-1 (stromal cell-derived factor 1), including SDF-1 and SDF-1 β stimulates the proliferation of B cell progenitors, and attracts mature dendritic cells (Finkel et al. *Immunobiology* 1998, 198:490-500). Synthetic human SDF-1 stimulates monocytes, neutrophils, and peripheral blood lymphocytes, as is indicated by [Ca²⁺]_i changes and chemotaxis. SDF-1 is also a powerful HIV-suppressive factor (See Baggiolini et al.

Annu. Rev. Immunol. 1997, 15:675-705).

The amino terminus of the β chemokines RANTES, MCP-1, and MCP-3 has been implicated in the mediation of cell migration and inflammation induced by these chemokines. This involvement is suggested by the observation that the deletion of the amino terminal 8 residues of MCP-1, amino terminal 9 residues of MCP-3, and amino terminal 8 residues of RANTES and the addition of a methionine to the amino terminus of RANTES, antagonize the chemotaxis, calcium mobilization and/or enzyme release stimulated by their native counterparts (Gong et al., 1996, *J. Biol. Chem.* 271:10521-10527; Proudfoot et al., 1996 *J. Biol. Chem.* 271:2599-2603). Additionally, α chemokine-like chemotactic activity has been introduced into MCP-1 via a double mutation of Tyr 28 and Arg 30 to leucine and valine, respectively, indicating that internal regions of this protein also play a role in regulating chemotactic activity (Beall et al., 1992, *J. Biol. Chem.* 267:3455-3459).

The monomeric forms of all chemokines characterized thus far share significant structural homology, although the quaternary structures of α and β groups are distinct. While the monomeric structures of the β and α chemokines are very similar, the dimeric structures of the two groups are completely different. An additional chemokine, lymphotactin, which has only one N terminal cysteine has also been identified and may represent an additional subgroup (γ) of chemokines (Yoshida et al., 1995, *FEBS Lett.* 360:155-159; and Kelner et al., 1994, *Science* 266:1395-1399).

Receptors for chemokines belong to the large family of G-protein coupled, 7 transmembrane domain receptors (GCR's) (See, reviews by Horuk, R., 1994, *Trends Pharmacol. Sci.* 15:159-165; and Murphy, P.M., 1994, *Annu. Rev. Immunol.* 12:593-633). Competition binding and cross-desensitization studies have shown that chemokine receptors exhibit considerable promiscuity in ligand binding. Examples demonstrating the promiscuity among β chemokine receptors include: CCR-1, which binds RANTES and MIP-1 α (Neote et al., 1993, *Cell* 72:415-425), CCR-4, which binds RANTES, MIP-1 α , and MCP-1 (Power et al., 1995, *J. Biol. Chem.* 270:19495-19500), and CCR-5, which binds RANTES, MIP-1 α , and MIP-1 β (Alkhatib et al., 1996, *Science* 272:1955-1958 and Dragic et al., 1996, *Nature* 381:667-674). Erythrocytes possess a receptor (known as the Duffy antigen) which binds both α and β chemokines (Horuk et al., 1994, *J. Biol. Chem.* 269:17730-17733; Neote et al., 1994, *Blood* 84:44-52; and Neote et al., 1993, *J. Biol. Chem.* 268:12247-12249). Thus the sequence and

structural homologies evident among chemokines and their receptors allow some overlap in receptor-ligand interactions.

Godiska et al. identified and described the nucleic acid and amino acid sequences of an additional β chemokine designated macrophage derived chemokine (MDC) (PCT Publication WO 96/40923 dated December 19, 1996, and 1997, *J. Exp. Med.* 185:1595-1604). PCT publication WO 96/40923 further provides materials and methods for the recombinant production of the chemokine, the purified and isolated chemokine protein, and polypeptide analogues thereof. The PCT publication WO 96/40923 does not disclose that the human MDC has chemotactic activity upon DC. While Godiska et al. (1997, *J. Exp. Med.* 185:1595-1604) showed that, in a microchamber migration assay, monocyte-derived DC migrated toward the human MDC, the reference fails to teach that MDC can enhance an immune response to an antigen *in vivo*.

Chang et al. (1997, *J. Biol. Chem.* 272(40):25229-25237), isolated a stimulated T cell chemotactic protein (STCP-1) from an activated macrophage cDNA library. The nucleotide sequence of the STCP-1 is identical to that of the MDC isolated by Godiska et al. (PCT Publication WO 96/40923 dated December 19, 1996, and 1997, *J. Exp. Med.* 185:1595-1604). However, unlike the results observed by Godiska et al. (1997, *J. Exp. Med.* 185:1595-1604), Chang et al. (1997, *J. Biol. Chem.* 272(40):25229-25237) showed that although the STCP-1 acted as a mild chemoattractant for primary activated T lymphocytes and a potent chemoattractant for chronically activated T lymphocytes, the STCP-1 has no chemoattractant activity for monocytes, neutrophils, eosinophils and resting T lymphocytes. Chang et al. further showed that the STCP-1 does not induce Ca^{2+} mobilization in monocytes, dendritic cells, neutrophils, eosinophils, lipopolysaccharide-activated B lymphocytes, and freshly isolated resting T lymphocytes.

2.4. HIV VACCINES

Human immunodeficiency virus (HIV) induces a persistent and progressive infection leading, in the vast majority of cases, to the development of the acquired immunodeficiency syndrome (AIDS) (Barre-Sinoussi et al., 1983, *Science* 220:868-870; Gallo et al., 1984, *Science* 224:500-503). The HIV envelope surface glycoproteins are

synthesized as a single 160 kilodalton precursor protein which is cleaved by a cellular protease during viral budding into two glycoproteins, gp41 and gp120. gp41 is a transmembrane glycoprotein and gp120 is an extracellular glycoprotein which remains non-covalently associated with gp41, possibly in a trimeric or multimeric form (Hammerskjold, M. and Rekosh, D., 1989, *Biochem. Biophys. Acta* 989:269-280). The V3 loop of gp120 is the major determinant of sensitivity to chemokine inhibition of infection or replication (Cocchi et al., 1996, *Nature Medicine* 2:1244-1247; and Oravec et al., 1996, *J. Immunol.* 157:1329-1332).

Although considerable effort is being put into the design of effective therapeutics, currently no curative anti-retroviral drugs against AIDS exist. The HIV-1 envelope proteins (gp160, gp120, gp41) have been shown to be the major antigens for neutralizing anti-HIV antibodies present in AIDS patients (Barin et al., 1985, *Science* 228:1094-1096). Thus far, therefore, these proteins seem to be the most promising candidates to act as antigens for anti-HIV vaccine development. Several groups have begun to use various portions of gp160, gp120, and/or gp41 as immunogenic targets for the host immune system (see, for example, Ivanoff et al., U.S. Pat. No. 5,141,867; Saith et al., PCT publication WO 92/22654; Shafferman, A., PCT publication WO 91/09872; Formoso et al., PCT publication WO 90/07119). Therefore, methods to increase the efficacy of vaccines against HIV, especially vaccines using gp120 as the antigen, are needed.

Additionally a novel vaccine technology, designated genetic vaccination, nucleic acid vaccination or DNA vaccination, has been explored to induce immune responses *in vivo*. Injection of cDNA expression cassettes results in *in vivo* expression of the encoded proteins (Dubensky et al., 1984, *Proc. Natl. Acad. Sci. USA* 81:7529-7533; Raz et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:4523; Wolff et al., 1990, *Science* 247:1465-1468), with the concomitant development of specific cellular and humoral immune responses directed against the encoded antigen(s) (Wang et al., 1995, *Hum. Gene Ther.* 6:407-418; Ulmer et al., 1993, *Science* 259:1745-1749; Tang et al., 1992, *Nature* 356:152-154; Michel et al., 1995, *Proc. Natl. Acad. Sci. USA* 92:5307-5311; and Lowrie et al., 1994, *Vaccine* 12:1537-1540). Humoral and cellular responses have been induced to HIV-1 and SIV antigens through various applications of this technology in macaques (Wang et al., 1995, *Virology* 221:102-112; Wang et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:4156-4160; and Boyer et al., 1996, *J. Med.*

Primatol. 25:242-250) as well as mice (Wang et al., 1995, *Virology* 221:102-112; Lu et al., 1995, *Virology* 209:147-154; Haynes et al., 1994, *AIDS Res. Hum. Retroviruses* 10 (Suppl. 2):S43-S45; Okuda et al., 1995, *AIDS Res. Hum. Retroviruses* 11:933-943).

Recently, Lekutis et al. (1997, *J. Immunol.* 158:4471-4477), assessed the TH cell response elicited by an HIV-1 gp120 DNA vaccine in rhesus monkeys by isolation of gp120-specific, MHC class II-restricted CD4⁺ T cell lines from the vaccinated animals. Lekutis et al. showed that the isolated cell lines proliferated in response to APC in the presence of recombinant gp120, as well as to APC expressing HIV encoded env protein. Lekutis et al. further showed that these cell lines responded to env by secreting IFN- γ and IFN- α without appreciable IL-4 production. These results demonstrate that the animals exhibited a cellular immune response to the DNA vaccine.

Boyer et al. (1997, *Nature Medicine* 3:625-532), inoculated chimpanzees with an HIV-1 DNA vaccine encoding env, rev, and gag/pol, and found that the immunized animals developed specific cellular and humoral immune responses to these proteins. After challenging the immunized animals with a heterologous chimpanzee titrated stock of HIV-1 SF2, Boyer et al. further found, using a Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) assay, that those animals vaccinated with the DNA vaccine were protected against infection whereas the control animals were not so protected.

Kim et al., (1997 *J. Immunol.* 158:816-826), investigated the role of co-delivery of genes for IL-12 and GM-CSF along with DNA vaccine formulation for HIV-1 antigens env and gag/pol in mice. Kim et al. observed a dramatic increase in specific CTL response from the mice immunized with the HIV-1 DNA vaccine and IL-12. Kim et al. also observed that the co-delivery of IL-12 genes resulted in the reduction of specific antibody response, whereas the codelivery of GM-CSF genes resulted in the enhancement of specific antibody response. Kim et al. further observed that co-delivery of IL-12 gene with a HIV DNA vaccine results in splenomegaly (Kim et al. 1997, *J. Immunol.*, 158:816-826), which has been shown in mice to have toxic effects such as weight reduction or even death (Eng et al., 1995, *J. Exp. Med.* 181:1893; Stevensen et al., 1995, *J. Immunol.* 155:2545; and Orange et al., 1995, *J. Exp. Med.* 181:901).

Notwithstanding the recent developments of the HIV DNA vaccine, there still

exists a need for a method to enhance the efficacy of a vaccine, especially an HIV DNA vaccine. For instance, for efficacious vaccine against HIV-1 one preferably induces both cellular and humoral immune responses to control the infection (Boyer et al., 1997, *Nature Medicine* 3:625-532). The induction of both cellular and humoral immune response by the Berjer et al. method is still quite low because only one of the three immunized chimpanzees developed both cellular and humoral responses. Similarly, although co-delivery of an IL-12 encoding gene with a HIV DNA vaccine, as described in Kim et al. (1997, *J. Immun.* 158:816-826), may have enhanced the cellular immune response, this co-delivery also decreased the humoral response.

Citation of a reference hereinabove shall not be construed as an admission that such reference is prior art to the present invention.

3. SUMMARY OF THE INVENTION. SUMMARY OF THE INVENTION. . SUMMARY OF THE INVENTION

The present invention is based upon the ability of chemokines, such as MDC, Rantes, MIP-1d, MIP-1B, and I-309, to enhance the immune response to an antigen, particularly a vaccine. Accordingly, in a first aspect, the present invention provides a method for enhancing the efficacy of a vaccine, which method comprises administration to a subject of one or more purified chemokines, or biologically active fragments, analogues or derivatives thereof, either concurrently with one or more purified antigens against which an immune response is desired or within a time period either before or after administration of the antigens such that the immune response against the antigens is enhanced.

In a second aspect, the present invention provides a method to enhance the efficacy of a vaccine, which method comprises administration to a subject of a first set of one or more purified nucleic acids comprising one or more nucleotide sequences encoding one or more chemokines, or fragments, derivatives, analogues, and/or truncation isoforms thereof, and a second purified nucleic acid comprising a nucleotide sequence encoding one or more antigens against which an immune response is desired, such that, the one or more chemokine(s) and the antigen(s) are expressed in a coordinated manner upon introduction into a suitable cell. Alternatively, the nucleotide sequences encoding one or more chemokines, or

fragments, derivatives, and/or analogues thereof, and the antigens against which an immune response is desired are present on the same nucleic acid.

In a preferred embodiment, the invention provides a method to enhance the efficacy of an HIV vaccine.

In yet another aspect, the present invention provides a composition comprising an immunogenic amount of one or more purified antigens, an amount of one or more purified chemokines, or a fragments, derivatives, analogues and/or truncation isoforms thereof, effective to enhance the immune response to the antigen. In another aspect, the present invention provides a composition comprising a first set of one or more purified nucleic acids comprising one or more nucleotide sequences encoding one or more chemokines, fragments, derivatives analogues and or truncation isoforms thereof, and a second set of purified nucleic acids comprising one or more nucleotide sequences encoding one or more antigens against which an immune response is desired, such that, the chemokine(s) and the antigen are expressed in a coordinated manner upon introduction into a suitable cell. In a preferred embodiment, the antigen is an HIV antigen. In another preferred embodiment, the chemokine is selected from the group consisting of: Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EB11-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha

chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

4. DESCRIPTION OF FIGURES

Figures 1A and 1B. The nucleotide and amino acid sequences of MDC. 1A depicts the nucleotide sequence of MDC (SEQ ID NO:1), with the coding region indicated by the appearance of the amino acid sequence in the line below; and 1B depicts the amino acid of MDC (SEQ ID NO:2) from GenBank accession no. U83171 (Godiska et al., 1997, *J. Exp. Med.* 185:1595-1604).

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method for enhancing the efficacy of a vaccine in a subject comprising administering to the subject one or more purified antigens in conjunction with one or more purified chemokines, or more purified fragments, derivatives or analogues and/or truncation isoforms thereof.

While any chemokine may be employed according to the present invention, the chemokine is preferably selected from the following table:

| Chemokine Class | Chemokines | Abbreviations | Accession Number |
|-----------------|---|----------------|--------------------------|
| CC Chemokines | Macrophage-derived chemokine | MDC/STCP-1 | u83171; u83239 |
| | Monocyte chemotactic protein 1 | MCP-1 | x14768 |
| | Monocyte chemotactic protein 2 | MCP-2 | X99886 |
| | Monocyte chemotactic protein 3 | MCP-3 | x72308; s57464 |
| | Monocyte chemotactic protein 4 | MCP-4 | u46767 |
| | activated macrophage specific chemokine 1 | AMAC-1 | Y13710 |
| | Macrophage inflammatory protein 1 alpha | MIP-1 α | AF043339; X03754; D90144 |

| Chemokine Class | Chemokines | Abbreviations | Accession Number |
|------------------------------|--|-----------------------------------|--------------------------------|
| CC Chemokines (continued) | Macrophage inflammatory protein 1 beta | MIP-1 β | j04130; d90145 |
| | Macrophage inflammatory protein 1 gamma | MIP-1 γ | |
| | Macrophage inflammatory protein 1 delta | MIP-1 δ | AF031587 |
| | Macrophage inflammatory protein 2 alpha | MIP-2 α | AF043340 |
| | Macrophage inflammatory protein 3 alpha | MIP-3 α | u77035 |
| | Macrophage inflammatory protein 3 beta | MIP-3 β | u77180 |
| | Regulated upon activation, normal T cell expressed and secreted (and its variants) | RANTES | M21211 |
| | I-309 | | M57502 |
| | EBI1-ligand chemokine | ELC | AB000887 |
| | Pulmonary and activation regulated chemokine | PARC/DC-CK-1/MIP4 | AB000221 |
| | Liver and activation-regulated chemokine | LARC | D86955 |
| | Thymus and activation regulated chemokine | TARC | D43767 |
| | Eotaxin (and variants) | | D49372; Z69291; Z75669; Z75668 |
| | Human chemokine 1 | HCC1; NCC2 | Z49270; z49269 |
| | Human chemokine 2 | HCC2; NCC3, MIP-5, MIP-1 δ | Z70292 |
| | Human chemokine 3 | HCC3 | Z70293 |
| | IL-10-inducible chemokine | HCC4 | U91746 |
| | liver-expressed chemokine. | LEC; HCC4;NCC4 | AB007454 |
| | 6Ckine | | AF001979 |
| | Exodus 1 | | u64197 |
| | Exodus 2 | | U88320 |
| | Exodus 3 | | U88321 |
| | thymus-expressed chemokine | TECK | U86358 |
| | Secondary Lymphoid tissue chemokine | SLC | AB002409 |

| Chemokine Class | Chemokines | Abbreviations | Accession Number |
|------------------------------|---|--------------------------|------------------|
| CC Chemokines (continued) | Lymphocyte and Monocyte chemoattractant; Monotactin | LMC | AF055467 |
| | Activation-induced, chemokine-related molecule | ATAC | x86474 |
| | Myeloid progenitor inhibitory factor-1 | MPIF-1; MIP-3 or ckbeta8 | u85767 |
| | Myeloid progenitor inhibitory factor-2 | MPIF-2 | u85768 |
| | Stromal cell-derived factor 1 alpha | SDF-1 α ; PBSF | L36034 |
| CXC chemokines | Stromal cell-derived factor 1 beta | SDF-1 β ; PBSF | L36033 |
| | B-cell-attracting chemokine 1 | BLC | AJ002211 |
| | HuMIG | | x72755 s60728 |
| | H174 | | AF002985 |
| | Interferon-stimulated T-cell alpha chemoattractant | I-TAC | AF030514 |
| | Interleukin-8 | IL-8 | m17017; y00787 |
| | IP-10 | | X02530 |
| | platelet factor 4 | PF4 | M20901 |
| | growth-regulated gene-alpha | GRO- α | J03561 |
| | growth-regulated gene-beta | GRO- β | M36820 |
| | growth-regulated gene-gamma | GRO- γ | M36821 |
| | Neutrophil-activating protein 2 | NAP-2; CTAP-3 | M54995; M38441 |
| | ENA-78 | | L37036 |
| | granulocyte chemotactic protein 2 | GCP-2 | Y08770 |
| C-CHEMOKINES | LYMPHOTACTIN | SCM-1 | D63789 D63790 |
| CX3C-CHEMOKINES | Fractalkine/neurotactin | | U91835 U84487 |

The present invention also relates to the use of fragments, analogues and derivatives of the foregoing chemokines, as well as truncation isoforms of such chemokines which are known in the art.

The present invention also relates to therapeutic compositions comprising one or more chemokines, nucleic acids encoding one or more chemokines, derivatives, analogues, and/or truncation isoforms thereof, and nucleic acids encoding the same, that are effective to enhance the immune response of a subject to a vaccine.

In another preferred embodiment of the invention, nucleic acids comprising

nucleotide sequences encoding one or more chemokines or fragments or derivatives, including truncation isoforms, thereof, and encoding one or more antigens against which an immune response is desired, which coding sequences are operatively linked to gene regulatory sequences capable of directing the expression of the one or more chemokines and the one or more antigens upon introduction into a suitable cell, for example, but not limited to, the cell (of a subject), are administered to a subject such that the one or more chemokines, or fragments or derivatives, including truncation isoforms, thereof, and one or more antigens, are expressed in the subject.

For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections which follow.

5.1. METHODS AND COMPOSITIONS TO ENHANCE THE EFFICACY OF A VACCINE

The present invention provides methods for enhancing the efficacy of a vaccine in a subject, which methods comprise administering to a subject an immunogenic amount of one or more purified antigens against which an immune response is desired in the subject in conjunction with an amount of one or more purified chemokines, or fragments, derivatives, analogues and/or truncation isoforms thereof, effective to enhance the immune response against the antigen. In one aspect, the purified chemokine(s), or fragment(s), derivative(s), analogue(s) and/or truncation isoforms thereof, are administered to the subject concurrently with (e.g., in the same composition with) the purified antigen or antigens against which an immune response is desired. In another, aspect, the purified chemokine(s), or fragment(s), derivative(s), analogue(s) and/or truncation isoforms thereof, are administered either before or after the administration of one or more purified antigens against which immunity is desired in the subject, but is administered within such time that the chemokine(s) enhance the immune response to the one or more antigens. For example, but not by way of limitation, the purified chemokine(s) are administered during the time that the subject mounts an immune response against the administered one or more antigens, or, the purified MDC is administered within, for example, but not limited to, 30 minutes, 1 hour, 5 hours, 10 hours, 1 day, 2 days of (preferably, after) administration of the one or more purified antigens against which immunity is desired.

In a preferred embodiment, the present invention provides compositions comprising an immunogenic amount of one or more purified antigens and an amount of purified MDC, or one or more fragments, derivatives or analogues thereof, effective to enhance the immune response to said antigen and, preferably, the composition further comprises a pharmaceutically acceptable carrier.

A preferred chemokine for use in the methods and compositions of the present invention is any MDC protein, fragment or derivative thereof, that is capable of enhancing the efficacy of a vaccine (for example, but not limited to, as determined by the assays described in Section 5.4, *infra*). In one specific embodiment, the MDC is purified full length MDC, preferably full length MDC having the amino acid sequence of SEQ ID NO: 2 (Figure 1B). In another embodiment, the MDC is a purified protein, the amino acid sequence of which consists of amino acid numbers 2-69 of SEQ ID NO: 2 (Figure 1B). In another specific embodiment, the MDC is a purified protein, the amino acid sequence of which consists of amino acid numbers 3-69 of SEQ ID NO: 2 (Figure 1B). In still another specific embodiment, the MDC is a purified protein, the N-terminal amino acid sequence of which consists of the amino acid sequence Tyr-Gly-Ala-Asn-Met-Glu-Asp-Ser-Val-Cys-Cys-Arg-Asp-Tyr-Val-Arg-Tyr-Arg-Leu (portion of SEQ ID NO: 2). In yet another specific embodiment, the MDC is a purified protein, the N-terminal amino acid sequence of which consists of the amino acid sequence Pro-Tyr-Gly-Ala-Asn-Met-Glu-Asp-Ser-Val-Cys-Cys-Arg (portion of SEQ ID NO: 2). In yet another specific embodiment, the MDC is a purified derivative of a protein, the N-terminal amino acid sequence of which protein consists of the amino acid sequence Tyr-Gly-Ala-Asn-Met-Glu-Asp-Ser-Val-Cys-Cys-Arg-Asp-Tyr-Val-Arg-Tyr-Arg-Leu (SEQ ID NO:2), which derivative has activity to enhance the efficacy of the vaccine. In yet another specific embodiment, the MDC is a purified derivative of a protein, the N-terminal amino acid sequence of which protein consists of the amino acid sequence Pro-Tyr-Gly-Ala-Asn-Met-Glu-Asp-Ser-Val-Cys-Cys-Arg (SEQ ID NO:2), which derivative has activity to enhance the efficacy of the vaccine.

In yet another specific embodiment, the chemokine is a purified derivative of the protein, which derivative has one or more insertions of or substitutions with one or more non-classical amino acids relative to a corresponding wildtype chemokine, which derivative will enhance the efficacy of the vaccine. In yet another specific

embodiment, the chemokine is a purified derivative of the protein that has only one or more conservative substitutions in sequence relative a corresponding wildtype chemokine, which derivative will enhance the efficacy of the vaccine. The chemokines useful in the present invention may be derived from any suitable source and obtained by any method known in the art, for example but not limited to the methods described in Section 5.2 *infra*.

Preferably, the chemokine(s) are of the same species as the subject to which the vaccine is administered. In a preferred embodiment, one or more human chemokines are administered to a human subject, e.g., human MDC is administered to a human subject, alone or in combination with another chemokine.

The present invention also provides a method to enhance the efficacy of a vaccine in a subject, which method comprises administering to a subject a purified first nucleic acid comprising a nucleotide sequence encoding an antigen against which an immune response is desired in a subject and a purified second nucleic acid comprising a nucleotide sequence encoding one or more chemokines, or fragment(s), derivative(s) or analogue(s) thereof, where the expression of the encoded antigen(s) and chemokine(s), or fragment(s), derivative(s) or analogue(s) thereof, are under control of one or more appropriate gene regulatory elements (which regulatory elements can be any regulatory element known in the art, for example, but not limited to, those regulatory elements described in Section 5.2 *supra*), such that, upon introduction of said first and second nucleic acids into a suitable cell (e.g., a cell of the subject), the antigen and chemokine(s), or fragment(s), derivative(s) or analogue(s) thereof, are coordinately expressed, *i.e.*, are expressed either at the same time or within an appropriate time period (*i.e.*, sufficient for the chemokine(s) to enhance the immune response against the antigen relative to a corresponding immune response in the absence of the chemokine) and the antigen(s) are expressed in an immunogenic amount and the chemokine(s), or fragment(s), derivative(s) or analogue(s) thereof, are expressed in an amount sufficient to enhance the immune response against the antigen(s). In a specific embodiment, the nucleotide sequences encoding the chemokine(s) and the antigen are present on separate nucleic acids. In another embodiment, the nucleotide sequences encoding the chemokine(s) and the antigen(s) are present on the same nucleic acid.

The present invention also provides compositions to enhance the

efficacy of a vaccine in a subject, which compositions comprise a purified first nucleic acid comprising a nucleotide sequence encoding one or more antigen(s) and a purified second nucleic acid comprising a nucleotide sequence encoding one or more chemokines, or fragments or derivatives, including truncation isoforms, thereof, wherein the nucleotide sequences encoding the antigens and the chemokine(s) are operably linked to one or more gene regulatory elements such that, upon introduction of said first and second nucleic acids into a suitable cell (e.g., a cell of the subject), the antigen(s) and chemokine(s) are expressed in a coordinated manner and the antigen(s) are expressed in an immunogenic amount and the chemokine(s) are expressed in an amount effective to enhance the immune response against the antigen, relative to a corresponding immune response in the absence of such chemokine(s).

The present invention also provides compositions to enhance the efficacy of a vaccine in a subject, which compositions comprise a purified first set of one or more purified nucleic acids comprising one or more nucleotide sequences encoding one or more antigens and a purified second set of one or more purified nucleic acids comprising a nucleotide sequence encoding one or more chemokines, or fragments, analogues, derivatives, (including truncation isoforms) thereof, wherein the nucleotide sequence(s) encoding the antigen(s) and the chemokine(s) are operably linked to one or more gene regulatory elements such that, upon introduction of said first and second sets of nucleic acids into a suitable cell (e.g., a cell of the subject), the antigen(s) and chemokine(s) are expressed in a coordinated manner and the antigen(s) are expressed in an immunogenic amount and the chemokine(s) are expressed in an amount effective to enhance the immune response against the antigen, relative to a corresponding immune response in the absence of such chemokine(s).

The present invention also provides compositions to enhance the efficacy of a vaccine in a subject, which compositions comprise a purified nucleic acid comprising a first set of one or more nucleotide sequences encoding one or more antigens and a second set of one or more nucleotide sequence encoding one or more chemokines, or fragments, derivatives, or analogues thereof (including truncation isoforms), wherein the first and second sets of nucleotide sequences are operably linked to one or more gene regulatory elements such that, upon introduction into a suitable cell, the antigen(s) and the chemokine(s) are expressed in a coordinated manner and the antigen(s) are expressed in an immunogenic amount and the chemokine(s) are

expressed in an amount effective to enhance the immune response against the antigen(s).

Any nucleic acid comprising a nucleotide sequence encoding one or more chemokine proteins, or fragments or derivatives, thereof (including truncation isoforms), that are capable of enhancing the immune response to the antigen (for example, but not limited to, as determined by any of the assays described in Section 5.2., *infra*) can be used in the methods and compositions of the present invention.

In a preferred embodiment, the nucleotide sequence encodes MDC. In another embodiment, the MDC-encoding nucleotide consists of the nucleotide sequence of SEQ ID NO:1 (Figure 1A). In another specific embodiment, the method or composition of the invention uses a nucleic acid encoding an MDC derivative having deletional, insertional or substitutional mutations and combination thereof, which derivative has activity to enhance the immune response against an antigen in a subject.

Such compositions of nucleic acids encoding an antigen are often referred to as DNA vaccines.

Such DNA vaccines are produced by any method known in the art for constructing an expression plasmid vector containing the nucleotide sequences of the antigen(s) and/or chemokine(s) to be expressed which vector is suitable for expression of the encoded proteins in the subject or in cells recombinant for the expression vector, which cells are to be provided to the subject. Such expression vectors may contain various promoters, terminators and polyadenylation coding regions to control the expression of the encoded protein.

The DNA vaccine can be administered by any method known in the art for administration of DNA. The DNA vaccine may be delivered either directly, in which case the subject is directly exposed to the DNA vaccine such that the DNA enters and is expressed in cells of the subject, or indirectly, in which case, the DNA vaccine is first introduced into suitable cells by any method known in the art *in vitro*, then the cells containing the DNA vaccine are transplanted into the subject.

In a specific embodiment, the DNA vaccine is directly administered *in vivo*, where it is expressed to produce the encoded antigens and chemokine(s). This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or

other viral vector (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In a preferred embodiment, the nucleic acid of a DNA vaccine is injected into the muscle of the subject to be immunized.

Another approach is to introduce the nucleic acid of the DNA vaccine into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign nucleic acid into cells (see e.g., Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92 (1985)) and may be used in accordance with the present invention. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene.

Cells into which a DNA vaccine can be introduced for purposes of immunization encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

The resulting recombinant cells can be delivered to a subject by various

methods known in the art. In a preferred embodiment, the recombinant cells are injected, e.g., subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the patient. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The cells can also be encapsulated in a suitable vehicle and then implanted in the subject (see, e.g., Dionne et al. PCT Publication WO 92/19195, dated November 12, 1992). The amount of cells envisioned for use depends on the desired effect, subject state, etc., and can be determined by one skilled in the art.

By way of example, and not by way of limitation a DNA vaccine may be generated as described by Lekutis et al. for an HIV DNA vaccine (1997, *J. Immunol.* 158:4471-4477). Briefly, an expression vector is constructed with the promoter, enhancer and intron A of human cytomegalovirus (CMV) and the termination and polyadenylation sequences of bovine growth hormone in a plasmid backbone. Additionally, the nucleotide sequence for signal sequence of tissue plasminogen activator is either substituted for the signal sequence of the antigen, if the antigen has a signal sequence or is added onto the amino-terminus of the antigen, thereby eliminating the dependence on viral proteins for expression (e.g., in the case of gp120 expression, rev and env proteins are required unless the HIV-1 signal sequence is so substituted). The resulting formulation is then injected intra-muscularly.

Further examples of DNA vaccines are set forth in Boyer et al. (1996, *J. Med. Primatol.*, 25:242-250), which describes the construction of a plasmid encoding the HIV-1 gp160 envelope glycoprotein as well as the rev-tax region cloned into pMAMneoBlue vector (Clonetech, Inc., Palo Alto, CA), and a vector encoding the envelope glycoprotein and rev from HIV-1 strain MN under the control of the CMV promoter. Another vector which can be used in the present invention is as described in Boyer et al. (1997, *Nature Medicine* 3:526-532) and contains expression cassettes encoding the envelope and Rev proteins of HIV-1 strain MN, and encoding the Gag/Pol proteins of HIV-1 strain IIIB.

For the practice of the present invention, the nucleotide sequence for the one or more chemokines, or fragments, derivatives, or analogues thereof, can either be incorporated into the same expression vector containing the nucleotide sequence encoding the antigen in such a manner that the chemokine(s) are expressed. Alternatively, the nucleotide sequence encoding the chemokine(s), or fragment(s),

derivative(s) or analogue(s) thereof, can be cloned into a separate expression vector (e.g., as described above for the expression vector containing the sequences coding for antigen) and the expression vector that expresses the antigen(s) mixed with the expression vector that expresses the chemokine(s). The mixture of the two expression vectors can then be administered to the subject.

The methods and compositions of the present invention may be used as a vaccine in a subject in which immunity for the antigen(s) is desired. Such antigens can be any antigen known in the art to be useful in a vaccine formulation. The methods and compositions of the present invention can be used to enhance the efficacy of any vaccine known in the art. The vaccine of the present invention may be used to enhance an immune response to infectious agents and diseased or abnormal cells, such as but not limited to bacteria, parasites, fungi, viruses, tumors and cancers. The compositions of the invention may be used to either treat or prevent a disease or disorder amenable to treatment or prevention by generating an immune response to the antigen provided in the composition. In one preferred embodiment, the antigen(s) are proteins, fragments or derivatives, including truncation isoforms, thereof, encoded by any genes of the HIV genome including the *env*, *gag*, *pol*, *nef*, *vif*, *rev*, and *tat* genes. In a more preferred embodiment, the antigen is an HIV-associated gp120 protein.

The methods and compositions of the present invention may be used to elicit a humoral and/or a cell-mediated response against the antigen(s) of the vaccine in a subject. In one specific embodiment, the methods and compositions elicit a humoral response against the administered antigen in a subject. In another specific embodiment, the methods and compositions elicit a cell-mediated response against the administered antigen in a subject. In a preferred embodiment, the methods and compositions elicit both a humoral and a cell-mediated response.

The subjects to which the present invention is applicable may be any mammalian or vertebrate species, which include, but are not limited to, cows, horses, sheep, pigs, fowl (e.g., chickens), goats, cats, dogs, hamsters, mice and rats, monkeys, rabbits, chimpanzees, and humans. In a preferred embodiment, the subject is a human. The compositions and methods of the invention can be used to either prevent a disease or disorder, or to treat a particular disease or disorder, where an immune response against a particular antigen or antigens is effective to treat or prevent the

disease or disorder. Such diseases and disorders include, but are not limited to, viral infections, such as HIV, CMV, hepatitis, herpes virus, measles, etc, bacterial infections, fungal and parasitic infections, cancers, and any other disease or disorder amenable to treatment or prevention by eliciting an immune response against a particular antigen or antigens. In another preferred embodiment, the subject is infected or at risk of being infected with HIV virus.

In another preferred embodiment the invention provides methods and compositions to enhance the efficacy of an HIV vaccine, such a vaccine can be administered to either prevent or treat HIV.

5.2. CHEMOKINE GENES AND PROTEINS

Chemokine proteins and nucleic acids can be obtained by any method known in the art. Chemokine nucleotide and amino acid sequences are available in public databases such as Genbank and are also published in various references known to those of skill in the art. The gene bank accession numbers for the preferred chemokines of the present invention are provided in Table I, in Section 5 above. The ensuing discussion uses MDC by way of example, but applies equally to other chemokines as well.

The MDC nucleotide and amino acid sequences for, *inter alia*, human, are available in the public databases (e.g. Genbank accession No. U83171) also published in Godiska et al., 1997, *J. Exp. Med.* 185:1595-1604. The nucleotide sequence and the amino acid sequence for the human MDC are provided in Figures 1A and B (SEQ ID NOS:1 and 2, respectively).

Chemokines used herein include, but are not limited to, chemokines from mice, hamsters, dogs, cats, monkeys, rabbits, chimpanzees, and human. In one preferred embodiment, the chemokine is of human origin.

Any vertebrate cell potentially can serve as the nucleic acid source for the isolation of chemokine nucleic acids. The nucleic acid sequences encoding the chemokine(s) can be isolated from vertebrate, mammalian, human, porcine, bovine, feline, avian, equine, canine, as well as additional primate sources, etc. The DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a

DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (see, for example, Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II.) Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences. Whatever the source, the gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from cDNA, cDNA is generated from totally cellular RNA or mRNA by methods that are well known in the art. The gene may also be obtained from genomic DNA, where DNA fragments are generated (e.g. using restriction enzymes or by mechanical shearing), some of which will encode the desired gene. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing all or a portion of the chemokine gene may be accomplished in a number of ways.

A preferred method for isolating a chemokine gene is by the polymerase chain reaction (PCR), which can be used to amplify the desired chemokine sequence in a genomic or cDNA library or from genomic DNA or cDNA that has not been incorporated into a library. Oligonucleotide primers which would hybridize to chemokine sequences can be used as primers in PCR.

Additionally, a portion of the chemokine (of any species) gene or its specific RNA, or a fragment thereof, can be purified (or an oligonucleotide synthesized) and labeled, the generated DNA fragments may be screened by nucleic acid hybridization to the labeled probe (Benton, W. and Davis, R., 1977, *Science* 196:180; Grunstein, M. And Hogness, D., 1975, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961). Those DNA fragments with substantial homology to the probe will hybridize. Chemokine nucleic acids can be also identified and isolated by expression cloning using, for example, anti-chemokine antibodies for selection.

Alternatives to obtaining the chemokine DNA by cloning or amplification

include, but are not limited to, chemically synthesizing the gene sequence itself from the known chemokine sequence or making cDNA to the mRNA which encodes the chemokine protein. Other methods are possible and within the scope of the invention. Once a clone has been obtained, its identity can be confirmed by nucleic acid sequencing (by any method well known in the art) and comparison to known chemokine sequences. DNA sequence analysis can be performed by any techniques known in the art, including but not limited to the method of Maxam and Gilbert (1980, *Meth. Enzymol.* 65:499-560), the Sanger dideoxy method (Sanger, F., et al., 1977, *Proc. Natl. Acad. Sci. U.S.A.* 74:5463), the use of T7 DNA polymerase (Tabor and Richardson, U.S. Patent No. 4,795,699), use of an automated DNA sequencer (e.g., Applied Biosystems, Foster City, CA) or the method described in PCT Publication WO 97/ 15690.

Nucleic acids which are hybridizable to a chemokine nucleic acid, or to a nucleic acid encoding a chemokine derivative can be isolated, by nucleic acid hybridization under conditions of low, high, or moderate stringency (see also Shilo and Weinberg, 1981, *Proc. Natl. Acad. Sci. USA* 78:6789-6792). For example, the nucleic acid of SEQ ID No: 1 is hybridizable to an MDC nucleic acid.

Chemokine proteins and derivatives, analogs and fragments of chemokine proteins can be obtained by any method known in the art, including but not limited to recombinant expression methods, purification from natural sources, and chemical synthesis.

For example, chemokines can be obtained by recombinant protein expression techniques. For recombinant expression, the chemokine gene or portion thereof is inserted into an appropriate cloning vector for expression in a particular host cell. A large number of vector-host systems known in the art may be used. Possible vectors include, but are not limited to, plasmids or modified viruses, but the vector system must be compatible with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene). The insertion into a cloning vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary cohesive termini. However, if the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site

desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative method, the cleaved vector and chemokine gene may be modified by homopolymeric tailing. Recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, etc., so that many copies of the gene sequence are generated.

In an alternative method, the desired gene may be identified and isolated after insertion into a suitable cloning vector in a "shot gun" approach. Enrichment for the desired gene, for example, by size fractionation, can be done before insertion into the cloning vector.

In specific embodiments, transformation of host cells with recombinant DNA molecules that incorporate the isolated chemokine gene, cDNA, or synthesized DNA sequence enables generation of multiple copies of the gene. Thus, the gene may be obtained in large quantities by growing transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

The nucleotide sequence coding for a chemokine protein or a functionally active analog or fragment or other derivative thereof, can be inserted into an appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The necessary transcriptional and translational signals can also be supplied by the native chemokine gene and/or its flanking regions. A variety of host-vector systems may be utilized to express the protein-coding sequence. These include but are not limited to mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used.

Any of the methods previously described for the insertion of DNA fragments into a vector may be used to construct expression vectors containing a chimeric gene consisting of appropriate transcriptional/translational control signals and the protein

coding sequences. These methods may include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of nucleic acid sequence encoding a chemokine protein or peptide fragment may be regulated by a second nucleic acid sequence so that the chemokine protein or peptide is expressed in a host transformed with the recombinant DNA molecule. For example, expression of a chemokine protein may be controlled by any promoter/enhancer element known in the art. Promoters which may be used to control chemokine expression include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Kamaroff, et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731), or the tac promoter (DeBoer, et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25); see also "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242:74-94; promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171), beta-globin gene control region

which is active in myeloid cells (Mogram et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94), myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712), myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, Nature 314:283-286), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

For example, a vector can be used that comprises a promoter operably linked to an chemokine-encoding nucleic acid, one or more origins of replication, and, optionally, one or more selectable markers (e.g., an antibiotic resistance gene).

In a specific embodiment, an expression construct is made by subcloning a chemokine coding sequence into the *EcoRI* restriction site of each of the three pGEX vectors (Glutathione S-Transferase expression vectors; Smith and Johnson, 1988, Gene 7:31-40). This allows for the expression of the chemokine protein product from the subclone in the correct reading frame.

Expression vectors containing chemokine gene inserts can be identified by three general approaches: (a) nucleic acid hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the first approach, the presence of a chemokine gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted chemokine gene. In the second approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of a chemokine gene in the vector. For example, if the chemokine gene is inserted within the marker gene sequence of the vector, recombinants containing the chemokine insert can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the product expressed by the recombinant. Such assays can be based, for example, on the physical or functional properties of the chemokine protein in *in vitro* assay systems, e.g., binding with anti-chemokine antibody or the chemokine's receptor.

Once a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host

system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (e.g., lambda), and plasmid and cosmid DNA vectors, to name but a few.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered protein may be controlled. Furthermore, different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, phosphorylation of proteins). Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce an unglycosylated core protein product. Expression in yeast will produce a glycosylated product. Expression in mammalian cells can be used to ensure "native" glycosylation of a heterologous protein. Furthermore, different vector/host expression systems may effect processing reactions to different extents.

In other specific embodiments, the chemokine protein(s), fragment(s), analogue(s), or derivative(s) may be expressed as a fusion, or chimeric protein product (comprising the protein, fragment, analog, or derivative joined via a peptide bond to a heterologous protein sequence (of a different protein)). Such a chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the chimeric product by methods commonly known in the art. Alternatively, such a chimeric product may be made by protein synthetic techniques, e.g., by use of a peptide synthesizer. In a specific embodiment, a chimeric protein containing all or a portion of the chemokine is joined via a peptide bond to all or a portion of an antigen against which immunity is desired.

Both cDNA and genomic sequences can be cloned and expressed.

The chemokine protein(s) may also be isolated and purified by standard methods including chromatography (e.g., ion exchange, affinity, and sizing column

chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. The functional properties may be evaluated using any suitable assay (see Section 5.5). Alternatively, the protein can be synthesized by standard chemical methods known in the art (e.g., see Hunkapiller, M., et al., 1984, *Nature* 310:105-111). The chemokine-encoding nucleic acid sequence(s) can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions. Any technique for mutagenesis known in the art can be used, including, but not limited to, *in vitro* site-directed mutagenesis (Hutchinson et al., 1978, *J. Biol. Chem.* 253:6551), use of TAB linkers (Pharmacia), mutation-containing PCR primers, etc.

The experimentation involved in mutagenesis consists primarily of site-directed mutagenesis followed by phenotypic testing of the altered gene product. Some of the more commonly employed site-directed mutagenesis protocols take advantage of vectors that can provide single stranded as well as double stranded DNA, as needed. Generally, the mutagenesis protocol with such vectors is as follows. A mutagenic primer, i.e., a primer complementary to the sequence to be changed, but consisting of one or a small number of altered, added, or deleted bases, is synthesized. The primer is extended *in vitro* by a DNA polymerase and, after some additional manipulations, the now double-stranded DNA is transfected into bacterial cells. Next, by a variety of methods, the desired mutated DNA is identified, and the desired protein is purified from clones containing the mutated sequence. For longer sequences, additional cloning steps are often required because long inserts (longer than 2 kilobases) are unstable in those vectors. Protocols are known to those skilled in the art and kits for site-directed mutagenesis are widely available from biotechnology supply companies, for example from Amersham Life Science, Inc. (Arlington Heights, IL) and Stratagene Cloning Systems (La Jolla, CA).

In other specific embodiments, the chemokine derivative(s) or analogue(s) may be expressed as a fusion, or chimeric protein product (comprising the protein, fragment, analogue, or derivative joined via a peptide bond to a heterologous protein sequence (of a different protein)). Such a chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the chimeric product by methods commonly known in the art.

In addition, chemokine proteins, derivatives (including fragments and chimeric proteins), and analogues can be chemically synthesized. See, e.g., Clark-Lewis et al., 1991, *Biochem.* 30:3128-3135 and Merrifield, 1963, *J. Amer. Chem. Soc.* 85:2149-2156. For example, chemokines, derivatives and analogues can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography (e.g., see Creighton, 1983, *Proteins, Structures and Molecular Principles*, W.H. Freeman and Co., N.Y., pp. 50-60). Chemokines, derivatives and analogues that are proteins can also be synthesized by use of a peptide synthesizer. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, 1983, *Proteins, Structures and Molecular Principles*, W.H. Freeman and Co., N.Y., pp. 34-49).

The chemokine proteins, derivatives, or analogues of the invention may be synthesized in their entirety by the sequential addition of amino acid residues or alternatively as fragment subcomponents which may be combined using techniques well known in the art, such as, for example, fragment condensation (Shin et al., 1992, *Biosci. Biotech. Biochem.* 56:404-408; Nyfeler et al., 1992, *Peptides*, Proc. 12th Amer. Pep. Soc., Smith and Rivier (eds), Leiden, pp 661-663); and Nokihara et al., 1990, *Protein Research Foundation*, Yanaihara (ed), Osaka, pp 315-320).

In a less preferred embodiment, chemokine derivatives can be obtained by proteolysis of the protein followed by purification using standard methods such as those described above (e.g., immunoaffinity purification).

In another alternate embodiment, native chemokine proteins can be purified from natural sources, by standard methods such as those described above (e.g., immunoaffinity purification).

5.3. COMPOSITION FORMULATIONS AND METHODS OF ADMINISTRATION

The composition formulations of the invention comprise an effective immunizing amount of an immunologically active ingredient, i.e., one or more antigens, and an amount of one or more chemokine(s), or fragment(s) or derivative thereof, effective to enhance the immune response against the antigen in a subject, and a pharmaceutically acceptable carrier or excipient. In a specific embodiment, the

chemokines are selected from the group consisting of Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EBI1-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

Pharmaceutically acceptable carriers or excipients are well known in the art and include but are not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, sterile isotonic aqueous buffer, and combinations thereof. One example of such an acceptable carrier is a physiologically balanced culture medium containing one or more stabilizing agents such as stabilized, hydrolyzed proteins, lactose, etc. The carrier is preferably sterile. The formulation should suit the mode of administration.

In addition, if desired, the vaccine or composition preparation may also include minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine or composition. Suitable adjuvants may include, but are not limited to: mineral gels,

e.g., aluminum hydroxide; surface active substances such as lysolecithin, pluronic polyols; polyanions; peptides; oil emulsions; alum, MDP, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine, and N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine. The effectiveness of an adjuvant may be determined by comparing the induction of antibodies directed against a MDC-containing composition in the presence and in the absence of various adjuvants.

In instances where the recombinant antigen is a hapten, *i.e.*, a molecule that is antigenic in that it can react selectively with cognate antibodies, but not immunogenic in that it cannot elicit an immune response, the hapten may be covalently bound to a carrier or immunogenic molecule; for instance, a large protein such as serum albumin will confer immunogenicity to the hapten coupled to it. The hapten-carrier may be formulated for use as a vaccine.

The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc.

The chemokine(s), or fragment(s) or derivative(s) thereof, and/or the antigen(s) may be formulated into the composition as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids, such as, for example, hydrochloric or phosphoric acids, or organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with free carboxyl groups may also be derived from inorganic bases, such as, for example, sodium potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

The vaccines of the invention may be multivalent or univalent. Multivalent vaccines are made from recombinant viruses that direct the expression of more than one antigen.

An effective dose (immunizing amount) is that amount sufficient to produce an immune response to the antigen(s) in the host to which the vaccine preparation is administered. The precise dose of the composition to be employed in the formulation will depend on the route of administration, and the nature of the subject to be

immunized, and should be decided by the practitioner according to standard clinical techniques. Effective doses of the vaccines or compositions of the present invention may also be extrapolated from dose-response curves derived from animal model test systems.

The invention also provides a pharmaceutical pack or kit comprising one or more containers comprising one or more of the ingredients of the composition formulations of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is administered by injection, an ampoule of sterile diluent can be provided so that the ingredients may be mixed prior to administration.

In a specific embodiment, a lyophilized immunologically active ingredient and one or more chemokine polypeptide(s) of the invention are provided in a first container; a second container comprises diluent consisting of an aqueous solution of 50% glycerin, 0.25% phenol, and an antiseptic (e.g., 0.005% brilliant green).

Many methods may be used to introduce the composition formulations of the invention; these include but are not limited to oral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal routes, and via scarification (scratching through the top layers of skin, e.g., using a bifurcated needle).

The DNA vaccines of the invention can be administered by any method known in the art for delivery of DNA to subject (for example, as described in Section 5.3 supra)

5.4. DETERMINATION OF COMPOSITION EFFICACY

The activity of one or more chemokines, or a fragment, derivative or analogue thereof, to enhance immune response to an antigen can be determined by monitoring the immune response in test animals following immunization with a composition containing the chemokine(s) and an antigen and comparing the response to that following immunization with the antigen in the absence of the chemokine(s). Generation of a humoral (antibody) response and/or cell-mediated immunity, may be taken as an indication of an immune response. Test animals may include mice, hamsters, dogs, cats, monkeys, rabbits, chimpanzees, etc., and eventually human subjects. Assays for humoral and cell-mediated immunity are well known in the art.

Methods of introducing the composition may include oral, intracerebral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal or any other standard routes of immunization. The immune response of the test subjects can be analyzed by various approaches well known in the art, such as but not limited to: testing the reactivity of the resultant immune serum to the antigen of the chemokine-containing vaccine, as assayed by known techniques, e.g., immunosorbant assay (ELISA), immunoblots, radioimmunoprecipitations, etc.

As one example of suitable animal testing, a composition of the present invention may be tested in mice for the ability to enhance an antibody response to an antigen (using for example, but not limited to, the method as described in Section 6, *infra*) and the delayed-type hypersensitivity (DTH) response (also described in Section 6 *infra*), measured by an increase in footpad swelling after inoculation in the footpad of the test animal, as compared to the measurements in animals administered the antigen in a composition not containing chemokine. For example, as test animals BALB/c mice may be used. The test group each receives an inoculation with fixed amount of antigen and varying amount of one or more chemokines. The control group receives an inoculation of comparable amount of antigen alone.

Serum samples may be drawn from the mice after the final inoculation (for example every one or two weeks after inoculation), and serum is analyzed for antibodies against the antigen using known methods in the art, e.g., using an ELISA. DTH responses to the antigen may be measured after the final inoculation (e.g. within 1-7 days). An increase in the serum titer of antibodies recognizing the antigen and/or

an increase in footpad swelling in the animals receiving the antigen-compositions containing the chemokine(s) as compared to the serum titer of antibodies against the antigen and/or the footpad swelling in the animals receiving the antigen composition not containing the chemokine(s), indicates that the chemokine(s) enhance the immune response to antigen. An increase in the serum titer of antibodies recognizing the antigen and/or an increase in footpad swelling in the animals receiving the antigen-compositions containing the chemokines as compared to the serum titer of antibodies against the antigen and/or the footpad swelling in the animals receiving the antigen composition not containing chemokine(s), indicates that the chemokine(s) enhances the immune response to antigen. An increase in the serum titer of antibodies recognizing the antigen and/or an increase in footpad swelling in the animals receiving the antigen-compositions containing MDC as compared to the serum titer of antibodies against the antigen and/or the footpad swelling in the animals receiving the antigen composition not containing MDC, indicates that the MDC enhances the immune response to antigen. An increase in the serum titer of antibodies recognizing the antigen and/or an increase in footpad swelling in the animals receiving the antigen-compositions containing MDC as compared to the serum titer of antibodies against the antigen and/or the footpad swelling in the animals receiving the antigen composition not containing MDC, indicates that the MDC enhances the immune response to antigen.

6. EXAMPLE: IMMUNIZATION WITH MDC-CONTAINING COMPOSITION

The following experiment illustrates the evaluation of whether MDC will act as an adjuvant for a protein antigen and enhance the efficacy of a vaccine. However, it will be appreciated that the description applies equally to other chemokines and combinations of chemokines.

6.1. MATERIALS AND METHODS

6.1.1. ANIMALS AND REAGENTS

BALB/c mice are purchased from Harlan-Sprague-Dawley (Indianapolis, IN).

Human MDC (hMDC) was obtained from CD8⁺ T cell clones immortalized *in vitro* prepared as previously described (Markham et al., 1983 *Int. J. Cancer* 31:413; Markham et al. 1984, *Int. J. Cancer* 33:13). One such immortalized CD8⁺ T cell clone, F3b Clone 19, was adapted to growth in serum-free medium by the following procedure and used for further studies. F3b Clone 19 cells were grown in complete medium containing rIL-2 (16 ng/ml) at 37°C in a CO₂ incubator. After expanding the culture to 200 ml, the cells were pelleted and resuspended in RPMI medium containing HB101 (Irvine Scientific) supplemented with 16 ng/ml of rIL-2, 1% glutamine and 1% penicillin/streptomycin. The cells were grown to full confluence and the medium harvested by centrifugation at 670 x g for 10 minutes.

Human MDC (hMDC) was purified from F3b Clone 19 as described in Pal et al., 1997, *Science* 278:695-698. Briefly, the cell free culture supernatant from F3b Clone 19 was clarified by high speed centrifugation and fractionated by heparin affinity chromatography, taking advantage of the heparin binding characteristics of chemokines (Witt and Lander, 1994, *Current Biology* 4:394; Proost et al., 1996, *Method: A Companion to Methods in Enzymology* 10:82). Culture supernatant (1200 ml) from F3b Clone 19, grown to high cell density in serum-free medium supplemented with rIL-2 was clarified by high speed centrifugation (100,000 x g for 60 minutes at 4°C) and applied to a 5 ml HiTrap heparin affinity FPLC column (Pharmacia) equilibrated in 10 mM Tris-HCl, pH 7.6 containing 0.1 M NaCl (column buffer). The column was then washed extensively with column buffer and the bound proteins eluted from the column with 10 mM Tris-HCl, pH 7.6 containing 2.0 M NaCl at a flow rate of 0.5 to 1 ml/minute. Virtually all of the HIV suppressive activity effective against primary NSI and SI isolates and HIV-1_{IIIb} was recovered in the column eluate (data not shown). The heparin affinity column eluate was brought to pH 2.0 by addition of trifluoroacetic acid (TFA) and subjected to reversed phase HPLC on a PEEK C-18 column (Waters Instruments) equilibrated in H₂O containing 0.1 % TFA. Proteins bound to the column were eluted with a 5 minute linear gradient of aqueous acetonitrile (0 to 35 %) containing 0.1% TFA. After 10 minutes at 35% acetonitrile, the column was further developed with a 60 minute linear gradient of 35-70% aqueous acetonitrile in TFA. The flow rate was maintained at 0.5 to 1 ml/minute. The fractions obtained were then tested for suppressor activity in the acute infectivity assay using HIV-1_{IIIb}. Active fractions were pooled, diluted twofold in H₂O with 0.1 % TFA

and reapplied to the column. The column was then developed with a 30 minute linear aqueous acetonitrile gradient (0-60%) containing 0.1% TFA at a flow rate of 0.5 to 1 ml/minute. The fractions obtained were assayed as above. Active fractions were pooled, diluted with H₂O/0.1 % TFA and fractionated under the same conditions to obtain a single protein peak. The fraction corresponding to the peak and flanking fractions were tested in the infectivity assay to verify that suppressor activity was cofractionated with the protein.

Suppressive activity against HIV-1_{IIIB} in the absence of cytotoxic effects consistently copurified with a single protein peak that appeared as a homogeneous 8 kDa band when analyzed by SDS-polyacrylamide gel electrophoresis. This protein was not reactive in ELISAs for RANTES, MIP-1 α or MIP-1 β (R&D Systems).

Recombinant gp120 protein derived from HIV-1 IIIB isolate is purchased from Intracel (Foster City, CA).

6.1.2 IMMUNIZATION OF MICE

The hMDC and the gp120 is resuspended in a total volume of 50 μ l of phosphate-buffered saline (PBS). Mice are divided into 5 groups with 3-4 mice in each group. Groups 1-4 are inoculated with 10 μ g gp120 and 0.3 μ g, 0.1 μ g, 0.03 μ g, and 0.01 μ g of hMDC, respectively. As a control, group 5 is inoculated with 10 μ g of gp120 in the absence of hMDC. For primary inoculation, each group of mice is inoculated with 10 μ l of the hMDC and gp120 solution via footpad. Two to three weeks after the primary inoculation, each mouse is given the same dose of hMDC/gp120 that is used in primary inoculation.

6.1.3 ELISA ASSAY

Serum samples are collected one week after the second inoculation via tail vein bleed. gp120 serum responses are measured using standard gp120 antibody ELISA assays.

6.1.4 DTH ASSAY

The delayed-type hypersensitivity (DTH) response is measured from 1-7 days after the second inoculation. A caliper is to be used to measure footpad swelling.

6.2. RESULTS

Mice inoculated with hMDC/gp120 are expected to have greater serum antibody and DTH responses than mice inoculated with gp120 alone. The improved responses will be reflected in either increased titers of serum antibody responses or increased footpad swelling. A dose response effect is expected - increasing the dose of hMDC used is expected to cause a corresponding improvement in the serum and DTH gp120-specific responses.

7. EXAMPLE: OTHER CHEMOKINES AND COMBINATIONS OF CHEMOKINES

The foregoing experiments can be repeated using other chemokines and combinations of chemokines. For example, the experiments are preferably repeated using one or more chemokines selected from the group consisting of: Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EB11-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine., 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

Various references are cited herein, the disclosures of which are incorporated by reference in their entireties.

THE CLAIMS:

1. A method to enhance the efficacy of a vaccine in a subject comprising administering to the subject an immunogenic amount of one or more purified antigens against which an immune response is desired in the subject and an amount of one or more chemokines, or purified fragments or derivatives thereof, effective to enhance the efficacy of said vaccine.
2. The method of claim 1, wherein the one or more chemokines are selected from a chemokine class selected from the group consisting of: CC, CXC, C-C and CX3C.
3. The method of claim 1, wherein the one or more chemokines are selected from the group consisting of: Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EBI1-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-

regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

4. The method of claim 1, wherein the one or more chemokines are selected from the group consisting of: MDC, SDF-1, BLC, and MCP-1.
5. The method of claim 1 wherein the fragment(s) or derivative(s) are truncation isoforms.
6. The method of claim 1, wherein the one or more chemokines include MDC comprising the amino acid sequence of SEQ ID NO: 2.
7. The method of claim 1, wherein the one or more chemokine fragment includes an MDC fragment selected from the group consisting of amino acid numbers 2-69, 3-69, 5-69, 7-69 and 9-69 of SEQ ID NO: 2.
8. The method of claim 1, wherein the one or more chemokine fragment includes an MDC fragment selected from the group consisting of amino acid numbers 2-69, 3-69, 5-69, 7-69 and 9-69 of SEQ ID NO: 2., which derivative has activity to enhance the efficacy of the vaccine.
9. The method of claim 1, wherein the one or more chemokine derivatives has one or more insertions or substitutions with one or more non-classical amino acids relative to a corresponding wildtype chemokine, which derivative has activity to enhance the efficacy of the vaccine.
10. The method of claim 1, including a chemokine derivative having one or more conservative substitutions in sequence relative a wildtype MDC, which derivative has activity to enhance the efficacy of the vaccine.
11. The method of claim 1, wherein the one or more chemokines include a human chemokine.

12. The method of claim 1, wherein the purified chemokine(s) or purified fragment(s) or derivative(s) thereof is/are administered concurrently with the purified antigen(s).
13. The method of claim 1 wherein the purified chemokine(s) or purified fragment(s) or derivative(s) thereof, are administered within a time period before or after administration of the purified antigen, which time period permits the purified MDC or purified fragment or derivative thereof MDC to enhance the efficacy of the vaccine.
14. The method of claim 1, wherein the antigen is an HIV antigen.
15. The method of claim 14, wherein the HIV antigen is HIV-associated gp120 protein.
16. The method of claim 1, wherein the subject is a human.
17. The method of claim 1, wherein the subject is infected or at risk of being infected with HIV virus.
18. The method of claim 1, wherein the vaccine elicits a humoral response against the antigen in the subject.
19. The method of claim 1, wherein the vaccine elicits a cell-mediated response against the antigen in the subject.
20. The method of claim 1, wherein the vaccine elicits both a humoral and a cell-mediated response against the antigen in the subject.
21. The method of claim 1, wherein the vaccine further comprises pharmaceutically acceptable excipient, auxiliary substance, adjuvant, wetting or emulsifying agent, or pH buffering agent.

22. A method to enhance the efficacy of a vaccine in a subject comprising administering to the subject a first amount of a first set of one or more purified nucleotide sequences encoding one or more antigens against which an immune response is desired in the subject and a second second set of one or more purified nucleic acids, each comprising a nucleotide sequence encoding one or more chemokines, or fragments or derivatives thereof, wherein the antigen(s) and the chemokine(s) are expressed in a coordinated manner upon introduction into a suitable cell, said first amount is immunogenic and said second amount is effective in enhancing the efficacy of the vaccine.
23. The method of claim 22, wherein the one or more chemokines are selected from a chemokine class selected from the group consisting of: CC, CXC, C-C and CX3C.
24. The method of claim 22, wherein the one or more chemokines are selected from the group consisting of: Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EBI1-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine

- 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.
25. The method of claim 22, wherein the one or more chemokines are selected from the group consisting of: MDC, SDF-1, BLC, and MCP-1.
26. The method of claim 22 wherein the fragment(s) or derivative(s) are truncation isoforms.
27. The method of claim 22, wherein the nucleotide sequence encoding one or more chemokines comprises the nucleotide sequence of SEQ ID NO:1.
28. The method of claim 22, wherein one or more of the chemokine derivative(s) have deletional, insertional or substitutional mutations and combination thereof, which derivative has activity to enhance the efficacy of the vaccine.
29. The method of claim 22, wherein the vaccine elicits a humoral response against the antigen in the subject.
30. The method of claim 22, wherein the vaccine elicits a cell-mediated response against the antigen in the subject.
31. The method of claim 22, wherein the vaccine elicits both a humoral and a cell-mediated response against the antigen in the subject.
32. The method of claim 22, wherein the vaccine further comprises pharmaceutically acceptable excipient, auxiliary substance, adjuvant, wetting or emulsifying agent, or pH buffering agent.
33. A composition comprising: an immunogenic amount of one or more purified antigens and an amount of one or more purified chemokines, or purified

fragments or derivatives thereof, effective to enhance the immune response to said antigen(s); and a pharmaceutically acceptable carrier.

34. The composition of claim 33, wherein the one or more chemokines are selected from the group consisting of: MDC, SDF-1, BLC, and MCP-1.
35. The composition of claim 33, wherein the one or more chemokines are selected from a chemokine class selected from the group consisting of: CC, CXC, C-C and CX3C.
36. The composition of claim 33, wherein the one or more chemokines are selected from the group consisting of: Macrophage-derived chemokine, Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EBI1-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating

protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

37. The composition of claim 33, wherein the fragment(s) or derivative(s) are truncation isoforms.
38. The composition of claim 33, wherein the one or more chemokine fragment includes an MDC fragment selected from the group consisting of amino acid numbers 2-69, 3-69, 5-69, 7-69 and 9-69 of SEQ ID NO: 2.
39. The composition of claim 33, wherein the one or more chemokine fragment includes an MDC fragment selected from the group consisting of amino acid numbers 2-69, 3-69, 5-69, 7-69 and 9-69 of SEQ ID NO: 2, which derivative has activity to enhance the efficacy of the vaccine.
40. The composition of claim 33, wherein the one or more chemokine derivatives has one or more insertions of or substitutions with one or more non-classical amino acids relative to a corresponding wildtype chemokine, which derivative has activity to enhance the efficacy of the vaccine.
41. The composition of claim 33, wherein the one or more chemokine derivatives has one or more conservative substitutions in sequence relative to a corresponding wildtype chemokine, which derivative has activity to enhance the efficacy of the vaccine.
42. The composition of claim 33, wherein the chemokine is a human chemokine.
43. The composition of claim 33, wherein the antigen is an HIV antigen.
44. The composition of claim 43, wherein the antigen is HIV associated gp120 protein.
45. A composition comprising an amount of a first set of purified nucleic acids comprising one or more nucleotide sequences encoding one or more antigens

and a second set of purified nucleic acids comprising one or more nucleotide sequences encoding one or more chemokines, or fragments or derivatives thereof, wherein the antigen(s) and the chemokine(s), or fragment(s) or derivative(s) thereof, are expressed from said first set of nucleic acid(s) and second set of nucleic acid(s) in a coordinated manner such that upon introduction into a suitable cell, the amount of said first set of nucleic acid(s) is sufficient to express an immunogenic amount of the antigen and the amount of the said second set of nucleic acid(s) is effective in enhancing the efficacy of the vaccine; and a pharmaceutically acceptable carrier.

46. The composition of claim 45, wherein the chemokine is MDC and the nucleic acid encoding the MDC comprises the nucleotide sequence of SEQ ID NO: 1.
47. The composition of claim 45, wherein the chemokine derivative(s) have deletional, insertional or substitutional mutations and/or combinations thereof, and the derivative(s) have activity to enhance the efficacy of the vaccine.
48. The composition of claim 45, further comprising pharmaceutically acceptable excipient, auxiliary substance, adjuvant, wetting or emulsifying agent, or pH buffering agent.
49. A composition comprising a first set of purified nucleotide sequences encoding one or more antigens and a second set of purified nucleotide sequences encoding one or more chemokines, or fragments or derivatives thereof, wherein the antigen(s) and the chemokine(s) are expressed in a coordinated manner such that upon introduction into a suitable cell, the sets produce an amount of said antigen(s) that is immunogenic and an amount of chemokine(s), or fragment(s) or derivative(s) thereof, that is effective in enhancing the efficacy of the vaccine relative to a corresponding vaccine composition without such chemokine(s), fragment(s) or derivative(s) thereof.
50. The composition of claim 49, wherein the one or more chemokines are selected from the group consisting of: Macrophage-derived chemokine,

Monocyte chemotactic protein 1, Monocyte chemotactic protein 2, Monocyte chemotactic protein 3, Monocyte chemotactic protein 4, activated macrophage specific chemokine 1, Macrophage inflammatory protein 1 alpha, Macrophage inflammatory protein 1 beta, Macrophage inflammatory protein 1 gamma, Macrophage inflammatory protein 1 delta, Macrophage inflammatory protein 2 alpha, Macrophage inflammatory protein 3 alpha, Macrophage inflammatory protein 3 beta, Regulated upon activation, normal T cell expressed and secreted (and its variants), I-309, EBI1-ligand chemokine, Pulmonary and activation regulated chemokine, Liver and activation-regulated chemokine, Thymus and activation regulated chemokine, Eotaxin (and variants), Human CC chemokine 1, Human CC chemokine 2, Human CC chemokine 3, IL-10-inducible chemokine, liver-expressed chemokine, 6Ckine, Exodus 1, Exodus 2, Exodus 3, thymus-expressed chemokine, Secondary Lymphoid tissue chemokine, Lymphocyte and Monocyte chemoattractant; Monotactin, Activation-induced, chemokine-related molecule, Myeloid progenitor inhibitory factor-1, Myeloid progenitor inhibitory factor-2, Stromal cell-derived factor 1 alpha, Stromal cell-derived factor 1 beta, B-cell-attracting chemokine 1, HuMIG, H174, Interferon-stimulated T-cell alpha chemoattractant, Interleukin-8, IP-10, platelet factor 4, growth-regulated gene-alpha, growth-regulated gene-beta, growth-regulated gene-gamma, Neutrophil-activating protein 2, ENA-78, granulocyte chemotactic protein 2, LYMPHOTACTIN, and Fractalkine/neurotactin.

51. The method of claim 49, wherein the one or more chemokines are selected from a chemokine class selected from the group consisting of: CC, CXC, C-C and CX3C.
52. The method of claim 49, wherein the one or more chemokines are selected from the group consisting of: MDC, SDF-1, BLC, and MCP-1.
53. The composition of claim 49, wherein the fragment(s) or derivative(s) are truncation isoforms.

54. The composition of claim 49, wherein the nucleic acid is administered directly to the subject.
55. The composition of claim 49, wherein the nucleic acid is introduced into a suitable host cell and said suitable host cell is introduced into the subject.

| | |
|--|-----|
| AGC CAA TGAAGAGCCT ACTCTGATGA CCGTGGCCTT GGCTCCTCCA GGAAGGCTCA | 348 |
| Ser Gln | |
| GGAGCCCTAC CTCCCCTGCCA TTATAGCTGC TCCCCGCCAG AAGCCTGTGC CAACTCTCTG | 408 |
| CATTCCCTGA TCTCCATCCC TGTGGCTGTC ACCCTTGGTC ACCTCCGTGC TGTCACTGCC | 468 |
| ATCTCCCCC TGACCCCTCT AACCATCCT CTGCCTCCCT CCCTGCAGTC AGAGGGTCCT | 528 |
| GTTCCCATCA GCGATTCCCC TGCTTAAACC CTTCCATGAC TCCCCACTGC CCTAAGCTGA | 588 |
| GGTCAGTCTC CCAAGCCTGG CATGTGGCCC TCTGGATCTG GGTTCATCT CTGTCTCCAG | 648 |
| CCTGCCCACT TCCCTTCATG AATGTTGGGT TCTAGCTCCC TGTCTCCAA ACCCATACTA | 708 |
| CACATCCAC TTCTGGGTCT TTGCCCTGGA TGTGTGCTGAC ACTCAGAAAG TCCCACCACC | 768 |
| TGCACATGTG TAGCCCCACC AGCCCTCCAA GGCATTGCTC GCCCAAGCAG CTGGTAATTC | 828 |
| CATTTCATGT ATTAGATGTC CCCTGGCCCT CTGTCCCCCTC TTAATAACCC TAGTCACAGT | 888 |
| CTCCGCAGAT TCTTGGGATT TGGGGGTTT CTCCCCCACC TCTCCACTAG TTGGACCAAG | 948 |

FIG. 1A-2

| | | | | | | |
|-------------------|-------------------|-------------------|-------------------|-------------------|------------------|-------------|
| <u>GTTTCTAGCT</u> | <u>AAGTTACTCT</u> | <u>AGTCTCCAAG</u> | <u>CCTCTAGCAT</u> | <u>AGAGCACTGC</u> | <u>AGACAGGCC</u> | <u>1008</u> |
| TGGCTCAGAA | TCAGAGCCCA | GAAAGTGGCT | GCAGACAAA | TCAATAAAC | TAATGTCCCT | 1068 |
| CCCCTCTCCC | TGCCAAAAGG | CAGTTACATA | TCAATACAGA | GACTCAAGT | CACTAGAAAT | 1128 |
| GGGCCAGCTG | GGTCAATGTG | AAGCCCCAAA | TTTGCCCAGA | TTCACCTTTC | TTCCCCCACT | 1188 |
| CCCTTTTTT | TTTTTTTTT | TTTGAGATGG | AGTTTCGCTC | TTGTCACCCA | CGCTGGAGTG | 1248 |
| CAATGGGTG | GTCTTGGCTT | ATTGAAGCCT | CTGCCTCCTG | GGTTCAAGTG | ATTCTCTTGC | 1308 |
| CTCAGCCTCC | TGAGTAGCTG | GGATTACAGG | TTCTTGCTAC | CACGCCCAGC | TAATTTTGT | 1368 |
| ATTTTTAGTA | GAGACGAGGC | TTCAACCATGT | TGGCCAGGCT | GGTCTCGAAC | TCCTGTCCCTC | 1428 |
| AGGTAATCCG | CCCACCTCAG | CCTCCCCAAG | TGCTGGGATT | ACAGGCGTGA | GCCACAGTGC | 1488 |
| CTGGCCCTCT | CCCTCTCCCC | ACTGCCCCCC | CCAACTTTTT | TTTTTTTTTT | ATGGCAGGGT | 1548 |
| CTCACTCTGT | CGCCCAGGCT | GGAGTGCAGT | GGCGTGATCT | CGGCTCACTA | CAACCTCGAC | 1608 |
| CTCCTGGGTT | CAAGTGATTC | TCCCACCCCA | GCCTCCCAAG | TAGCTGGGAT | TACAGGTGTG | 1668 |

FIG. 1A-3

| | | | | | | |
|-------------|------------|------------|------------|-------------|-------------|------|
| TGCCACTACG | GCTGGCTAAT | TTTTGTATTT | TTAGTAGAGA | CAGGTTTCAC | CATATTGGCC | 1728 |
| AGGCTGGTCT | TGAACTCCTG | ACCTCAAGTG | ATCCACCTTC | CTTG TGCTCC | CAAAGTGCTG | 1788 |
| AGATTACAGG | CGTGAGCTAT | CACACCCAGC | CTCCCCCTTT | TTTTCCCTAAT | AGGAGACTCC | 1848 |
| TGTACCTTTC | TTCGTTTTAC | CTATGTGTCG | TGCTGCTTA | CATTTCCTTC | TCCCCCTCAGG | 1908 |
| CTTTTTTTGG | GTGGTCCTCC | AACCTCCAAT | ACCCAGGCCT | GGCCTCTTCA | GAGTACCCCC | 1968 |
| CATTCCACTT | TCCCTGCCTC | CTTCCTTAAA | TAGCTGACAA | TCAAATTCTAT | GCTATGGTGT | 2028 |
| GAAAGACTAC | CTTTGACTTG | GTATTATAAG | CTGGAGTTAT | ATATGTATTT | GAAAACAGAG | 2088 |
| TAAATACTTA | AGAGGCCCAA | TAGATGAATG | GAAGAATTTT | AGGAACTGTG | AGAGGGGGAC | 2148 |
| AAGGTGAAGC | TTTCCTGGCC | CTGGGAGGAA | GCTGGCTGTG | GTAGCGTAGC | GCTCTCTCTC | 2208 |
| TCTGTCTGTG | GCAGGAGCCA | AAGAGTAGGG | TGTAATTGAG | TGAAGGAATC | CTGGGTAGAG | 2268 |
| ACCATTCCTCA | GGTGGTTGGG | CCAGGCTAAA | GACTGGGAGT | TGGGTCTATC | TATGCCTTTC | 2328 |
| TGGCTGATTT | TTGTAGAGAC | GGGGTTTTGC | CATGTTACCC | AGGCTGGTCT | CAAACTCCTG | 2388 |

FIG. 1A-4

| | |
|--|------|
| GGCTCAAGCG ATCCTCCTGG CTCAGCCTCC CAAAGTGCTG GGATTACAGG CGTGAATCAC | 2448 |
| TGGGCCCTGGC TTCCTCTTCC TCTTGAGAAA TATCTTTTC ATACAGCAAG TATGGGACAG | 2508 |
| CAGTGTCCTCA GGTAAGGAC ATAAATGTTA CAAGTGCTG GTCCTTTCTG AGGGAGGCTG | 2568 |
| GTGCCGCTCT GCAGGGTATT TGAACCTGTG GAATTGGAGG AGGCCATTTC ACTCCCTGAA | 2628 |
| CCCAGCCTGA CAAATCACAG TGAGAATGTT CACCTTATAG GCTTGCTGTG GGGCTCAGGT | 2688 |
| TGAAAGTGTG GGGAGTGACA CTGCCCTAGGC ATCCAGCTCA GTGTCATCCA GGGCCTGTGT | 2748 |
| CCCTCCCGAA CCCAGGGTCA ACCTGCCCTGC CACAGGCACT AGAAGGACGA ATCTGCCTAC | 2808 |
| TGCCCATGAA CGGGGCCCTC AAGCGTCCTG GGATCTCCTT CTCCCTCCTG TCCTGTCCCT | 2868 |
| GCCCCTCAGG ACTGCTGGAA AATAAATCCT TTAAATAGT AAAAAAAAAA AAAA | 2923 |

5/6

FIG. 1A-5

FIG. 1A

| |
|-----------|
| FIG. 1A-1 |
| FIG. 1A-2 |
| FIG. 1A-3 |
| FIG. 1A-4 |
| FIG. 1A-5 |

6/6

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Ala | Arg | Leu | Gln | Thr | Ala | Leu | Leu | Val | Val | Leu | Val | Leu | Leu | Ala |
| -24 | | | | -20 | | | | | -15 | | | | | -10 | |
| Val | Ala | Leu | Gln | Ala | Thr | Glu | Ala | Gly | Pro | Tyr | Gly | Ala | Asn | Met | Glu |
| | | | -5 | | | | | 1 | | | | 5 | | | |
| Asp | Ser | Val | Cys | Cys | Arg | Arg | Asp | Tyr | Val | Arg | Tyr | Arg | Leu | Pro | Leu |
| | 10 | | | | | 15 | | | | | | 20 | | | Arg |
| Val | Lys | His | Phe | Tyr | Tyr | Trp | Thr | Ser | Asp | Ser | Cys | Pro | Arg | Pro | Gly |
| 25 | | | | 30 | | | | | 35 | | | | | | 40 |
| Val | Val | Leu | Leu | Thr | Phe | Arg | Asp | Lys | Glu | Ile | Cys | Ala | Asp | Pro | Arg |
| | | | | 45 | | | | | 50 | | | | | 55 | |
| Val | Pro | Trp | Val | Lys | Met | Ile | Leu | Asn | Lys | Leu | Ser | Gln | | | |
| | | | | 60 | | | | 65 | | | | | | | |

FIG. 1B

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANT: Gallo, Robert C.
DeVico, Anthony L.
Garzino, Alfredo

(ii) TITLE OF THE INVENTION: METHOD AND COMPOSITION TO ENHANCE
THE EFFICACY OF A VACCINE USING MACROPHAGE DERIVED CHEMOKINE

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Pennie & Edmonds LLP
(B) STREET: 1155 Avenue of the Americas
(C) CITY: New York
(D) STATE: New York
(E) COUNTRY: USA
(F) ZIP: 10036/2711

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ Version 2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: To be assigned
(B) FILING DATE: Herewith
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Misrock, S. Leslie
(B) REGISTRATION NUMBER: 18,872
(C) REFERENCE/DOCKET NUMBER: 8769-029

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 212-790-9090
(B) TELEFAX: 212-869-8864
(C) TELEX: 66141 PENNIE

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2923 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: mat_peptide
(B) LOCATION: 92..298

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

| | |
|--|-----|
| GAGACATACA GGACAGAGC ATG GCT CGC CTA CAG ACT GCA CTC CTG GTT GTC | 52 |
| Met Ala Arg Leu Gln Thr Ala Leu Val | |
| -24 -20 -15 | |
| CTC GTC CTC CTT GCT GTG GCG CTT CAA GCA ACT GAG GCA GGC CCC TAC | 100 |
| Leu Val Leu Ala Val Ala Leu Gln Ala Thr Glu Ala Gly Pro Tyr | |
| -10 -5 1 | |
| GGC GCC AAC ATG GAA GAC AGC GTC TGC CGT GAT TAC GTC CGT TAC | 148 |
| Gly Ala Asn Met Glu Asp Ser Val Cys Arg Asp Tyr Val Arg Tyr | |
| 5 10 15 | |
| CGT CTG CCC CTG CGC GTG AAA CAC TTC TAC TGG ACC TCA GAC TCC | 196 |
| Arg Leu Pro Leu Arg Val Lys His Phe Tyr Trp Thr Ser Asp Ser | |
| 20 25 30 35 | |
| TGC CCG AGG CCT GGC GTG TTG CTA ACC TTC AGG GAT AAG GAG ATC | 244 |
| Cys Pro Arg Pro Gly Val Leu Thr Phe Arg Asp Lys Glu Ile | |
| 40 45 50 | |
| TGT GCC GAT CCC AGA GTG CCC TGG GTG AAG ATG ATT CTC AAT AAG CTG | 292 |
| Cys Ala Asp Pro Arg Val Pro Trp Val Lys Met Ile Leu Asn Lys Leu | |
| 55 60 65 | |
| AGC CAA TGAAGAGCCT ACTCTGATGA CCGTGGCCTT GGCTCCTCCA GGAAGGCTCA | 348 |
| Ser Gln | |

GGAGCCCTAC CTCCCTGCCA TTATAGCTGC TCCCCGCCAG AAGCCTGTGC CAACTCTCTG 408
 CATTCCCTGA TCTCCATCCC TGTGGCTGTC ACCCTTGGTC ACCTCCGTGC TGTCAGTGCC 468
 ATCTCCCCC TGACCCCTCT AACCCTATCCT CTGCCCTCCCT CCCTGCAGTC AGAGGGTCTT 528
 GTTCCCATCA GCGATTCCCC TGCTTAAACC CTTCATGAC TCCCCACTGC CCTAAGCTGA 588
 GGTCAGTCTC CCAAGCCTGG CATGTGGCCC TCTGGATCTG GGTTCATCT CTGTCTCCAG 648
 CCTGCCCACT TCCCTTCATG AATGTTGGGT TCTAGCTCCC TGTCTCCAA ACCCATACTA 708
 CACATCCAC TTCTGGGTCT TTGCTGGGA TGTGCTGAC ACTCAGAAAG TCCCACCACC 768
 TGCACATGTG TAGCCCCACC AGCCCTCCAA GGCATTGCTC GCCCAAGCAG CTGGTAATT 828
 CATTTCATGT ATTAGATGTC CCCTGGCCCT CTGTCCCTC TTAATAACCC TAGTCACAGT 888
 CTCCGCAGAT TCTTGGGATT TGGGGGTTTT CTCCCCACC TCTCCACTAG TTGGACCAAG 948
 GTTCTAGCT AAGTTACTCT AGTCTCCAAG CCTCTAGCAT AGAGCACTGC AGACAGGCC 1008
 TGCAGCATGA TCAGAGCCCA GAAAGTGGCT GCAGACAAA TCAATAAAAC TAATGTCCCT 1068
 CCCCTCTCCC TGCCAAAAGG CAGTTACATA TCAATACAGA GACTCAAGGT CACTAGAAAT 1128
 GGGCCAGCTG GGTCAATGTG AAGCCCCAAA TTTGCCCAGA TTCACCTTTC TTCCCCCACT 1188
 CCCTTTTTTT TTTTTTTTTT TTTGAGATGG AGTTTCGCTC TTGTACCCA CGCTGGAGTG 1248
 CAATGGTGTG GTCTTGGCTT ATTGAAGCCT CTGCCCTCTG GGTTCAGTG ATTCTCTTGC 1308
 CTCAGCCTCC TGAGTAGCTG GGATTACAGG TTTCTGCTAC CACGCCCAGC TAATTTTTCT 1368
 ATTTTATAGT GACACGAGG TTCACCATGT TGGCCAGGCT GGTCTCGAAC TCCTGTCTGT 1428
 AGGTAATCCG CCACCTCAG CCTCCCAAAG TGCTGGGATT ACAGGCGTGA GCCACAGTGC 1488
 CTGGCTCTT CCTCTCCCC ACTGCCCCCC CCAACTTTTT TTTTTTTTTT ATGGCAGGGT 1548
 CTCACCTGT CGCCAGGCT GGAGTGCAGT GGCGTGATCT CGGCTCACTA CAACCTCGAC 1608
 CTCCTGGGTT CAAGTGATT TCCCACCCCA GCCTCCCAAG TAGCTGGGAT TACAGGTGTG 1668
 TGCCACTACG GCTGGCTAAT TTTGTATTT TTAGTAGAGA CAGGTTTCAC CATATTGGCC 1728
 AGGTGGTCT TGAACCTCTG ACCTCAAGTG ATCCACCTTC CTGTGTCTCC CAAAGTGTG 1788
 AGATTACAGG CGTGAGCTAT CACACCCAGC CTCCCCCTT TTTTCTAAT AGGAGACTCC 1848
 TGTACCTTTC TTCGTTTAC CTATGTGTCG TGTCTGCTTA CATTTCTTTC TCCCCTCAGG 1908
 CTTTTTTTGG GTGGTCTCC AACCTCCAAT ACCCAGGCCT GGCCTCTTCA GAGTACCCCC 1968
 CATTCCACTT TCCCTGCCTC CTCTCTTAA TAGCTGACAA TCAAATTCAT GCTATGGTGT 2028
 GAAAGACTAC CTTTGACTTG GTATTATAAG CTGGAGTTAT ATATGTATTT GAAAACAGAG 2088
 TAAATACTTA AGAGGCCAAA TAGATGAATG GAAGAATTT AGGAACGTG AGAGGGGGAC 2148
 AAGGTGAAGC TTTCTGGCC CTGGGAGGAA GCTGGCTGTG GTAGCGTAGC GCTCTCTCTC 2208
 TCTGTCTGTG GCAGGAGCCA AAGAGTAGGG TGTAATTGAG TGAAGGAATC CTGGGTAGAG 2268
 ACCATTCTCA GGTGGTTGGG CCAGGCTAAA GACTGGGAGT TGGGTCTATC TATGCTTTC 2328
 TGGCTGATTT TTGTAGAGAC GGGGTTTTGC CATGTTACCC AGGCTGGTCT CAACTCCTG 2388
 GGCTCAAGCG ATCCTCTTGG CTCAGCCTCC CAAAGTGTG GGATTACAGG CGTGAATCAC 2448
 TGCGCTGGC TTCCTCTTCC TCTTGAGAAA TATTCTTTTC ATACAGCAAG TATGGGACAG 2508
 CAGTGTCCTA GGTAAAGGAC ATAAATGTTA CAAGTGTCTG GTCCTTCTG AGGGAGGCTG 2568
 GTGCCGCTCT GCAGGGTATT TGAACCTGTG GAATTGGAGG AGGCCATTTT ACTCCCTGAA 2628
 CCCAGCCTGA CAAATCACAG TGAGAATGTT CACCTTATAG GCTTGTCTGT GGGCTCAGGT 2688
 TGAAAGTGTG GGGAGTGACA CTGCCTAGGC ATCCAGCTCA GTGTCAATCCA GGGCTGTGT 2748
 CCCTCCCGAA CCCAGGGTCA ACCTGCCTGC CACAGGCACT AGAAGGACGA ATCTGCCTAC 2808
 TGCCCATGAA CGGGGCCCTC AAGCGTCTG GGAATCTCTT CTCCCTCTG TCCTGTCTT 2868
 GCCCTCAGG ACTGCTGGAA AATAATCCT TTAATAAGT AAAAAA AAAAA 2923

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 93 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Arg Leu Gln Thr Ala Leu Val Leu Val Leu Ala
 -24 -20 -15 -10
 Val Ala Leu Gln Ala Thr Glu Ala Gly Pro Tyr Gly Ala Asn Met Glu
 -5 1 5
 Asp Ser Val Cys Arg Asp Tyr Val Arg Tyr Arg Leu Pro Leu Arg
 10 15 20
 Val Lys His Phe Tyr Trp Thr Ser Asp Ser Cys Pro Arg Pro Gly
 25 30 35 40
 Val Leu Thr Phe Arg Asp Lys Glu Ile Cys Ala Asp Pro Arg
 45 50 55
 Val Pro Trp Val Lys Met Ile Leu Asn Lys Leu Ser Gln
 60 65

C-CHEMOKINES

LYMPHOTACTIN

(SCM-1)

D63789 D63790

CX3C-chemokines

Fractalkine/neurotactin

U91835 U84487

LOCUS HSU83171 2923 bp mRNA PRI 31-MAY-1997
 DEFINITION Human macrophage-derived chemokine precursor (MDC) mRNA,
 complete
 cds.
 ACCESSION U83171
 NID g1931580
 KEYWORDS
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
 Hominidae;
 Homo.
 REFERENCE 1 (bases 1 to 2923)
 AUTHORS Godiska,R., Chantry,D., Raport,C.J., Sozzani,S., Allavena,P.,
 Leviten,D., Mantovani,A. and Gray,P.W.
 TITLE Human macrophage-derived chemokine (MDC), a novel
 chemoattractant
 for monocytes, monocyte-derived dendritic cells, and natural
 killer
 cells
 JOURNAL J. Exp. Med. 185 (9), 1595-1604 (1997)
 MEDLINE 97296313
 REFERENCE 2 (bases 1 to 2923)
 AUTHORS Godiska,R. and Gray,P.W.
 TITLE Direct Submission
 JOURNAL Submitted (23-DEC-1996) ICOS Corporation, 22021 20th Avenue SE,
 Bothell, WA 98021, USA
 FEATURES Location/Qualifiers
 source 1..2923
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="16"
 gene 20..301
 /gene="MDC"
 sig_peptide 20..91
 /gene="MDC"
 CDS 20..301
 /gene="MDC"
 /function="chemotactic for dendritic cells and natural
 killer cells"
 /codon_start=1
 /product="macrophage-derived chemokine precursor"
 /db_xref="PID:g1931581"
 /translation="MARLQTALLVVLVLLAVALQATEAGPYGANMEDSVCCRDYVRYR
 LPLRVVKHFYWTSDSCPRPGVLLTFRDKEICADPRVPWVKMILNKLQ"
 mat_peptide 92..298
 /gene="MDC"
 /product="macrophage-derived chemokine"
 repeat_region complement(1194..1805)
 /rpt_family="ALU"
 repeat_region complement(2335..2443)
 /rpt_family="ALU"
 BASE COUNT 605 a 861 c 669 g 788 t
 ORIGIN
 1 gagacataca ggacagagca tggetcgcc acagactgca ctcttggttg tctctgtcct
 61 ccttgctgtg gcgcttcaag caactgaggc aggcccttac ggcgccaaca tgggaagacag
 121 cgtctgtgc cgtgattacg tccgttaccg tctgccccctg cgcgtggtga aacacttcta
 181 ctggacctca gactcctgcc cgaggcctgg cgtggtgttg ctaaccttca gggataagga
 241 gatctgtgcc gatcccagag tgcccctggg gaagatgatt ctcaataagc tgagccaatg
 301 aagagccctac tctgatgacc gtggccttgg ctctccagg aaggctcagg agccctacct
 361 ccttgccatt atagctgctc cccgccagaa gcctgtgcca actctctgca ttccctgatc
 421 tccatccctg tggctgtcac ccttggtcac ctccgtgctg tcaactgcat ctccccctg
 481 accctcttaa cccatcctct gcctccctcc ctgcagtcag agggctcctg tcccatcagc
 541 gattccccctg cttaaaccct tccatgactc cccactgccc taagctgagg tcagctctcc
 601 aagcctggca tgtggccctc tggatctggg ttccatctct gtctccagcc tgcccacttc
 661 ccttcatgaa tgttgggttc tagctccctg ttctccaaac ccatactaca catccactt
 721 ctgggtcttt gcctgggatg ttgctgacac tcagaaagtc ccaccactg cacatgtgta
 781 gccccaccag cctccaagg cattgctcgc ccaagcagct ggtaattcca tttcatgtat
 841 tagatgtccc ctggccctct gtccctctt aataacccta gtcacagtct ccgcagattc

```

901 ttgggatttg ggggttttct cccccacctc tccactagtt ggaccaaggt ttctagctaa
961 gttactctag tctccaagcc tctagcatag agcactgcag acaggccctg gctcagaatc
1021 agagccccaga aagtgggtgc agacaaaatc aataaaacta atgtccctcc cctctccctg
1081 ccaaaaggca gttacatc aatacagaga ctcaaggta ctagaaatgg gccagctggg
1141 tcaatgtgaa gcccacaatt tgcccagatt cacctttctt cccccactcc cttttttttt
1201 tttttttttt tgagatggag ttctgcctct gtacccacag ctggagtgca atggtgtggt
1261 cttggcttat tgaagcctct gcctcctggg ttcaagtgat tctcttgctc cagcctcctg
1321 agtagctggg attacaggtt cctgctacca cgcccagcta atttttgtat ttttagtaga
1381 gacgaggctt caccatgttg gccaggctgg tctcgaactc ctgtcctcag gtaatccgcc
1441 cacctcagcc tcccaaagtg ctgggattac aggcgtgagc cacagtgcct ggcctcttcc
1501 ctctcccccac tgcccccccc aacttttttt ttttttttat ggcagggtct cactctgtcg
1561 ccaggctggg agtgcagtg cgtagctctg gctcactaca acctcgacct cctgggttca
1621 agtgattctc ccaccccagc ctcccaagta gctgggatta cagggtgtgtg ccactacggc
1681 tggctaattt ttgtattttt agtagagaca ggtttcacca tattggccag gctgggtctg
1741 aactcctgac ctcaagtgat ccaccttctt tgtgctccca aagtgtgtg attacaggcg
1801 tgagctatca caccagcctt cccctttttt ttccataatg gagactcctg tacctttctt
1861 cgttttacct atgtgtcgtg tctgcttaca ttctcttctc cctcaggctt ttttttgggt
1921 ggtctcccaa cctccaatac ccaggcctgg cctcttcaga gtacccccca ttccactttc
1981 cctgctcctt tcctaaata gctgacaatc aaatttcatg tatggtgtga aagactacct
2041 ttgacttggg attataagct ggagtatat atgtatttga aaacagagta aatacttaag
2101 aggccaaata gatgaatgga agaatttttag gaactgtgag agggggacaa ggtgaagctt
2161 tcttgccctt gggaggaagc tggctgtggt agcgtagcgc tctctctctc tgtctgtggc
2221 aggagccaaa gaggtaggtg taattgagtg aaggaaatctt gggtagagac cattctcagg
2281 tgggtgggctc aggcataaaga ctgggagttg ggtctatcta tgctttctg gctgattttt
2341 gatagagcgg ggttttgcca tgttaccag gctgggtctc aactcctggg ctcaagcgat
2401 cctcctggct cagcctccca aagtgtcggg attacaggcg tgaatcactg cgctggctt
2461 cctcttctct ttgagaaata ttcttttcat acagcaagta tgggacagca gtgtcccagg
2521 taaaggacat aaatgttaca agtgtctggt cctttctgag ggaggctggt gccgctctgc
2581 aggggtattg aacctgtgga attggaggag gccatttcac tccctgaacc cagcctgaca
2641 aatcacagtg agaattgtca ccttataggc ttgctgtggg gctcagggtg aaagtgtggg
2701 gagtgcact gcctaggcat ccagctcagt gtcacccagg gcctgtgtcc ctcccgaacc
2761 cagggtcaac ctgcttgcca caggcactag aaggacgaat ctgcctactg cccatgaacg
2821 gggccctcaa gcgtcctggg atctccttct ccctcctgtc ctgtccttgc ccctcaggac
2881 tgctggaaaa taaatccttt aaaatagtaa aaaaaaaaaa aaa

```

```

//
LOCUS      HSU83239      932 bp      mRNA      PRI      02-MAY-1997
DEFINITION Human CC chemokine STCP-1 mRNA, complete cds.
ACCESSION  U83239
NID        g2062424
KEYWORDS
SOURCE
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 932)
AUTHORS    Chang,M.S., McNinch,J., Elias III,C., Manthey,C.L.,
            Grosshans,D.,
            Meng,T., Boone,T. and Andrew,D.P.
TITLE      Molecular cloning and functional characterization of a novel CC
            chemokine STCP-1 which specifically acts on activated T
lymphocytes
JOURNAL     Unpublished
REFERENCE  2 (bases 1 to 932)
AUTHORS    Chang,M.S., McNinch,J., Elias III,C., Manthey,C.L.,
            Grosshans,D.,
            Meng,T., Boone,T. and Andrew,D.P.
TITLE      Direct Submission
JOURNAL     Submitted (26-DEC-1996) Research Computing, Amgen Institute,
620
            University Ave, Suite 706, Toronto, ON M5G 2C1, Canada
FEATURES
            source      1..932
                        /organism="Homo sapiens"
                        /note="Amgen EST program"
                        /db_xref="taxon:9606"
            CDS          15..296
                        /codon_start=1
                        /product="CC chemokine STCP-1"
                        /db_xref="PID:g2062425"

/translation="MARLQTALLVVLVLLAVALQATEAGPYGANMEDSVCCRDIYVRYR

```

```

                                LPLRVVKHFYWTSDSCP RP GVVLLTFRDKEICADPRVPWVKMILNKLSQ"
BASE COUNT      166 a      330 c      201 g      235 t
ORIGIN
    1 atacaggaca gagcatggct cgcctacaga ctgcactcct ggttgctcct gtcctccttg
   61 ctgtggcgct tcaagcaact gaggcaggcc cctacggcgc caacatggaa gacagcgtct
  121 gctgccgtga ttacgtccgt taccgtctgc ccctgcgcgt ggtgaaacac ttctactgga
  181 cctcagactc ctgcccaggg cctggcgtgg tgttgctaac cttcagggat aaggagatct
  241 gtgccgatcc cagagtggcc tgggtgaaga tgattctcaa taagctgagc caatgaagag
  301 cctactctga tgaccgtggc cttggctcct ccaggaaggc tcaggagccc tacctccctg
  361 ccattatagc tgctccccgc cagaagcctg tgccaactct ctgcattccc tgatctccat
  421 ccctgtggct gtcacccttg gtcaccctcg tgcgtgctact gccatctccc ccctgacccc
  481 tetaacccat cctctgcctc cctccctgca gtcagagggt cctgttccca tcagcgattc
  541 ccctgcttaa acccttccat gactccccac tgccctaagc tgagggtcagt ctcccaagcc
  601 tggcatgtgg ccctctggat ctgggttcca tctctgtctc cagcctgccc acttcccttc
  661 atgaatgttg ggttctagct ccctgttctc caaacccata ctacacatcc cacttctggg
  721 tctttgccctg ggatgttgct gacactcaga aagtccacc acctgccatc gtgtagcccc
  781 accagccctc caaggcattg ctgcaccaag cagctggtaa ttccatttca tgtattagat
  841 gtccctggc cctctgtccc ctcttaataa ccctagtcac agtctccgca gattcttggg
  901 atttgggggt tttctcccc accctccac ta

//
LOCUS      HSMCP1          725 bp      RNA          PRI          03-APR-1995
DEFINITION H.sapiens mRNA for monocyte chemoattractant protein 1 (MCP-1).
ACCESSION  X14768
NID        g34513
KEYWORDS   monocyte chemoattractant protein 1.
SOURCE     human.
  ORGANISM Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 725)
AUTHORS    Yoshimura,T., Yuhki,N., Moore,S.K., Appella,E., Lerman,M.I. and
            Leonard,E.J.
TITLE      Human monocyte chemoattractant protein-1 (MCP-1). Full-length
cdna
            cloning, expression in mitogen-stimulated blood mononuclear
            leukocytes, and sequence similarity to mouse competence gene JE
JOURNAL    FEBS Lett. 244 (2), 487-493 (1989)
MEDLINE    89153605
COMMENT    ZAP11.
FEATURES   Location/Qualifiers
            source          1..725
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /cell_type="glioma cells"
                        /cell_line="U105MG"
                        /clone_lib="lambda"
            sig_peptide     54..122
                        /note="signal peptide (AA -23 to -1)"
            CDS              54..353
                        /codon_start=1
                        /product="monocyte chemoattractant preprotein"
                        /db_xref="PID:g34514"
                        /db_xref="SWISS-PROT:P13500"

/translation="MKVSAALLCLLLIAATFIPQGLAQPDAINAPVTCCYNFTNRKIS
VQRLASYRRITSSKCPKEAVIFKTIVAKEICADPKQKWVQDSMDHLDKQTQTPKT"
            mat_peptide     123..350
                        /note="MCP-1 (AA 1 - 76)"
            misc_feature     162..170
                        /note="pot. N-linked glycosylation site"
            misc_feature     707..712
                        /note="pot. polyA signal"
            polyA_site       725
                        /note="polyA site"
BASE COUNT      208 a      171 c      126 g      220 t
ORIGIN
    1 ctaaccacaga aacatccaat tctcaaactg aagctcgcac tctcgctccc agcatgaaag
   61 tctctgccgc ccttctgtgc ctgctgtctc tagcagccac cttcattccc caagggtctg
  121 ctcagccaga tgcaatcaat gccccagtc a cctgctgtta taacttccc aataggaaga
  181 tctcagtgc a gaggtcgcg agctatagaa gaatcaccag cagcaagtgt cccaaagaag

```

```

241 ctgtgatctt caagaccatt gtggccaagg agatctgtgc tgacccaag cagaagtggg
301 ttcaggattc catggaccac ctggacaagc aaacccaac tccgaagact tgaacactca
361 ctccacaacc caagaatctg cagctaactt attttccct agctttcccc agacaccctg
421 ttttatttta ttataatgaa tttgtgtgtg tgatgtgaaa cattatgcct taagtaatgt
481 taattcttat ttaagttatt gatgttttaa gtttatcttt catgggtacta gtgtttttta
541 gatacagaga cttggggaaa ttgcttttcc tcttgaacca cagttctacc cctgggatgt
601 tttgagggtc tttgcaagaa tcattaatac aaagaatttt ttttaacatt ccaatgcatt
661 gctaaaatat tattgtggaa atgaatattt tgtaactatt acaccaaata aatatatttt
721 tgtac

```

//

```

LOCUS      HSMCP2      2991 bp      DNA      PRI      20-MAR-1997
DEFINITION H.sapiens MCP-2 gene.
ACCESSION  X99886
NID        g1905800
KEYWORDS   MCP-2 gene; monocyte chemotactic protein 2; SCYA10 gene.
SOURCE     human.
  ORGANISM Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 2991)
AUTHORS    Van Coillie,E., Fiten,P., Nomiyama,H., Sakaki,Y., Miura,R.,
            Yoshie,O., Van Damme,J. and Opdenakker,G.
TITLE      The human MCP-2 gene (SCYA8): cloning, sequence analysis,
tissue     expression, and assignment to the CC chemokine gene contig on
            chromosome 17q11.2
JOURNAL    Genomics 40 (2), 323-331 (1997)
MEDLINE    97237052
REFERENCE  2 (bases 1 to 2991)
AUTHORS    Opdenakker,G.M.M.
TITLE      Direct Submission
JOURNAL    Submitted (07-AUG-1996) G.M.M. Opdenakker, Rega Institute for
            Medical Research, Minderbroedersstraat 10, B 3000 Leuven,

```

BELGIUM

```

FEATURES             Location/Qualifiers
    source             1..2991
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /chromosome="17"
                        /map="q11.2"
    repeat_region      209..219
                        /note="DR-A"
                        /rpt_type=DIRECT
    repeat_region      240..248
                        /note="DR-B"
                        /rpt_type=DIRECT
    CAAT_signal        296..300
    repeat_region      310..318
                        /note="IR-A"
                        /rpt_type=INVERTED
    repeat_region      406..415
                        /note="DR-B"
                        /rpt_type=DIRECT
    repeat_region      407..416
                        /note="IR-B"
                        /rpt_type=INVERTED
    repeat_region      425..435
                        /note="DR-A"
                        /rpt_type=DIRECT
    repeat_region      429..437
                        /note="IR-B"
                        /rpt_type=INVERTED
    repeat_region      455..465
                        /note="IR-C"
                        /rpt_type=INVERTED
    TATA_signal        467..472
    repeat_region      492..502
                        /note="IR-C"
                        /rpt_type=INVERTED
    repeat_region      492..500
                        /note="IR-A"

```

```

        /rpt_type=INVERTED
exon      534..639
          /gene="MCP-2 (SCYA10)"
          /number=1
gene      534..1969
          /gene="MCP-2 (SCYA10)"
CDS       join(534..639,1331..1448,1864..1969)
          /gene="MCP-2 (SCYA10)"
          /codon_start=1
          /product="monocyte chemotactic protein-2"
          /db_xref="PID:e279930"
          /db_xref="PID:g1905801"

/translation="MLKLTPPSKMKVSAALLCLLLMAATFSPQGLAQPDVSVIPITC
CFNVINRKIPIQRLESYTRITNIQCPKEAVIFKTQRGKEVCADPKERWVRDSMKHLDQ
IFQNLKP"
intron    640..1330
          /gene="MCP-2 (SCYA10)"
          /number=1
exon      1331..1448
          /gene="MCP-2 (SCYA10)"
          /number=2
intron    1449..1863
          /gene="MCP-2 (SCYA10)"
          /number=2
exon      1864..1969
          /gene="MCP-2 (SCYA10)"
          /number=3
BASE COUNT      799 a      709 c      632 g      851 t
ORIGIN
1 agattctggg gcattaagac ttagttccag gattctgtca ttctgccaac gttctgtggc
61 tgggggttcta aaggagcttg cctggccttag aactgcaagt gactctagtg tgatggagag
121 caccagcaaa gccttagggc ccatcccttg cctctctgta ccacagagg ggtaagcctt
181 ggctctcttc cactatgacg tcagcttcca ttcttctttt cttatagaca attttccatt
241 tcaaggaaat cagagccctt aatagttcag tgaggtcact ttgctgagca caatcccata
301 ccttccagcc tctgtctcac agagcctaag caaaagatag aaactcacia cttccttggt
361 ttgttatctg gaaattatcc caggatctgg tgcttactca gcatattcaa ggaaggtctt
421 acttcattct tcttgattg tgaccatgcc caggctctct gctccctata aaaggcaggc
481 agagccaccg aggagcagag aggttgagaa caaccagaa accttcacct ctcatgctga
541 agctcacacc cttgccctcc aagatgaagg tttctgcagc gcttctgtgc ctgctgtca
601 tggcagccac tttcagccct cagggacttg ctacagccagg taagacctct ccttttttaa
661 ggggagacca aaagaggaat taagaagagc cattatgtca cagctcata ggaacaaaac
721 cagaactaaa ggctcaggtc actgaggctg gttcccttga tcttctctga cccagtttt
781 gggaggagac agtggagccg ctacagcaac aaccttccca ttgtttgggg aaataatcca
841 gaacgaagaa ctgtttctca ctgtgggtgt aaaggacatt tcaggccgta gtggagaggg
901 agaaaactatt gcctgaagct tcaaattttg gttatgggtt agtgtaacct ccagaaacat
961 ggctgtgtaa agaggatgag gaccagagg aatctcagcg tatggcatag gctaactcta
1021 aagcccatga ggatgaaaga ctgggaagca aggtattgga acttatgttc ccagtgtcag
1081 aagtttttgg ttagtagaca aggactagct tgttactcaa aatgtttcca cccagctga
1141 acaatgacgg gccgcagagt tcaatagagg aaagagactc acaggcaaca ttttatctct
1201 gggatctgga ctaagacact gaacttgagg tggtgacttc ttgggtctct ccttcttctt
1261 cttcttttcc ttacaaatgc acacttacgg tgggtcctaa atgtctcatt ctttgcaaaa
1321 tttctttcag attcagtttc cattccaatc acctgctgct ttaacgtgat caataggaaa
1381 attcctatcc agaggctgga gagctacaca agaatcacca acatccaatg tcccaaggaa
1441 gctgtgatgt gagtggacag tgcctggcac ccccatccta aagttctgat ggacaacata
1501 gagaagtcaa gattcatgtc catatgagtc ggaatgcatat aacttctatc caaaggggac
1561 ctctacccca tagagaaact cagtcctgta gaaggagtec ataactgttc taggattccc
1621 ttctaggggc ttgtgaaac taacccaata tctgtagcca ggacctgga gggtttcacc
1681 tggacagcaa gagcagagct tcttcttgga gcttcttctt cccactcttc ccttctcttc
1741 tgcctccggg cgggtctctt cacttaagga ccaagggtctg atcagttcta gggaccaatg
1801 gcccacagtc ctgtgcagga tcttcaaagt cttccatcta attgtgccct ctctcccca
1861 cagcttcaag acccaacggg gcaaggaggt ctgtgctgac ccaaggaga gatgggtcag
1921 ggattccatg aagcatctgg accaaatatt tcaaaatctg aagccatgag ccttcataca
1981 tggactgaga gtcagagctt gaagaaaagc ttatttattt tcccaacctt cccccagggt
2041 cagtgtgaca ttattttatt ataacatcca caaagagatt atttttaaat aatttaaaagc
2101 ataattttt ttaaaaagta ttaattata ttaagttgt tgatgtttta actctatctg
2161 tcatacatcc tagtgaatgt aaaatgcata atcctgggtg tgtgtttttt gttttgtgtt
2221 tctgtgagc tcaactaagt tcacggcaaa atgtcattgt tctcctctct acctgtctgt
2281 agtgtgttgg ggtcctccca tggatcatca aggtgaaaca ctttggtatt ctttggaat
2341 cagtgtcctt gtaagtcaaa tgtgtgcttt gtactgtctg tgtgaaatt gatgttactg

```

```

2401 tatataacta tggaattttg aaaaaaaatt tcaaaaagaa aaaaatatat ataatttaac
2461 actacttagt cttattcttc ttggggtaac atttagctgg gaggtagttt tgggcatcat
2521 gggtagacagt ttgggcatgg acggggccatt tttcaagaat gtcttctggc tacgctggac
2581 tcaaccaagg ttctcagaga acttggtggg accaggccag gatgttccag ctctctgact
2641 ctagtcctta acttcagcag ccctgattcg ctagcctctc ttgtttctct tgtttatata
2701 ttatccagcc taaggatatt tggtatagct gcccataaag actaagataa tctccatcac
2761 tctaccccca accccaatcc caagaacttg caagcatcca tttaaaggcg tggaacctct
2821 tctttttgac agccttttaa ggtcaagatt ccctgtact tagtgagctt agctgaatct
2881 tcttacaac atgtgaccgc ccatattgag ccatacatac cgagcttatt attttccag
2941 cttattggga aaacacgtct aaggcaaca aatttattgt actgttgaac c
//LOCUS      HSY16645      1368 bp      mRNA      PRI      25-SEP-1998
DEFINITION   Homo sapiens mRNA for monocyte chemotactic protein-2.
ACCESSION    Y16645
NID          g2916795
KEYWORDS     MCP-2 gene; monocyte chemotactic protein 2.
SOURCE       human.
ORGANISM     Homo sapiens
              Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
              Primates; Catarrhini; Hominidae; Homo.
REFERENCE    1 (bases 1 to 1368)
AUTHORS      Van Coillie,E.
TITLE        Functional comparison of two human monocyte chemotactic
protein-2
              isoforms, role of the amino-terminal pyroglutamic acid and
              processing by CD26/dipeptidyl peptidase IV
JOURNAL      Biochemistry 37, 12672-12680 (1998)
REFERENCE    2 (bases 1 to 1368)
AUTHORS      Van Coillie,E.
TITLE        Direct Submission
JOURNAL      Submitted (23-FEB-1998) E. Van Coillie, Rega Institute for
Medical
              Research, Minderbroedersstraat 10, 3000 Leuven, BELGIUM
COMMENT      Related sequences: X99886, Y10802.
FEATURES
  source      1..1368
              /organism="Homo sapiens"
              /db_xref="taxon:9606"
              /chromosome="17"
              /tissue_type="testis"
              /clone_lib="Clontech"
              /clone="HL1142q"
              /map="ql1.2"
  gene        473..772
              /gene="MCP-2"
  sig_peptide 473..541
              /gene="MCP-2"
  CDS         473..772
              /gene="MCP-2"
              /codon_start=1
              /product="monocyte chemotactic protein-2"
              /db_xref="PID:e1253690"
              /db_xref="PID:g2916796"

/translation="MKVSAALLCLLLMAATFSPQGLAQPDVSIPITCCFNVINRKIP
IQRLESYTRITNIQCPKEAVIFKTKRGKEVCADPKERWVRDSMKHLDQIFQNLKP"
  mat_peptide 542..769
              /gene="MCP-2"
  variation    677
              /gene="MCP-2"
              /note="polymorphism, Lys -> Gln"
              /replace="c"
BASE COUNT   457 a      292 c      243 g      376 t
ORIGIN
  1 atccattgtg ctctaaagtg atggagagca ccagcaaagc cttagggccc atccctggcc
  61 tcctgttacc cacagagggg taggccttg gctctcttc actatgacgt cagcttccat
  121 tcttctcttc ttatagacaa ttttccattt caaggaaatc agagccctta atagttcagt
  181 gaggtcactt tgctgagcac aatccatac ccttcagcct ctgctccaca gaggcctaagc
  241 aaaagataga aactcacaac ttccttgttt tggtatctgg aaattatccc aggatctggg
  301 gcttactcag catattcaag gaaggtctta cttcattctt ccttgattgt gaccatgccc
  361 aggcctctctg ctccctataa aaggcaggca gagccaccga ggagcagaga ggttgagaac

```



```

421 aacccagaaa ctttcacctc tcatgctgaa gctcacaccc ttgccctcca agatgaaggt
481 ttctgcagcg cttctgtgcc tgctgctcat ggcagccact ttcagccctc agggacttgc
541 tcagccagat tcagtttcca ttccaatcac ctgctgcttt aacgtgatca ataggaaaat
601 tcctatccag aggtctggaga gctacacaag aatcaccaac atccaatgtc ccaaggaagc
661 tgtgatcttc aagaccaaac ggggcaagga ggtctgtgct gacccaagg agagatgggt
721 cagggattcc atgaagcatc tggaccaaat atttcaaaat ctgaagccat gagccttcat
781 acatggactg agagtcagag cttgaagaaa agcttattta ttttcccaa cctccccag
841 gtgcagtgtg acattatttt attataacat ccacaaagag attattttta aataatttaa
901 agcataatat ttcttaaaaa gtatttaatt atatttaagt tgttgatgtt ttaactctat
961 ctgtcatata tcctagttaa tgtaaaaatgc aaaatcctgg tgatgtgttt ttgtttttg
1021 ttttcctgtg agctcaacta agttcacggc aaaatgtcat tgttctcct cctacctgtc
1081 tgtagtgttg tggggtcctc ccatggatca tcaagggtgaa acactttggt attctttggc
1141 aatcagtgtc cctgtaagtc aaatgtgtgc ttgtactgct tgttgttgaa attgatgtta
1201 ctgtatataa ctatggaatt ttgaaaaaaa atttcaaaaa gaaaaaata tatataattt
1261 aaaactaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
1321 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
//LOCUS      HSMCP3A      1085 bp      DNA      PRI      25-JUL-1994
DEFINITION   H.sapiens MCP-3 mRNA for monocyte chemotactic protein-3.
ACCESSION    X72308 S57464
NID          g313707
KEYWORDS     monocyte chemotactic protein 3.
SOURCE       human.
ORGANISM     Homo sapiens
              Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
              Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE    1 (bases 1 to 1085)
AUTHORS      Opdenakker,G., Froyen,G., Fiten,P., Proost,P. and Van Damme,J.
TITLE        Human monocyte chemotactic protein-3 (MCP-3): molecular cloning
of           the cDNA and comparison with other chemokines
JOURNAL      Biochem. Biophys. Res. Commun. 191 (2), 535-542 (1993)
MEDLINE      93213290
REFERENCE    2 (bases 1 to 1085)
AUTHORS      Opdenakker,G.M.
TITLE        Direct Submission
JOURNAL      Submitted (27-MAY-1993) G.M. Opdenakker, Rega Institute,
University   of Leuven, Minderbroedersstraat 10, B-3000 Leuven, BELGIUM
REFERENCE    3 (bases 1 to 1085)
AUTHORS      Opdenakker,G., Fiten,P., Nys,G., Froyen,G., Van Roy,N.,
              Speleman,F., Laureys,G. and Van Damme,J.
TITLE        The human MCP-3 gene (SCYA7): cloning, sequence analysis, and
              assignment to the C-C chemokine gene cluster on chromosome
              17q11.2-q12
JOURNAL      Genomics 21 (2), 403-408 (1994)
MEDLINE      94375065
FEATURES
  source      Location/Qualifiers
              1..1085
              /organism="Homo sapiens"
              /db_xref="taxon:9606"
  gene        299..810
              /gene="MCP-3"
  CDS         299..628
              /gene="MCP-3"
              /codon_start=1
              /product="monocyte chemotactic protein-3"
              /db_xref="PID:g313708"
              /db_xref="SWISS-PROT:P80098"

/translation="MWKPMPSPSNMKASAALLCLLLTAAAFSPQGLAQPVGINTSTTC
CYRFINKKIPKQRLESYRRTTSSHCPREAVIFKTKLDKEICADPTQKWVQDFMKHLDK
              KTQTPKL"
  sig_peptide 299..397
              /gene="MCP-3"
  mat_peptide 398..625
              /gene="MCP-3"
              /product="monocyte chemotactic protein-3"
  polyA_signal 806..810
              /gene="MCP-3"
BASE COUNT   314 a      214 c      229 g      328 t

```

ORIGIN

```

1 ggtttctatt gacttgggtt aatcgtgtga ccgcggtggc tggcacgaaa ttgaccaacc
61 ctgggggttag tatagcttag ttaaactttc gtttattgct aaaggttaat cactgctggt
121 tcccgtagggg gtgtgggctag gctaagcgtt ttgagctgca ttgctgcgtg cttgatgctt
181 gtcccttttg atcgtgggtga tttagagggt gaactcactg gaatggggat gcttgcattg
241 gtaatcttac taagagctaa tagaaaggct aggaccaaac cagaaacctc caattctcat
301 gtggaagccc atgcccctcac cctccaacat gaaagcctct gcagcacttc tgtgtctgct
361 gctcacagca gctgctttca gccccagggt gcttgctcag ccagttggga ttaatacttc
421 aactacctgc tgctacagat ttatcaataa gaaaatccct aagcagaggtc tggagagcta
481 cagaaggacc accagtagcc actgtccccg ggaagctgta atcttcaaga ccaacttgga
541 caaggagatc tgtgctgacc ccacacagaa gtgggtccag gactttatga agcacttgga
601 caagaaaacc caaactccaa agctttgaac attcatgact gaactgaaaa caagccatga
661 cttgagaaac aaataatttg tataccctgt cctttctcag agtggttctg agattatttt
721 aatctaattc taaggaatat gagctttatg taataatgtg aatcatggtt tttcttagta
781 gattttaaaa gttattaata ttttaattta atcttccatg gattttgggt gggtttgaac
841 ataaagcctt ggatgtatat gtcattctcag tgctgtaaaa actgtgggat gctcctcctt
901 tctctacctc atgggggttat tgtataagtc cttgcaagaa tcagtgcata gatttgcttt
961 aattgttaag atatgatgtc cctatggaag catattgtta ttatataatt acatatttgc
1021 atatgtatga ctcccaaatt ttcacataaa atagattttt gtataacaaa aaaaaaaaaa
1081 aaaaa

```

//

```

LOCUS      HSMCP3A      1085 bp      DNA      PRI      25-JUL-1994
DEFINITION H.sapiens MCP-3 mRNA for monocyte chemotactic protein-3.
ACCESSION  X72308 S57464
NID        g313707
KEYWORDS   monocyte chemotactic protein 3.
SOURCE      human.
ORGANISM    Homo sapiens
             Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
             Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1 (bases 1 to 1085)
AUTHORS     Opdenakker,G., Froyen,G., Fiten,P., Proost,P. and Van Damme,J.
TITLE       Human monocyte chemotactic protein-3 (MCP-3): molecular cloning
of          the cDNA and comparison with other chemokines
JOURNAL     Biochem. Biophys. Res. Commun. 191 (2), 535-542 (1993)
MEDLINE     93213290
REFERENCE   2 (bases 1 to 1085)
AUTHORS     Opdenakker,G.M.
TITLE       Direct Submission
JOURNAL     Submitted (27-MAY-1993) G.M. Opdenakker, Rega Institute,
University  of Leuven, Minderbroedersstraat 10, B-3000 Leuven, BELGIUM
REFERENCE   3 (bases 1 to 1085)
AUTHORS     Opdenakker,G., Fiten,P., Nys,G., Froyen,G., Van Roy,N.,
             Speleman,F., Laureys,G. and Van Damme,J.
TITLE       The human MCP-3 gene (SCYA7): cloning, sequence analysis, and
             assignment to the C-C chemokine gene cluster on chromosome
             17ql1.2-ql2
JOURNAL     Genomics 21 (2), 403-408 (1994)
MEDLINE     94375065
FEATURES    Location/Qualifiers
             source          1..1085
                           /organism="Homo sapiens"
                           /db_xref="taxon:9606"
             gene            299..810
                           /gene="MCP-3"
             CDS              299..628
                           /gene="MCP-3"
                           /codon_start=1
                           /product="monocyte chemotactic protein-3"
                           /db_xref="PID:g313708"
                           /db_xref="SWISS-PROT:P80098"

/translation="MWKPMPSPSNMKASALLCLLLTAAAFSPQGLAQPVGINTSTTC
CYRFINKKIPKQRLESYRRTTSSHCPREAVIFKTKLDKEICADPTQKWVQDFMKHLDK
KTQTPKL"
sig_peptide 299..397
             /gene="MCP-3"
mat_peptide 398..625

```

```

                /gene="MCP-3"
                /product="monocyte chemotactic protein-3"
polyA_signal    806..810
                /gene="MCP-3"
BASE COUNT      314 a    214 c    229 g    328 t
ORIGIN
1  gggtttctatt gacttgggtt aatcgtgtga ccgcggtggc tggcacgaaa ttgaccaacc
61 ctgggggttag tatagcttag ttaaactttc gtttattgct aaagggtaat cactgctgtt
121 tcccgtgggg gtgtggctag gctaagcgtt ttgagctgca ttgctgcgtg cttgatgctt
181 gtcccttttg atcgtggtga tttagagggt gaactcactg gaatggggat gcttgcattg
241 gtaatcttac taagagctaa tagaaaaggct aggaccaaac cagaaacctc caattctcat
301 gtggaagccc atgccctcac cctccaacat gaaagcctct gcagcacttc tgtgtctgct
361 gctcacagca gctgctttca gccccagggt gcttgctcag ccagttggga ttaatacttc
421 aactacctgc tgctacagat ttatcaataa gaaaatccct aagcagaggtc tggagagcta
481 cagaaggacc accagtagcc actgtccccg ggaagctgta atcttcaaga ccaaactgga
541 caaggagatc tgtgctgacc ccacacagaa gtgggtccag gactttatga agcacctgga
601 caagaaaacc caaactccaa agctttgaac attcatgact gaactgaaaa caagccatga
661 cttgagaaac aaataatttg tataacctgt cctttctcag agtggttctg agattatttt
721 aatctaattc taagggaatat gagctttatg taataatgtg aatcatgggt tttcttagta
781 gattttaaaa gttattaata ttttaattta atcttccatg gattttgggt ggttttgaac
841 ataagacctt ggatgtatat gtcattccag tgctgtaaaa acttgggag gctctccct
901 tctctacctc atgggggtat tgtataagtc cttgcaagaa tcagtgcata gatttgctt
961 aattgttaag atatgatgtc cctatggaag catattgtta ttatataatt acatatattg
1021 atatgtatga ctcccaaatt ttcacataaa atagattttt gtataacaaa aaaaaaaaaa
1081 aaaaaa
//LOCUS      HSU46767      825 bp      mRNA      PRI      16-DEC-1996
DEFINITION   Human monocyte chemoattractant protein-4 precursor (MCP-4)
mRNA,
complete cds.
ACCESSION    U46767
NID          gl732122
KEYWORDS
SOURCE       human.
ORGANISM     Homo sapiens
              Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
              Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE    1 (bases 1 to 825)
AUTHORS      Garcia-Zepeda, E.A., Combadiere, C.C., Rothenberg, M.E.,
Sarafi, M.N.,
              Lavigne, F., Hamid, Q., Murphy, P. and Luster, A.D.
TITLE        Human monocyte chemoattractant Protein (MCP)-4: A novel CC
              chemokine with activities on monocytes, eosinophils, and
basophils
              induced in allergic and non-allergic inflammation that signals
              through the CC chemokine receptors CCR-2 and 3
JOURNAL      J. Immunol. 158 (1996) In press
REFERENCE    2 (bases 1 to 825)
AUTHORS      Garcia-Zepeda, E.A. and Luster, A.D.
TITLE        Direct Submission
JOURNAL      Submitted (22-JAN-1996) Eduardo A. Garcia-Zepeda, Infectious
              Disease Unit, Massachusetts General Hospital, 149 13th St.,
              Charlestown, MA 02129, USA
FEATURES
source       Location/Qualifiers
              1..825
              /organism="Homo sapiens"
              /db_xref="taxon:9606"
              /tissue_type="heart"
              /clone_lib="EG3.16"
sig_peptide  34..102
              /gene="MCP-4"
CDS          34..330
              /gene="MCP-4"
              /note="small cytokine; intercrine/chemokine; C-C
subfamily    signature; chemoattractant for monocytes, eosinophils"
              /codon_start=1
              /product="monocyte chemoattractant protein-4
precursor"   /db_xref="PID:gl732123"
/translation="MKVSAVLLCLLLMTAAFPNQGGLAQPDALNVPSTCCFTFSSKKIS

```

LQRLKSYVITTSRCPQKAVIFRTKLGKEICADPKEKWVQNYMKHLGRKAHTLKT"

gene 34..330
/gene="MCP-4"
mat_peptide 103..327
/gene="MCP-4"

BASE COUNT 221 a 175 c 185 g 244 t
ORIGIN

```

1 acattgtgaa atctccaact cttaaccttc aacatgaaag tctctgcagt gcttctgtgc
61 ctgctgctca tgacagcagc ttccaacccc cagggacttg ctcagccaga tgcactcaac
121 gtcccatcta cttgctgctt cacatcttagc agtaagaaga tctccttgca gaggtgaag
181 agctatgtga tcaccaccag caggtgtccc cagaaggctg tcatcttcag aaccaactg
241 ggcaaggaga tctgtgctga cccaaaggag aagtgggtcc agaattatat gaaacacctg
301 ggccggaag ctcacaccct gaagacttga actctgctac ccctactgaa atcaagctgg
361 agtacgtgaa atgacttttc cattctcctc tggcctcctc ttctatgctt tggataactt
421 ctaccataat tttcaaatag gatgcattcg gttttgtgat tcaaaatgta ctatgtgtta
481 agtaaatatt gctattattt gacttggtgc tggtttgag tttatttgag tattgctgat
541 cttttctaaa gcaaggcctt gagcaagtag gttgctgtct ctaagccccc ttcccttcca
601 ctatgagctg ctggcagtggt gttgtattcg gttcccaggg gttgagagca tgcctgtggg
661 agtcatggac atgaagggat gctgcaatgt aggaaggaga gctctttgtg aatgtgaggt
721 tgttgctaaa ttattgttta ttgtggaaag atgaatgcaa tagtaggact gctgacattt
781 tgcagaaaat acattttatt taaaatctcc taaaaaaaaa aaaaa

```

//LOCUS HSAMAC1 803 bp RNA PRI 10-AUG-1997
DEFINITION Homo sapiens mRNA for alternative activated macrophage specific
CC

chemokine 1.
ACCESSION Y13710
NID g2326515
KEYWORDS AMAC-1 gene; CC-chemokine 1.
SOURCE human. ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;

Hominidae;

Homo.
REFERENCE 1 (bases 1 to 803)
AUTHORS Politz,O.
TITLE Direct Submission
JOURNAL Submitted (10-JUN-1997) Politz O., Dermatology, Free University
Benjamin Franklin Medical Center, Hindenburgdamm 30; 12200
Berlin

GERMANY
REFERENCE 2 (bases 1 to 803)
AUTHORS Kodelja,V., Mueller,C., Politz,O., Hakiy,N., Orfanos,C.E. and
Goerd,S.
TITLE Cloning of alternative activated macrophage associated CC
chemokine

1 (AMAC-1)
JOURNAL Unpublished
FEATURES Location/Qualifiers
source 1..803
/organism="Homo sapiens"
/db_xref="taxon:9606"
/cell_type="macrophage"
sig_peptide 71..133
/gene="amac-1"
CDS 71..340
/gene="amac-1"
/note="macrophage specific"
/codon_start=1
/product="CC-chemokine 1"
/db_xref="PID:e321838"
/db_xref="PID:g2326516"

/translation="MKGLAAALLVLVCTMALCSAQVGTNKLCLVYTSWQIPQKFI
VDYSETSPQCPKPGVILLTKRGRQICADPNKKWVQKYISDLKLNA"
gene 71..340
/gene="amac-1"
mat_peptide 134..337
/gene="amac-1"

BASE COUNT 214 a 213 c 160 g 216 t
ORIGIN

```

1  ccggcacgag  aggagttgtg  agtttccaag  cccagctca  ctctgaccac  ttctctgcct
61  gcccagcatc  atgaagggcc  ttgcagctgc  cctccttgtc  ctctgtctgca  ccattggccct
121  ctgctcctgt  gcacaagttg  gtaccaacaa  agagctctgc  tgctctgtct  atacctctgt
181  gcagattcca  caaaagtcca  tagttgacta  ttctgaaacc  agccccagct  gcccgaagcc
241  aggtgtcatc  ctcctaacca  agagaggccg  gcagatctgt  gctgacccca  ataagaagtg
301  ggtccagaaa  tacatcagcg  acctgaagct  gaatgcctga  ggggcctgga  agctgcgagg
361  gcccagtcaa  cttggtgggc  ccaggaggga  acaggagcct  gagccagggc  aatggccctg
421  ccaccctgga  ggccacctct  tctaagagtc  ccatctgcta  tgcccagcca  cattaactaa
481  ctttaatctt  agtttatgca  tcatatttca  ttttgaaatt  gatttctatt  gttgagctgc
541  attatgaaat  tagtattttc  tctgacatct  catgacattg  tctttatcat  cctttccctt
601  ttcccttcaa  ctcttcgtac  attcaatgca  tggatcaatc  agtgtgatta  gctttctcag
661  cagcattgtg  gccatatgta  tcaaatgaca  aatctttatt  gaatggtttt  gctcagcacc
721  accttttaat  atattggcag  tacttattat  ataaaaggta  aaccagcatt  ctactgtga
781  aaaaaaaaaa  aaaaaaaaaa  aaa

```

```

//
LOCUS      HUMLD78A      3176 bp      DNA      PRI      17-JAN-1992
DEFINITION Human LD78 alpha gene.
ACCESSION  D90144
NID        g219905
KEYWORDS   LD78; LD78 alpha; cytokine; inducible gene family; secreted
           peptide.
SOURCE     Human blood lymphocyte DNA, clone Lm LD-3.
ORGANISM   Homo sapiens
           Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
           Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE  1 (bases 1 to 3176)
AUTHORS   Nakao,M., Nomiyama,H. and Shimada,K.
TITLE     Structures of human genes coding for cytokine LD78 and their
           expression
JOURNAL    Mol. Cell. Biol. 10 (7), 3646-3658 (1990)
MEDLINE    90287155
COMMENT    These data kindly submitted in computer readable form by:
Hisayuki

```

```

Nomiyama
Department of Biochemistry
Kumamoto University Medical School
2-2-1 Honjo, Kumamoto 860
Japan
Phone: 096-344-2111
Fax: 096-372-6140.
FEATURES             Location/Qualifiers
     source            1..3176
                       /organism="Homo sapiens"
                       /db_xref="taxon:9606"
     TATA_signal        1041..1045
     exon               1069..1227
                       /number=1
     prim_transcript    1069..2957
                       /note="LD78 alpha mRNA and introns"
     sig_peptide        1155..1220
                       /note="LD78 alpha signal peptide"
     CDS                join(1155..1227,1916..2030,2451..2541)
                       /codon_start=1
                       /product="LD78 alpha precursor"
                       /db_xref="PID:d1014875"
                       /db_xref="PID:g219906"

```

```

/translation="MQVSTAALAVLLCTMALCNQFSASLAADTPTACCFSYTSRQIPQ
NFIADYFETSSQCSKPGVIFLTKRSRQVCADPSEEWVQKYVSDLELSA"
     mat_peptide       join(1218..1227,1916..2030,2451..2538)
                       /partial
                       /note="LD78 alpha mature peptide"
     intron            1228..1915
     exon              1916..2030
                       /partial
                       /number=2
     intron            2031..2450
     exon              2451..2957
                       /number=3

```

```

BASE COUNT      833 a    741 c    752 g    850 t
ORIGIN
1  acccagggac  ctatcacaca  aatataagaa  ctattcattc  ttttaaggcat  gtattttccaa
61  gcctttgtat  ttttttccat  gcttaggggt  ggcaagggaat  atatatatat  ttgtacaaat
121  atatatgtgt  atatgtacaa  atacatgtat  atatagtaca  aatatatata  tatattttga
181  caattcttca  gactttgtag  aattttgtata  atgtcgtatc  ttgtcttttt  taaccactga
241  tgttataagc  atatttatgc  cacttcattc  attttagaga  cttaataata  aatgatctag
301  tggataattt  atcattccct  gatggagaaa  aatttagctt  tgtttatatt  agagttataa
361  acgatgctgg  gtcagggtatc  tttatgtttg  aagatggctc  catatttggg  ttgtttccac
421  agaactcttt  cctagaaatg  ctttttctag  gttaatggct  acagatattt  ctaggcacct
481  gacatattga  caccacacct  taaagtattt  ttatgatcca  caactagcgt  ttaacacagc
541  gccctagtca  ctacatgact  aataaataga  caaatgactg  aaacatgacc  tcatgttttc
601  tattctccca  gctttcattc  agttctttgc  ctctgggagg  aggaagggtt  gtgcagccct
661  ccacagcatc  agcccatcaa  ccctatccct  gtggttatag  cagctgagga  agcagaattg
721  cagctctgtg  ggaaggaatg  gggcttgaga  gttcatgcac  agaccagttc  ttatgagaag
781  ggaatgacta  agaatagcct  tgggttgaca  tataccctc  ttcacactca  caggagaaac
841  catttcccta  tgaactata  acaagtcagt  agttgagagc  tgagagttag  agaataagctc
901  aaagatgcta  ttcttgata  tcctgagccc  ctgtggtcac  cagggaccct  gagttgtgca
961  acttagcatg  acagcatcac  tacgcttaaa  aatttccctc  ctacccccc  gattccattt
1021  ccccatccgc  cagggtctgc  tataaaggag  agagctgggt  tcagacttca  gaaggacacg
1081  ggcagcagac  agtggctcgt  cctttcttgg  ctctgctgac  actcgagccc  acattccgtc
1141  acctgctcag  aatcatgcag  gtctccactg  ctgcccttgc  tgtcctcctc  tgcaccattg
1201  ctctctgcaa  ccagttctct  gcctcacgtg  agtctgagtt  tcgttgtggg  taccaccact
1261  ctctggccat  ggttagacca  catcaatctt  ttcttgtggc  ctaaaagccc  ccaagagaaa
1321  agagaacttc  ttaaagggtc  gccaaacatc  ttggctcttc  tctttaagac  ttttattttt
1381  atctctagaa  ggggtcttag  ccccttagtc  tccaggtagc  agaacttagg  caggggcagg
1441  ggagttacag  tcccttttac  agatagaaaa  acagggttcg  aaacgaatca  gttagcaaga
1501  ggcagaatcc  agggctgctt  acttcccagt  ggggtatggt  gttcactctc  cagctcactc
1561  taggtctccc  aggagctctg  tcccttggat  gtcttatgag  agatgtccaa  ggcttctctt
1621  ggggttgggt  atgacttctt  gaaccagaca  aaattccctg  aagagaactg  agataagaga
1681  acagtcctgt  cagggtatct  gatcacacag  agaaacagag  aaccacttat  gaagagtcaa
1741  ggagaagaaa  ggatacagac  agaaacaaag  agacatttct  cagcaaaaaa  gcccaaatgc
1801  ctctcagtc  cttggtctga  gcaagcctgc  cttcctcaac  tgctcgggga  tcagaagctg
1861  cctggccttt  tcttctgagc  tgtgactcgg  gctcattctc  ttcttttctc  cacagttgct
1921  gctgacacgc  cgaccgcctg  ctgcttcagc  tacacctccc  ggcagattcc  acagaatttc
1981  atagctgact  actttgagac  gagcagccag  tgctccaagc  ccggtgtcat  gtaagtgcga
2041  gtcttctctg  tcacctctat  ggaggtaggg  aggggtcagg  ttggggcaga  gacaggccag
2101  aaggctatcc  tggaaaggcc  cagccttcag  gagcctatcg  gggatacagg  acgcagggct
2161  ccgaggtgtg  acctgacttg  gagctggagt  gaggcattgt  ttacagagtc  aggaagggtc
2221  cccccagccc  agaggaaaag  gacaggaaga  aggaggcagc  gggacactct  gagggccacc
2281  cctactgagt  cactgagaga  agctctctag  acagagatag  gcaggggggc  cctgaaagag
2341  gagcaagccc  tgagctgccc  aggacagaga  gcagaatggt  ggggccatgg  tgggccagg
2401  attccctgc  tggattcccc  agtgcttaac  tcttccacag  ttctccacag  cttcctaacc
2461  aagcgaagcc  ggcaggtctg  tgcagacccc  agtgaggagt  ggggtccagaa  atatgtcagc
2521  gacctggagc  tgagtgcctg  aggggtccag  aagcttcgag  gcccagcgac  ctcggtgggc
2581  ccagtgggga  ggagcaggag  cctgagcctt  ggggaacatg  gtgtgacctc  cacagctacc
2641  tctctatgg  actggttgtt  gccaaacagc  cacactgtgg  gactcttctt  aacttaatt
2701  ttaatttatt  tatactattt  agtttttgta  atttattttc  gatttcacag  tgtgtttgtg
2761  attgtttgct  ctgagagttc  cctgttcccc  tcccccttcc  ctacacccgc  gtctggtgac
2821  aaccgagtgg  ctgtcatcag  cctgtgtagg  cagtcattgg  accaaagcca  ccagactgac
2881  aaatgtgtat  cggatgcttt  tgttcagggc  tgtgatcggc  ctggggaaat  aataaagatg
2941  ctcttttaaa  aggtaaacca  gtattgagtt  tggttttgtt  tttctggcaa  atcaaaatca
3001  ctggttaaga  ggaatcatag  gcaaagatta  ggaagaggtg  aaatggaggg  aaattgggag
3061  agatggggag  ggctaccaca  gagtattcca  ctttacaacg  gagacacagt  tctggaacat
3121  tgaactacg  aatatgttat  aactcaaact  ataactgca  tgctctagga  gaattc

```

```

//
LOCUS      AF043339      225 bp      mRNA      PRI      23-FEB-1998
DEFINITION Homo sapiens macrophage inflammatory protein 1 alpha (MIP1a)
mRNA,
partial cds.
ACCESSION  AF043339
NID        g2905627
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 225)
AUTHORS    Jang, J.S. and Kim, B.E.
TITLE      Direct Submission
JOURNAL    Submitted (15-JAN-1998) Protein Engineering, General Institute

```

of Technology, Hyundai Pharm. Ind. Co., Ltd., 213 Sosa Bon 1-dong, Sosa-gu, Bucheon 422-231, Korea

COMMENT forward primer (5'-tgcgcatcacttgctgctgaca-3')
reverse primer (5'-cttctggaccctcaggcact-3').

FEATURES Location/Qualifiers
source 1..225
/organism="Homo sapiens"
/db_xref="taxon:9606"
/cell_type="PHA-treated peripheral blood leukocyte"
gene <1..225
/gene="MIP1a"
primer_bind <1..19
/gene="MIP1a"
/PCR_conditions="94C-1min, 50C-1min, 72C-3min, 30
cycles;
DeltaCycler II from Ericomp"
CDS <1..213
/gene="MIP1a"
/function="CC chemokine"
/function="proinflammatory cytokine involved in
inflammation"
/note="8-10 kDa"
/codon_start=1
/product="macrophage inflammatory protein 1 alpha"
/db_xref="PID:g2905628"

/translation="ASLAADTPTACCFSTSRQIPQNFADYFETSSQCSKPGVIFLT
KRSRQVCADPSEEWVQKYVSDLELSA"
primer_bind complement(205..225)
/gene="MIP1a"

BASE COUNT 50 a 68 c 62 g 45 t
ORIGIN
1 gcacacttg ctgctgacac gccgaccgcc tgctgcttca gctacacctc ccggcagatt
61 ccacagaatt tcatagctga ctactttgag acgagcagcc agtgctccaa gcccgggtgctc
121 atcttcttaa ccaagcgaag ccggcagggtc tgtgctgacc ccagtggagga gtgggtccag
181 aaatgtgtca gcgacctgga gctgagtgcc tgaggggtcc agaag

//
LOCUS HUMLD78B 3112 bp DNA PRI 17-JAN-1992
DEFINITION Human LD78 beta gene.
ACCESSION D90145
NID g219907
KEYWORDS LD78; LD78 beta; cytokine; inducible gene family; secreted
peptide.
SOURCE Human placenta DNA, clone Lm LD-1.
ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae;
Homo.
REFERENCE 1 (bases 1 to 3112)
AUTHORS Nakao,M., Nomiya,H. and Shimada,K.
TITLE Structures of human genes coding for cytokine LD78 and their
expression
JOURNAL Mol. Cell. Biol. 10 (7), 3646-3658 (1990)
MEDLINE 90287155
COMMENT These data kindly submitted in computer readable form by:
Hisayuki Nomiya
Department of Biochemistry
Kumamoto University Medical School
2-2-1 Honjo, Kumamoto 860
Japan
Phone: 096-344-2111
Fax: 096-372-6140.

FEATURES Location/Qualifiers
source 1..3112
/organism="Homo sapiens"
/db_xref="taxon:9606"
repeat_unit 498..797
/note="Alu repeat"

```

TATA_signal      1078..1082
prim_transcript  1106..2995
                  /note="LD78 beta mRNA and introns"
exon              1106..1267
                  /note="LD78 beta precursor, coding region of exon 1"
                  /number=1
CDS              join(1192..1267,1953..2067,2488..2578)
                  /partial
                  /codon_start=1
                  /product="LD78 beta precursor"
                  /db_xref="PID:d1014876"
                  /db_xref="PID:g219908"

/translation="MQVSTAALAVLLCTMALCNQVLSAPLAADTPTACCFSTSRQIP
              QNFIADYFETSSQCSKPSVIFLTRGRQVCADPSEEWVQKYVSDLELSA"
sig_peptide      1192..1260
                  /partial
                  /note="LD78 beta signal peptide"
mat_peptide      join(1258..1267,1953..2067,2488..2575)
                  /partial
                  /note="LD78 beta mature peptide"
intron           1268..1952
exon              1953..2067
                  /number=2
intron           2068..2487
exon              2488..2955
                  /number=3

BASE COUNT      756 a    775 c    780 g    801 t
ORIGIN
1  ttagagactt aataataaag gatcttgttg ataatttata attccctgat agagaaaaat
61 ttagctttgc ttatttttaga gttataaatg atgctgggtc aggtatcttt atgtttgaag
121 atggctccat atttgggttg tttccacaga actctttccc agaaatgctt tttctaggtt
181 aatggctaca catattttcta ggcacctgac atactgacac ccacctctaa agtattttta
241 tgatccacaa ctacggttta acacagcgcc ccagtcactc cgagactaat aaatagacaa
301 atgactgaaa cgtgacctca tgctttctat tctccagctt ttcattgagt tctttctctc
361 tgggaggact gggggttgte tagccctcca cagcatcagc ccattgacct tatccttggt
421 gttatagcag ctgaggaagc agaattacag ctctgtggga aggaatgggg ctggagagtt
481 atgcataga ccaattcttt tttttttttt tttttgagat ggagtctcac ttttgtgccc
541 caggctggag tgcaatggca tgatctcagc tcaccacagc cccacctctc tgggttcaag
601 cgattctcct gccctcagcc tcccagtagt ctgggattac aggcattgtg caccacgcct
661 gactactttt gtatttttag tagagatgga gtttctcttt ctgggtcagg ttggtctcaa
721 actcctgacc tcaggtgatc cgcagcctcg gcctccaaa gtgttgggat tacaggtgtg
781 agcgaccatg cctggctgca tagaccagtt cttatgagaa gggatcaact aagaatagcc
841 ttgggttgac acacaccctt cttcacactc acaggagaaa cccatgaaag ctagaaccag
901 tcatgagttg agagctgaga gttagagagt agctcagaga tgctattctt ggatattctg
961 agccccctgt gtcaccaggg accctgagtt gtgcaacact cagcatgaca gcatcactac
1021 acttaaaaaa ttcctctctc acccccagat tccatttccc catccgccag ggctgcctat
1081 aaagaggaga gatggcttca gacatcagaa ggacgcaggc agcaaagagt agtcagctcc
1141 ttcttggttc tgctgacact cgagcccaca ttccatcacc tgctcccaat catgcagggtc
1201 tccactgctg cccttgccgt cctcctctgc accatggctc tctgcaacca ggctcctctt
1261 gcaccacgtg agtccatggt gttgttgtgg gtatcaccac tctctggcca tgggttagacc
1321 acatcagttt ttttttgctg cctgagagcc ccgaagagaa aagaaggaag ttcttaaagc
1381 gctgcccac accttggtct ttttcttcac aacttttatt tttatctcta gaagggtctt
1441 tagccctcct agtctccagg tatgagaatc taggcagggg caggggagtt acagtcctt
1501 gtacagatag aaaaacaggg ttcaaaacga atcagtttgc aagaggcaga atccagggtt
1561 gcttacttcc cagtgggggtc tgttgttcac tctccagctc accctagggtc tcccaggagc
1621 cctgtccctt ggtgtcttta tgagagatgt ccagggtctc tcttgggctg gggatgactt
1681 tcttgaaccg acaaaattcc atgaagagag ctaagagaac agtccattca ggtatctgga
1741 tcacatagag aaacagagaa cccactatga agagtcaagg ggaagagga atatagacag
1801 aaacaaagag acatttctct gcaaaacccc ccaaagcctt tgagtcactt tggctctgagc
1861 aagcctgccc tctcaacca ctcagggtatc agaagctgcc tggccttttc tcttgagctg
1921 tgactcgggc ttattctctc ctttctccgc agttgtctgt gacacgccga ccgctgctg
1981 cttcagctac acctcccgac agattccaca gaatttcata gctgactact ttgagacgag
2041 cagccagtg cccaagcccc gtgtcatgta agtgccagtc ttctgtctca cctctaggga
2101 ggtaggaggt gtcagggtgg gggcagaaac aggcagaaag gccatcctgg aaggccagc
2161 ccttcaggag cctatcgggg atacaggagc cagggcactg aggtgtgacc tgacttgggg
2221 ctggagtgag gtgggtgtta cagagtcagg aagggtctgc ccaggccaga ggaaggaagc
2281 aggaagaagg aggcagcagg acactctgag ggcccccttg cctggagtcg ctgagagaag
2341 cctctagac ggagataggc agggggcccc tgagagagga gcaggccttg agctgccag
2401 gacagagagc aggatgtcag gccatggttg gccaggatt ccccggtctg attccccagt
2461 gcttaactct tctcccttc tccacagctt cctaaccaag agaggccggc aggtctgtgc

```



```

2521 tgacccagcaggagtgagg tccagaaata cgtcagtgac ctggagctga gtgcctgagg
2581 ggtccagaag cttcgaggcc cagcgacctc agtggggccca gtggggagga gcaggagcct
2641 gagccttggg aacatgcgtg tgacctctac agctacctct tctatggact gggtattgcc
2701 aaacagccac actgtgggac tcttcttaac ttaaatttta atttatttat actatttagt
2761 ttttataatt tatttttgat ttcacagtgt gtttgtgatt gtttgctctg agagttcccc
2821 ctgtccccc caccctccct cacagtgtgt ctggtgacga ccgagtggtc gtcacgggcc
2881 tgtgtaggca gtcattggc caaagccacc agactgacaa atgtgtatca gatgcttttg
2941 ttcagggtcg tgatcgccct ggggaaataa taaagatgtt cttttaaacg gtaaaccagt
3001 attgagtttg gttttgtttt tctggcaaat caaaatcact agttaagagg aatcataggg
3061 aaagattagg aagaggtgaa atggaggagg actgggagag atggggagcg ct

```

```

//
LOCUS      HUMACT2A      696 bp      mRNA      PRI      30-OCT-1994
DEFINITION Human activation (Act-2) mRNA, complete cds.
ACCESSION  J04130
NID        g178017
KEYWORDS   act2 gene; immune activation gene.
SOURCE     Human (Hut-102B2 library) activated T cells, cDNA to mRNA.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 696)
AUTHORS    Lipes,M.A., Napolitano,M., Jeang,K.T., Chang,N.T. and
Leonard,W.J.
TITLE      Identification, cloning, and characterization of an immune
            activation gene
JOURNAL     Proc. Natl. Acad. Sci. U.S.A. 85 (24), 9704-9708 (1988)
MEDLINE     89071764
COMMENT     Draft entry and computer-readable sequence [1] kindly submitted
            by
            W.Leonard, 09-JAN-1989.

```

```

FEATURES             Location/Qualifiers
     source            1..696
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /map="Unassigned"
     mRNA              <1..696
                        /note="act-2 mRNA"
     sig_peptide       109..177
                        /gene="LAG2"
                        /note="act-2 protein signal peptide"
     gene              109..387
                        /gene="LAG2"
     CDS               109..387
                        /gene="LAG2"
                        /note="act-2 protein precursor"
                        /codon_start=1
                        /db_xref="GDB:G00-127-452"
                        /db_xref="PID:g178018"

```

```

/translation="MKLCVTVLSLLMLVAAFCSPLSAPMGSDPPTACCFSTARKLP
            RNFVVDYYETSSSLCSQPAVVFQTKRSKQVCADPSESWSVQEVYDLELN"
     mat_peptide       178..384
                        /gene="LAG2"
                        /note="act-2 protein"

```

```

BASE COUNT      157 a      203 c      139 g      197 t
ORIGIN           Unreported.

```

```

1  tttccccccc ccccccccc ccccgcccga gcacaggaca cagctgggtt ctgaagcttc
61  tgagttctgc agcctcacct ctgagaaaac ctcttttcca ccaataccat gaagctctgc
121 gtgactgtcc tgtctctcct catgctagta gctgccttct gctctccagc gctctcagca
181 ccaatgggct cagaccctcc caccgctgct tgcttttctt acaccgcgag gaagcttctt
241 cgcaactttg tggtagatta ctatgagacc agcagcctct gctccagacc agctgtggta
301 ttccaaacca aaagaagcaa gcaagctctg gctgatccca gtgaatcttg gttccaggag
361 tacgtgtatg acctggaact gaactgagct gctcagagac aggaagtctt cagggaaggt
421 caccgtgagc cggtatgctt tccatgagac acatctcctc catactcagg actcctctcc
481 gcagtctctg tcccttctct taatttaata ttttttatgt gccgtgttat tgtattaggt
541 gtcatttcca ttatttatat tagtttagcc aaaggataag tgtcctatgg ggtgggtcca
601 ctgtcactgt ttctctgctg ttgcaaatat atggataaca catttgattc tgtgtgtttt
661 ccataataaa acttttaaat aaaatgcaga cagtta

```

```

//
LOCUS      AF031587      481 bp      mRNA      PRI      02-JAN-1998
DEFINITION Homo sapiens MIP-1 delta mRNA, complete cds.

```

ACCESSION AF031587
 NID g2739163
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
 Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 481)
 AUTHORS Wang, W.
 TITLE Molecular cloning and characterization of a new CC chemokine
 MIP-1
 delta
 JOURNAL Unpublished
 REFERENCE 2 (bases 1 to 481)
 AUTHORS Wang, W.
 TITLE Direct Submission
 JOURNAL Submitted (27-OCT-1997) Immunobiology, DNAX Research Institute,
 901
 California Ave, Palo Alto, CA 94304, USA
 FEATURES Location/Qualifiers
 source 1..481
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="17"
 CDS 1..342
 /note="CC or beta chemokine"
 /codon_start=1
 /product="MIP-1 delta"
 /db_xref="PID:g2739164"

/translation="MKVSVAAALSCLMLVAVLGSQAQFINDAETELMMSKLPLENPVVL

NSFHFAADCCTSYISQSIPLCSLMKSYFETSSECSKPGVIFLTKKGRQVCAKPSGPGVQ
 DCMKKLKPYSI"

BASE COUNT 140 a 112 c 100 g 129 t
 ORIGIN

1 atgaaggtct cctgtgctgc cctctctgc ctcatgcttg ttgctgtcct tggatcccag
 61 gcccagttca taaatgatgc agagacagag ttaatgatgt caaagcttcc actggaaaat
 121 ccagtagttc tgaacagctt tcactttgct gctgactgct gcacctccta catctcacia
 181 agcatcccggt gttcactcat gaaaagttat tttgaaacga gcagcgagtgc ctccaagcca
 241 ggtgtcatat tcctcaccaa gaaggggagg caagtctgtg ccaaaccag tgggtccggga
 301 gttcaggatt gcatgaaaaa gctgaagccc tactcaatat aataataaag agacaaaaaga
 361 gggcagccac ccacctccaa cacctcctgt gagtttcttg gtctgaaata cttaaaaaat
 421 atatatttg ttgtgtctgg taatgaaagt aatgcatcta ataaagagta ttcaattttt
 481 t

// LOCUS AF043340 234 bp mRNA PRI 23-FEB-1998
 DEFINITION Homo sapiens macrophage inflammatory protein 2 alpha (MIP2a)
 mRNA,

partial cds.
 ACCESSION AF043340
 NID g2905629
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
 Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 234)
 AUTHORS Jang, J.S. and Kim, B.E.
 TITLE Direct Submission
 JOURNAL Submitted (15-JAN-1998) Protein Engineering, General Institute
 of
 Technology, Hyundai Pharm. Ind. Co., Ltd., 213 Sosa Bon 1-dong,
 Sosa-gu, Bucheon 422-231, Korea
 COMMENT forward primer (5'-tgcgacccctggccactgaactg-3')
 reverse primer (5'-ccttccttctggtcagttgga-3').
 FEATURES Location/Qualifiers
 source 1..234
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /cell_type="PHA-treated peripheral blood leukocyte"

```

gene          <1..234
               /gene="MIP2a"
primer_bind   <1..21
               /gene="MIP2a"
               /PCR_conditions="94C-1min, 50C-1min, 72C-3min, 30
cycles;
               DeltaCycler II from Ericomp"
CDS           <1..222
               /gene="MIP2a"
               /function="CXC chemokine"
               /function="proinflammatory cytokine involved in
inflammation"
               /note="8-10 kDa"
               /codon_start=1
               /product="macrophage inflammatory protein 2 alpha"
               /db_xref="PID:g2905630"

/translation="APLATELRCQCLQTLQGIHLKNIQSVKVKSPGPHCAQTEVIATL
               KNGQKACLNPASPMVKIIEKMLKNGKSN"
primer_bind   complement(214..234)
               /gene="MIP2a"
BASE COUNT    74 a      70 c      54 g      36 t
ORIGIN
      1 gcacccctgg ccactgaact gcgctgccag tgcttcgaga ccctgcaggg aattcacctc
     61 aagaacatcc aaagtgtgaa ggtgaagtcc cccggacccc actgcgcccc aaccgaagtc
    121 atagccacac tcaagaatgg gcagaaaagct tgtctcaacc ccgcatcgcc catggttaag
    181 aaaatcatcg aaaagatgct gaaaaatggc aaatccaact gaccagaagg aagg

//
LOCUS          HSU77035      764 bp      mRNA          PRI          23-JAN-1997
DEFINITION     Human macrophage inflammatory protein 3 alpha (MIP-3a) mRNA,
               complete cds.
ACCESSION      U77035
NID            g1790924
KEYWORDS       .
SOURCE         human.
ORGANISM       Homo sapiens
               Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
               Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE      1 (bases 1 to 764)
AUTHORS        Rossi,D.L., Vicari,A.P., Franz-Bacon,K., McClanahan,T.K. and
               Zlotnik,A.
TITLE          Identification through bioinformatics of two new macrophage
               proinflammatory human chemokines: MIP-3alpha and MIP-3beta
JOURNAL        J. Immunol. 158 (3), 1033-1036 (1997)
MEDLINE        97166046
REFERENCE      2 (bases 1 to 764)
AUTHORS        Rossi,D.L. and Zlotnik,A.
TITLE          Direct Submission
JOURNAL        Submitted (31-OCT-1996) Immunology, DNAX Research Institute,
901
               California Ave., Palo Alto, CA 94304, USA
FEATURES       Location/Qualifiers
               source          1..764
                               /organism="Homo sapiens"
                               /db_xref="taxon:9606"
                               /cell_type="elutriated monocytes activated with
                               LPS/IFN-GAMMA"
               gene           1..291
                               /gene="MIP-3a"
               CDS            1..291
                               /gene="MIP-3a"
                               /note="chemokine"
                               /codon_start=1
                               /product="macrophage inflammatory protein 3 alpha"
                               /db_xref="PID:g1790925"

/translation="MCCTKSLLLLAALMSVLLLLHLCGESEAASNFDCCLGYTDRILHPK
               FIVGFTRQLANEGCDINAIIFHTKKKLSVCANPKQTWVKYIVRLLSKVKNM"
BASE COUNT     235 a      121 c      146 g      260 t      2 others
ORIGIN
      1 atgtgctgta ccaagagttt gctcctggct gctttgatgt cagtgtgct actccacctc

```

```

61  tgcggcgaat  cagaagcagc  aagcaacttt  gactgctgtc  ttggatacac  agaccgtatt
121 cttcatccta  aattttattgt  gggcttcaca  cggcagctgg  ccaatgaagg  ctgtgacatc
181 aatgctatca  tctttcacac  aaagaaaaag  ttgtctgtgt  gcgcaaatcc  aaaacagact
241 tgggtgaaat  atattgtgcg  tctcctcagt  aaaaaagtca  agaacatgta  aaaactgtgg
301 cttttctgga  atggaattgg  acatagccca  agaacagaaa  gaaccttgct  ggggttggag
361 gtttcacttg  cacatcatgg  agggtttagt  gcttatctaa  tttgtgcctc  actggacttg
421 tccaattaat  gaagttgatt  catattgcat  catagtttgc  tttgtttaag  catcacatta
481 aagttaaact  gtattttatg  ttatttatag  ctgtaggttt  tctgtgttta  gctatttaat
541 actaattttc  cataagctat  tttggtttag  tgcaaagtat  aaaattatat  ttggggggga
601 ataagattat  atggactttt  ttgcaagcaa  caagctattt  tttaaaamma  actatttaac
661 attcctttgt  ttatattgtt  ttgtctccta  aattgttgta  attgcattat  aaaataagaa
721 aaatattaat  aagacaaata  ttgaaaataa  agaaacaaaa  agtt

```

```

//
LOCUS      HSU77180      545 bp      mRNA      PRI      23-JAN-1997
DEFINITION Human macrophage inflammatory protein 3 beta (MIP-3beta) mRNA,
complete cds.
ACCESSION  U77180
NID        gl791002
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 545)
AUTHORS    Rossi, D.L., Vicari, A.P., Franz-Bacon, K., McClanahan, T.K. and
            Zlotnik, A.
TITLE      Identification through bioinformatics of two new macrophage
            proinflammatory human chemokines: MIP-3alpha and MIP-3beta
JOURNAL     J. Immunol. 158 (3), 1033-1036 (1997)
MEDLINE     97166046
REFERENCE  2 (bases 1 to 545)
AUTHORS    Vicari, A. and Zlotnik, A.
TITLE      Direct Submission
JOURNAL     Submitted (01-NOV-1996) Immunology, DNAX Research Institute,
901         California Ave, Palo Alto, CA 94304, USA
FEATURES   Location/Qualifiers
            source          1..545
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /cell_type="macrophages activated with LPS or IFNg"
                        /chromosome="9"
            gene            1..297
                        /gene="MIP-3beta"
            CDS              1..297
                        /gene="MIP-3beta"
                        /function="chemokine"
                        /codon_start=1
                        /product="macrophage inflammatory protein 3 beta"
                        /db_xref="PID:gl791003"

```

/translation="MALLLALSLLVLWTSPAPTLSGTNDACCLSVTKPIPGYIVR"

NFHYLLIKDGCVRPAVVFTTLRGRQLCAPDQPPVERIIQRLQRTSAKMKRRSS"

BASE COUNT 125 a 160 c 153 g 107 t

```

ORIGIN
1  atggccctgc  tactggccct  cagcctgtgt  gttctctgga  cttccccagc  cccaactctg
61  agtggcacca  atgatgctga  agactgctgc  ctgtctgtga  cccagaaacc  catccctggg
121  tacatcgtga  ggaacttcca  ctaccttctc  atcaaggatg  gctgcagggt  gcctgctgta
181  gtgttcacca  cactgagggg  ccgccagctc  tgtgcacccc  cagaccagcc  ctgggtagaa
241  cgcatcatcc  agagactgca  gaggacctca  gccaaagatga  agcgccgcag  cagttaacct
301  atgaccgtgc  agagggagcc  cggagtccga  gtcaagcatt  gtgaattatt  acctaacctg
361  gggaaccgag  gaccagaagg  aaggaccagg  cttccagctc  ctctgcacca  gacctgacca
421  gccaggacag  ggcctggggg  gtgtgtgagt  gtgagtgtga  gcgagagggt  gagtgtggtc
481  tagagtaaag  ctgctccacc  cccagattgc  aatgctacca  ataaagccgc  ctggtgttta
541  caact

```

```

//
LOCUS      HUMTCSM      1160 bp      mRNA      PRI      15-JUN-1989
DEFINITION Human T cell-specific protein (RANTES) mRNA, complete cds.
ACCESSION  M21121
NID        g339420

```

KEYWORDS Alu repeat; T-cell-specific protein.
SOURCE Human peripheral blood (T lymphocyte) cell line AH2, cDNA to mRNA,
 clone 228.
ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (bases 1 to 1160)
AUTHORS Schall,T.J., Jongstra,J., Dyer,B.J., Jorgensen,J.,
 Clayberger,C.,
 Davis,M.M. and Krensky,A.M.
TITLE A human T cell-specific molecule is a member of a new gene family
JOURNAL J. Immunol. 141, 1018-1025 (1988)
MEDLINE 88285659
COMMENT Draft entry and computer-readable sequence for [1] kindly provided
 by A.M.Krensky, 24-OCT-1988.
FEATURES Location/Qualifiers
 source 1..1160
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 CDS 27..302
 /note="T cell-specific protein precursor"
 /codon_start=1
 /db_xref="PID:g339421"
 /translation="MKVSAARLAVILIATLALCAPASASPYSSDTPCCFAYIARPLPR
 AHIKEYFYTSGKCSNPVVFVTRKNRQVCANPEKKWVREYINSLEMS"
 sig_peptide 27..95
 /note="T cell-specific protein signal peptide"
 mat_peptide 96..299
 /note="T cell-specific protein"
 repeat_region 450..950
 /note="Alu-related repeats"
BASE COUNT 298 a 332 c 295 g 235 t
ORIGIN 276 bp upstream of RsaI site.
 1 cctccgacag cctctccaca ggtaccatga aggtctccgc ggcacgcctc gctgtcatcc
 61 tcattgctac tgccctctgc gctcctgcat ctgcctcccc atattcctcg gacaccacac
 121 cctgctgctt tgccctacatt gcccgcacac tgccccgtgc ccacatcaag gagtatttct
 181 acaccagtgg caagtgtctc aaccacgacg tcgtctttgt caccgaaaag aaccgccaag
 241 tgtgtgccaa cccagagaag aaatgggttc gggagtacat caactctttg gagatgagct
 301 aggatggaga gtccttgaac ctgaacttac acaaatttgc ctgtttctgc ttgctcttgt
 361 cctagcttgg gaggcttccc ctcaactatcc taccacaccc gctccttgaa gggcccagat
 421 tctgaccacg acgagcagca gttacaaaaa cttcccccag gctggacgtg gtggctcagc
 481 cttgtaatcc cagcactttg ggaggccaag gtgggtggat cacttgaggt caggagttcg
 541 agacagcctg gccaacatga tgaaccccca tgtgtactaa aaatacaaaa aattagccgg
 601 gcgtggtagc gggcgctgtg agtccagct actcgggagg ctgaggcagg agaattggct
 661 gaaccgggga gcggagcttg cagtgaagcg agatcgcgcc actgcactcc agcctgggcg
 721 acagagcgag actccgtctc aaaaaaaaaa aaaaaaaaaa aaaaaataca aaaattagcc
 781 gcgtgggtgg ccacgcctgt aatccagct actcgggagg ctaaggcagg aaaattgttt
 841 gaaccagga ggtggaggct gcagtgaagc gagattgtgc cacttcactc cagcctgggt
 901 gacaaagtga gactccgtca caacaacaac aacaaaaagc ttccccaact aaagcctaga
 961 agagcttctg aggcgtgct ttgtcaaaag gaagtctcta ggttctgagc tctggcttgc
 1021 ccttggcttt gcaagggctc tgtgacaagg aaggaagtca gcatgcctct agaggcaagg
 1081 aagggaggaa cactgcactc ttaagcttcc gccgtctcaa cccctcacag gagcttactg
 1141 gcaaacatga aaaatcgggg
 //
LOCUS HUMTLI309 520 bp mRNA PRI 14-JAN-1995
DEFINITION Human secreted protein (I-309) mRNA, complete cds.
ACCESSION M57502
NID g339728
KEYWORDS secreted protein.
SOURCE Human T-cell, cDNA to mRNA.
ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (bases 1 to 520)
AUTHORS Miller,M.D., Hata,S., De Waal Malefyt,R. and Krangel,M.S.
TITLE A novel polypeptide secreted by activated human T lymphocytes
JOURNAL J. Immunol. 143 (9), 2907-2916 (1989)

MEDLINE 90038522

FEATURES

source 1..520
/organism="Homo sapiens"
/db_xref="taxon:9606"
/cell_type="T-cell"
/germline
/map="17"

mRNA <1..520
/gene="SCYA1"
/note="G00-118-872"

gene 1..520
/gene="SCYA1"

CDS 51..341
/gene="SCYA1"
/codon_start=1
/db_xref="GDB:G00-118-872"
/product="secreted protein I-309"
/db_xref="PID:g339729"

/translation="MQIITTAIVCLLLAGMWPEVDVDSKSMQVPPFSRCCFSFAEQEIPL
RAILCYRNTSSICSNEGLIFKLKRGKEACALDTVGWVQRHRKMLRHCPSKRK"

BASE COUNT 140 a 137 c 122 g 121 t

ORIGIN

1 accaggctca tcaaagctgc tccaggaagg cccaagccag accagaagac atgcagatca
61 tcaccacagc cctggtgtgc ttgctgctag ctgggatgtg gccggaagat gtggacagca
121 agagcatgca ggtacccttc tccagatgtt gcttctcatt tgcggagcaa gagattcccc
181 tgagggcaat cctgtgttac agaaatacca gctccatctg ctccaatgag ggcttaatat
241 tcaagctgaa gagaggcaaa gaggcctgcg ccttggaacac agttggatgg gttcagagggc
301 acagaaaaat gctgaggcac tgcccgtcaa aaagaaaatg agcagatttc ttccattgt
361 gggctctgga aaccacatgg cttcacctgt ccccgaaact accagcccta caccattcct
421 tctgccctgc ttttgctagg tcacagagga tctgcttggt cttgataagc tatgttggtg
481 cactttaaac atttaaatta tacaatcatc aacccccaac

//

LOCUS AB000887 687 bp mRNA PRI 05-JUN-1997

DEFINITION Human mRNA for EB11-ligand chemokine, complete cds.

ACCESSION AB000887

NID g2189952

KEYWORDS EB11-ligand chemokine; ELC.

SOURCE Homo sapiens fetal tissue_lib:lung cDNA to mRNA.

ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;

Hominidae;
Homo.

REFERENCE 1 (bases 1 to 687)

AUTHORS Yoshida,R., Imai,T., Hieshima,K., Kusuda,J., Baba,M.,
Kitaura,M., Nishimura,M., Kakizaki,M., Nomiyama,H. and Yoshie,O.

TITLE Direct Submission

JOURNAL Submitted (05-FEB-1997) to the DDBJ/EMBL/GenBank databases.
Hisayuki Nomiyama, Kumamoto University Medical School,

Department of Biochemistry; Honjo 2-2-1, Kumamoto, Kumamoto 860, Japan
(E-mail:nomiyama@gpo.kumamoto-u.ac.jp, Tel:+81-96-373-5063)

REFERENCE 2 (sites)

AUTHORS Yoshida,R., Imai,T., Hieshima,K., Kusuda,J., Baba,M.,
Kitaura,M., Nishimura,M., Kakizaki,M., Nomiyama,H. and Yoshie,O.

TITLE Molecular cloning of a novel human CC chemokine EB11-ligand
chemokine that is a specific functional ligand for EB11, CCR7

JOURNAL J. Biol. Chem. 272 (21), 13803-13809 (1997)

MEDLINE 97298088

FEATURES

source 1..687
/organism="Homo sapiens"
/db_xref="taxon:9606"
/dev_stage="fetal"
/tissue_lib="lung"

gene 139..435
/gene="ELC"

```

CDS                139..435
                   /gene="ELC"
                   /note="CC chemokine"
                   /codon_start=1
                   /product="EBI1-ligand chemokine"
                   /db_xref="PID:d1021215"
                   /db_xref="PID:g2189953"

/translation="MALLLLALLVLTSPAPTLSGTNDACCLSVTQKPIPGYIVR
NFHYLLIKDGCVRPAVVFTTLRGRQLCAPPDQPWVERIIQRLQRTSAKMKRRSS"
mat_peptide        202..432
                   /gene="ELC"
                   /product="EBI1-ligand chemokine"
polyA_signal        657..662
BASE COUNT         154 a    223 c    173 g    137 t
ORIGIN
    1 cattccacgc ctcacatcac tcacaccttg catttcaccc ctgcatccca gtcgccctgc
   61 agcctcacac agatcctgca cacaccaga cagctggcgc tcacacattc accgttggcc
  121 tgccctctgtt caccctccat ggccctgcta ctggccctca gcctgctggt tctctggact
  181 tccccagccc caactctgag tggcaccaat gatgctgaag actgctgcct gtctgtgacc
  241 cagaaaccca tccctgggta catcgtgagg aacttccact accttctcat caaggatggc
  301 tgcagggtgc ctgctgtagt gttcaccaca ctgaggggcc gccagctctg tgcaccccca
  361 gaccagccct gggtagaacg catcatccag agactgcaga ggacctcagc caagatgaag
  421 cgccgcagca gttaacctat gaccgtgcag agggagcccg gagtccgagt caagcattgt
  481 gaattattac ctaacctggg gaaccgagga ccagaaggaa ggaccaggct tccagctcct
  541 ctgcaccaga cctgaccagc caggacaggg cctgggggtg gtgtgagtgt gagtgtgagc
  601 gagagggtga gtgtggtcag agtaaagctg ctccaccccc agattgcaat gctaccaata
  661 aagccgcctg gtgtttacaa ctaattg

//
LOCUS               AB000221       760 bp    mRNA                PRI          31-JUL-1997
DEFINITION          Homo sapiens mRNA for CC chemokine, complete cds.
ACCESSION            AB000221
NID                  g2289718
KEYWORDS             CC chemokine; PARC; pulmonary and activation-regulated
                    chemokine.
SOURCE               Homo sapiens lung cDNA to mRNA.
ORGANISM              Homo sapiens
                    Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
                    Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae;
                    Homo.
REFERENCE            1 (bases 1 to 760)
AUTHORS              Nomiya,H.
TITLE                Direct Submission
JOURNAL              Submitted (04-JAN-1997) to the DDBJ/EMBL/GenBank databases.
                    Hisayuki Nomiya, Kumamoto University Medical School,
Department
                    of Biochemistry; Honjo 2-2-1, Kumamoto, Kumamoto 860, Japan
                    (E-mail:nomiya@gpo.kumamoto-u.ac.jp, Tel:81-96-373-5063,
                    Fax:81-96-372-6140)
REFERENCE            2 (sites)
AUTHORS              Hieshima,K., Imai,T., Baba,M., Shoudai,K., Ishizuka,K.,
                    Nakagawa,T., Tsuruta,J., Takeya,M., Sakaki,Y., Takatsuki,K.,
                    Miura,R., Opdenakker,G., Damme,J., Yoshie,O. and Nomiya,H.
TITLE                A novel human CC chemokine PARC that is most homologous to
                    macrophage-inflammatory protein-1alpha/LD78alpha and
chemotactic
                    for T lymphocytes, but not for monocytes
JOURNAL              J. Immunol. 159 (3), 1140-1149 (1997)
MEDLINE              97376836
FEATURES
    source            Location/Qualifiers
                    1..760
                    /organism="Homo sapiens"
                    /db_xref="taxon:9606"
                    /tissue_type="lung"
    gene              64..333
                    /gene="PARC"
    CDS                64..333
                    /gene="PARC"
                    /note="pulmonary and activation-regulated chemokine"

```

/codon_start=1
 /product="CC chemokine"
 /db_xref="PID:d1022520"
 /db_xref="PID:g2289719"

/translation="MKGLAALLVLVCTMALCSCAQVGTNKLCLVYTSWQIPQKFI
 VDYSETSPQCPKPGVILLTKRGRQICADPNKKWVQKYISDLKUNA"

BASE COUNT 186 a 208 c 155 g 211 t

ORIGIN

```

1 gccaggagtt gtgagtttcc aagccccagc tcactctgac cacttctctg cctgcccagc
61 atcatgaagg gccttgacgc tgccctcctt gtcctcgtct gcaccatggc cctctgctcc
121 tgtgcacaag ttggtaccaa caaagagctc tgctgcctcg tctatacttc ctggcagatt
181 ccacaaaagt tcatagtga ctattctgaa accagcccc agtgcccaa gccagggtgc
241 atcctcctaa ccaagagagg ccggcagatc tgtgctgacc ccaataagaa gtgggtccag
301 aaatacatca ggcacctgaa gctgaatgcc tgaggggctt ggaagctgag agggcccagt
361 gaacttgggt ggcccaggag ggaacaggag cctgagccag ggcaatggcc ctggcaccct
421 ggaggccacc tcttctaaga gtcccatctg ctatgcccag ccacattaac taactttaat
481 cttagtttat gcatcatatt tcattttgaa attgatttct attgttgagc tgcattatga
541 aattagtatt ttctctgaca tctcatgaca ttgtctttat catcctttcc cctttccctt
601 caactcttcg tacattcaat gcatggatca atcagtgtga ttagctttct cagcagacat
661 tgtgccatat gtatcaaatg acaaatcttt attgaatggt ttgtctcagc accacctttt
721 aatatattgg cagtacttat tatataaaag gtaaaccagc

```

//

LOCUS D86955 799 bp mRNA PRI 06-MAR-1997
 DEFINITION Human mRNA for CC chemokine LARC precursor, complete cds.
 ACCESSION D86955
 NID g1871138
 KEYWORDS CC chemokine LARC precursor.
 SOURCE Homo sapiens cDNA to mRNA.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;

Hominidae;

Homo.

REFERENCE 1 (sites)

AUTHORS Hieshima,K., Imai,T., Opdenakker,G., Van Damme,J., Kusuda,J.,
 Tei,H., Sakaki,Y., Takatsuki,K., Miura,R., Yoshie,O. and
 Nomiya,H.

TITLE Molecular cloning of a novel human CC chemokine liver and
 activation-regulated chemokine (LARC) expressed in liver.
 Chemotactic activity for lymphocytes and gene localization on
 chromosome 2

JOURNAL J. Biol. Chem. 272 (9), 5846-5853 (1997)

MEDLINE 97190319

REFERENCE 2 (bases 1 to 799)

AUTHORS Hieshima,K., Imai,T., Opdenakker,G., Van Damme,J., Kusuda,J.,
 Tei,H., Sakaki,Y., Takatsuki,K., Miura,R., Yoshie,O. and
 Nomiya,H.

JOURNAL Unpublished (1996)

REFERENCE 3 (bases 1 to 799)

AUTHORS Nomiya,H.

TITLE Direct Submission

JOURNAL Submitted (08-AUG-1996) to the DDBJ/EMBL/GenBank databases.

Hisayuki Nomiya, Kumamoto University Medical School,

Department

of Biochemistry; Honjo 2-2-1, Kumamoto, Kumamoto 860, Japan
 (E-mail:nomiya@gpo.kumamoto-u.ac.jp, Tel:+81-96-373-5063)

FEATURES

Location/Qualifiers

source

1..799

/organism="Homo sapiens"

/db_xref="taxon:9606"

/chromosome="2"

/map="q33-37"

sig_peptide

59..136

/gene="LARC"

CDS

59..349

/gene="LARC"

/codon_start=1

/product="CC chemokine LARC precursor"

/db_xref="PID:d1013880"

/db_xref="PID:g1871139"


```

/translation="MCCTKSLLLAALMSVLLHLGGESEAASNFDCCLGYTDRILHPK
              FIVGFTRQLANEGCDINAIIFHTKKLSVCANPKQTWVKYIVRLLSKKVKNM"
gene          59..349
              /gene="LARC"
mat_peptide   137..346
              /gene="LARC"
              /product="CC chemokine LARC"
BASE COUNT    240 a    138 c    153 g    268 t
ORIGIN
1 cactcccaaa gaactgggta ctcaacactg agcagatctg ttctttgagc taaaaacat
61 gtgctgtacc aagagtttgc tcctggctgc tttgatgtca gtgctgctac tccacctctg
121 cggcgaatca gaagcagcaa gcaactttga ctgctgtctt ggatacacag accgtattct
181 tcatcctaaa tttattgtgg gcttcacacg gcagctggcc aatgaaggct gtgacatcaa
241 tgctatcatc tttcacacaa agaaaaagtt gtcgtgtgct gcaaatccaa aacagacttg
301 ggtgaaatat attgtgcgtc tcctcagtaa aaaagtcaag aacatgtaaa aactgtggct
361 tttctggaat ggaattggac atagcccaag aacagaaaga accttgctgg ggttggaggt
421 ttcacttgca catcatggag ggtttagtgc ttatctaatt tgtgcctcac tggacttgct
481 caattaatga agttgattca tattgcatca tagtttgctt tgtttaagca tcacattaaa
541 gttaaactgt attttatggt atttatagct gtaggttttc tgtgtttage tatttaatac
601 taattttcca taagctatgt tggtttagtg caaagtataa aattatattt ggggggggaat
661 aagattatat ggactttctt gcaagcaaca agctattttt taaaaaaact atttaacatt
721 cttttgttta tattgttttg tctcctaaat tgttgtaatt gcattataaa ataagaaaaa
781 cattaataag acaaatatt

//
LOCUS          HUMAR          538 bp      mRNA          PRI          11-SEP-1996
DEFINITION     Human mRNA for chemokine, complete cds.
ACCESSION      D43767
NID            g1536878
KEYWORDS       chemokine, thymus and activation-regulated; chemokine.
SOURCE         Homo sapiens male peripheral blood cDNA to mRNA, clone:D3A.
ORGANISM       Homo sapiens
               Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
               Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
               Hominidae;
               Homo.
REFERENCE      1 (sites)
AUTHORS        Imai,T., Yoshida,T., Baba,M., Nishimura,M., Kakizaki,M. and
               Yoshie,O.
TITLE          Molecular cloning of a novel T cell-directed CC chemokine
expressed      in thymus by signal sequence trap using Epstein-Barr virus
vector
JOURNAL        J. Biol. Chem. 271 (35), 21514-21521 (1996)
MEDLINE        96355526
REFERENCE      2 (bases 1 to 538)
AUTHORS        Imai,T.
JOURNAL        Unpublished (1996)
REFERENCE      3 (bases 1 to 538)
AUTHORS        Imai,T.
TITLE          Direct Submission
JOURNAL        Submitted (07-DEC-1994) to the DDBJ/EMBL/GenBank databases.
Toshio
               Imai, Shionogi Institute for Medical Science; 2-5-1 Mishima,
               Settsu, Osaka 566, Japan (Tel:06-382-2612, Fax:06-382-2598)
FEATURES
  source       Location/Qualifiers
               1..538
               /organism="Homo sapiens"
               /db_xref="taxon:9606"
               /clone="D3A"
               /sex="male"
               /tissue_type="peripheral blood"
  CDS          53..337
               /note="thymus and activation regulated"
               /codon_start=1
               /product="chemokine"
               /db_xref="PID:d1008410"
               /db_xref="PID:g1536879"

/translation="MAPLKMLALVTLLLGASLQHIHAARGTNVGRECCLEYFKGA IPL
              RKLKTWYQTSSEDCSRDAIVFVTVQGRAICSDPNNKRVKNAVKYLSLERS"

```

```

BASE COUNT      118 a      168 c      149 g      103 t
ORIGIN
    1 cccctgagcag agggacctgc acacagagac tccctcctgg gctcctggca ccatggcccc
   61 actgaagatg ctggccctgg tcaccctcct cctggggggt tctctgcagc acatccacgc
  121 agctcgaggg accaatgttg gccgggagtg ctgcctggag tacttcaagg gagccattcc
  181 ccttagaaaag ctgaagacgt ggtaccagac atctgaggac tgctccaggg atgccatcgt
  241 ttttgtaact gtgcagggca gggccatctg ttcggacccc aacaacaaga gagtgaagaa
  301 tgcagttaaa tacctgcaa gccttgagag gtcttgaagc ctctcacc cagactcctg
  361 actgtctccc gggactacct gggacctcca ccgttggtgt tcaccgcccc caccctgagc
  421 gcctgggtcc aggggagggc ttccaggagc gaagaagagc cacagtgagg gagatcccat
  481 ccccttgtct gaactggagc catgggcaca aagggccagc attaaagtct ttatcctc

```

//

```

LOCUS      HUMEOTAXIN      807 bp      mRNA      PRI      25-SEP-1996
DEFINITION Human mRNA for eotaxin, complete cds.
ACCESSION  D49372
NID        gl552240
KEYWORDS   eotaxin; eosinophil-selective CC chemokine; chemoattractant.
SOURCE     Homo sapiens Small intestine, proximal cDNA to mRNA, clone:141.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE  1 (bases 1 to 807)
AUTHORS   Kitaura,M., Nakajima,T., Imai,T., Harada,S., Combadiere,C.,
            Tiffany,H.L., Murphy,P.M. and Yoshie,O.
TITLE     Molecular cloning of human eotaxin, an eosinophil-selective CC
            chemokine, and identification of a specific eosinophil eotaxin
            receptor, CC chemokine receptor 3
JOURNAL    J. Biol. Chem. 271 (13), 7725-7730 (1996)
MEDLINE    96205964
REFERENCE  2 (bases 1 to 807)
AUTHORS   Yoshie,O.
TITLE     Direct Submission
JOURNAL    Submitted (15-FEB-1995) to the DDBJ/EMBL/GenBank databases.
Osamu
            Yoshie, Shionogi Institute for Medical Science; 2-5-1 Mishima,
            Settsu, Osaka 566, Japan (E-mail:osamu.yoshie@shionogi.co.jp,
            Tel:06-382-2612, Fax:06-382-2598)
COMMENT    On Sep 20, 1996 this sequence version replaced gi:1313900.
FEATURES   Location/Qualifiers
            source          1..807
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /clone="141"
                        /tissue_type="Small intestine, proximal"
            CDS             99..392
                        /codon_start=1
                        /product="eotaxin"
                        /db_xref="PID:d1008966"
                        /db_xref="PID:gl552241"
            /translation="MKVSAALLWLLIIAAAFSPQGLAGPASVPTTCCFNLANRKIPLQ
            RLESYRRITSGKCPQKAVIFKTKLAKDICADPKKKWVQDSMKYLDQKSPTPKP"
            misc_signal      548..557
                        /note="mRNA destabilization signal"
            polyA_signal     775..780
            polyA_site       807

```

```

BASE COUNT      229 a      198 c      147 g      233 t
ORIGIN
    1 gcattttttc aagttttatg attttattaa cttgtggaac aaaaataaac cagaaaccac
   61 cactctctcac gccaaagctc acaccttcag cctccaacat gaaggtctcc gcagcacttc
  121 tgtggctgct gctcatagca gctgccttca gcccccaggg gctcgctggg ccagcttctg
  181 tcccaaccac ctgctgcttt aacctggcca ataggaagat accccttcag cgactagaga
  241 gctacaggag aatcaccagt ggcaaagtgc ccagaaagc tgtgatcttc aagaccaaac
  301 tggccaagga tatctgtgcc gacccaaga agaagtgggt gcaggattcc atgaagtatc
  361 tggaccaaaa atctccaact ccaaagccat aaataatcac catttttgaa accaaaccag
  421 agcctgagtg ttgcctaatt tgttttcct tcttacaatg cattctgagg taacctcatt
  481 atcagtccaa agggcatggg ttttattata tatatatata tttttttttt aaaaaaaaaa
  541 gtattgcatt taatttattg aggccttaaa acttatcttc catgaatatic agttattttt

```

```

601 aaactgtaaa gctttgtgca gattctttac cccctgggag cccaattcg atccccgtc
661 acgtgtgggc aatgttcccc ctctcctctc ttcctccctg gaatcttgta aaggctctgg
721 caaagatgat cagtatgaaa atgtcattgt tcttgtgaac ccaaagtgtg actcattaaa
781 tggaagtaaa tggtgtttta ggaatac

//
LOCUS       HSCCCHHEM      232 bp      RNA              PRI      10-SEP-1996
DEFINITION   H.sapiens mRNA for CC-chemokine.
ACCESSION    Z69291
NID          g1181148
KEYWORDS     CC-chemokine.
SOURCE       human.
  ORGANISM   Homo sapiens
              Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
              Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE    1 (bases 1 to 232)
  AUTHORS    Bartels,J.H., Schlueter,C., Richter,E., Christophers,E. and
              Schroeder,J.M.
  TITLE      Cloning of a novel human chemokine homologous to human monocyte
              chemoattractant proteins and rodent eotaxins
  JOURNAL     Unpublished
REFERENCE    2 (bases 1 to 232)
  AUTHORS    Bartels,J.H.
  TITLE      Direct Submission
  JOURNAL     Submitted (01-FEB-1996) Bartels J. H.,
              Christian-Albrechts-Universitaet          zu          Kiel,
Dermatology/Hautklinik,
Mol.Biol.Lab.609, Schittenhelmstr. 7, Kiel, Schleswig-Holstein,
Germany, D-24105
REFERENCE    3 (bases 1 to 232)
  AUTHORS    Bartels,J., Schluter,C., Richter,E., Noso,N., Kulke,R.,
              Christophers,E. and Schroder,J.M.
  TITLE      Human dermal fibroblasts express eotaxin: molecular cloning,
mRNA
              expression, and identification of eotaxin sequence variants
  JOURNAL     Biochem. Biophys. Res. Commun. 225 (3), 1045-1051 (1996)
  MEDLINE     96374440
FEATURES
  source      Location/Qualifiers
              1..232
              /organism="Homo sapiens"
              /db_xref="taxon:9606"
              /clone="clones 4(9512),
              14(9512),15(9512),10(9601),11(9601)"
              /tissue_type="foreskin"
              /cell_type="fibroblast"
              /sex="Male"
  mRNA        <1..>232
              /citation=[1]
              /product="CC-chemokine"
  sig_peptide 56..109
              /citation=[1]
  CDS         56..>232
              /function="putative chemoattractant protein"
              /note="sequence homology to human MCP-1, MCP-2 and
MCP-3
              and to rodent eotaxins"
              /citation=[1]
              /codon_start=1
              /product="CC-chemokine, preprotein"
              /db_xref="PID:e221070"
              /db_xref="PID:g1181149"
              /db_xref="SWISS-PROT:P50877"

/translation="MKVSAALLWLLIAAAFSPQGLAGPASVPTTCCFNLANRKIPLQ
              RLESYRRITSGKCPQ"
  mat_peptide 110..>232
              /citation=[1]
              /function="putative chemoattractant protein"
              /product="CC-chemokine"
BASE COUNT   55 a      82 c      50 g      42 t      3 others
ORIGIN
      1 accaaaccag aaaccwccam ytctcagcc aaagtcaca cttcagcct ccaacatgaa

```

```

        61 ggtctccgca gcgcttctgt ggctgtgtct catagcggct gccttcagcc cccaggggct
       121 cgctggggcca gcttctgtcc caaccacctg ctgctttaac ctggccaata ggaagatacc
       181 ccttcagcga ctagagagct acaggagaat caccagtggc aaatgtcccc ag
//
LOCUS      HSHCC1GEN      4037 bp      DNA      PRI      01-OCT-1995
DEFINITION H.sapiens gene for chemokine HCC-1.
ACCESSION  Z49269
NID        g1004266
KEYWORDS   chemokine.
SOURCE      human.
ORGANISM    Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 4037)
AUTHORS    Pardigol,A., Maegert,H.J., Cieslak,A., Hill,O., Schulz-
Knappe,P.
TITLE       and Forssmann,W.G.
            Nucleotide Sequence of the Gene for the Human Chemokine HCC-1
JOURNAL     Unpublished
REFERENCE  2 (bases 1 to 4037)
AUTHORS    Pardigol,A.
TITLE       Direct Submission
JOURNAL     Submitted (18-MAY-1995) Andreas Pardigol, Molecular Biology,
Lower
            Saxony Institute for Peptide Research, Feodor-Lynen-Strasse 31,
            Hannover, Lower Saxon, 30625, Germany
FEATURES    Location/Qualifiers
            source          1..4037
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /clone="ph3b7"
                        /dev_stage="adult"
                        /tissue_type="placenta"
                        /clone_lib="lambda FIX II, Cat.Nr. 946203, Stratagene"
                        /sex="male"
            TATA_signal     727..733
                        /note="putative, determined by consensus rules."
            5'UTR           764..833
                        /note="first base determined by means of consensus
rules"
            exon            764..912
                        /note="first base determined by means of consensus
rules;
                        base 780 is the first base of cDNA (Z49270)"
            CDS              join(834..912,3021..3135,3585..3672)
                        /codon_start=1
                        /product="chemokine HCC-1"
                        /db_xref="PID:g1004267"
            /translation="MKISVAaipffllitlalgktesssrgpyhpsccfitytthyki
            PRQRIMDYyETNSQCSKPGIVFITKRGHSVCTNPSPDKWVQDIYKDMKEN"
            intron           913..3020
                        /number=1
            exon            3021..3135
                        /number=2
            intron           3136..3584
                        /number=2
            exon            3585..3817
                        /number=3
            3'UTR           3673..3817
BASE COUNT  1023 a   1048 c   1004 g   962 t
ORIGIN
        1 gagctccggtt gggagtcacca tgtttcttta tggcataatg ggtgagaaca cagacttgga
       61 agccaaacca cctgaatttg aacccagtt ccatttacca actgtcaaaa gcttaggctt
      121 tgattctaag cctgttctct caactgctgt tctaaagatt aaataggcta atattcataa
      181 ggcaactggg acagtggcct gtgtgtatag caaccattat ataagtgaat tatctactga
      241 gcaccacagc acttcttcac tccatggtgt ggtgaccaga atggagatga gacagagaac
      301 tgcaggttct gcttcgagtt taagtttaga ttcccttga ccaatgagac ctgacttgga
      361 ggagtctctg cctcattcca ttaccccaaa caccctctag tctctagatg aacagatcct
      421 gaatgtccag gccccacgtg gcctgttcta aggcctgaga tggaattgga tacaggacac

```

```

481 atccagcctt gagatctttt gctaagtgtg acacagtgcc cccagccctg tgctcatgtt
541 catgcctagg gaaaggcttc tatcaaaaga gttgaacttc tccccactgg ggaagggaaga
601 ccatctcttc ccttaaaccct tggctctccc tgcttctctc aggccaccaa caacacatgt
661 gcaggatatg aaattgctga ggcactactg ctttctactt tcccttccaa gtctcagctc
721 ccttattttta aaaaatattt ggcctcaatg atcatctctc aacaattcct caccgcagga
781 gcctctgaag ctcccaccag gccagctctc ctcccacaac agcttcccac agcatgaaga
841 tctccgtggc tgccattccc ttcttctccc cttccttcca acctagagcc tgcagggaaa
901 aatcctcctc acgtgagtgc aatgccttgt cttccttcca acctagagcc gaaaggaggg
961 taagcaggag tgagggtggg gctcagggga agaccaggag cagggactca tctactgagc
1021 ctgggtatctt cttgaaattg tgtgtatagc aacattatat aaatgaatta tctactgagc
1081 accacagcac ttcaccccat ggtgtggtga gcaggatgga gatgagactt aggactgtag
1141 gttctgctta agagtttaag ttgggactct ccagccttga ccaatgagac ttgacttggg
1201 agaetccagg cttcattcca ctaccccaaa tgccctctag tctccaaata aacagatcct
1261 gaatctccag gcctcacatg gccttgatct cttatcattg cccccagga ccagtcctcc
1321 cttgcccctc aggacatgga gtgagaccag cctgcctctc tactccctca atttctctct
1381 ctttgcgcgt aagcaaaaga gtggcccacc ccatttgggg tatatttctt cagggagatt
1441 agggagcagt tcttgagccc ctcaagggca ttttcttatt ggccctctga ggtttgggct
1501 cagcctgctt ccagcgtcac ctgtgcccag aggtgagcgc attgcttggg atttcttaga
1561 ggggaaacac gacagtgtgg ggtccatcct agggccctct ttctcagctg atttcttaga
1621 ataagctgcc ttttagagata accaaaacta tttatcactc ttcctattta cctactctcc
1681 ttttcagaaa ctggggggaa accgaagggt gttaaaatac agctaaagt tgggtggtag
1741 tgcacagtgt gacttgcctt ctccgatgct atttgtcagc tcagaggaaac aagggtggag
1801 agtataggag ctctgactgg gtctcaggaa acagggggcc cttatgccgt tctttggatc
1861 gtgaggatgc tgccctggaat ggagctggaa aacaggatga gacccttcca cccagacatc
1921 tggccaccct cagtgccttc tgaggccatt gtgatgcaca tccatgattc tatgaagcag
1981 ggtcacataa catgcacaca cctgatttct ccactccata accacaacat gtgcctgttt
2041 gtacagggct cttggcctac aatgtccttc ctgtacctc tataattcaa gcttgggggtg
2101 gctgctgtca ccttgcctct cctataaaag ccatgaaact tctcaatcag aaaatagatg
2161 aaaaaatcac ccaatccagt gattttttaa acttttttaga ccacaaaacc tttcttcaa
2221 gcaatatctt ccacagaggc ccaatatgta aaacagaaaa aatgggttga gtagggtaca
2281 agacaccact ctcaaatgca gcaaggcctc cacaatagtc cctgaggccc ccagagctca
2341 gtgtaaaaac cactgatgca gtccaagggc ctcatctaca gaggaggaa cagggggaaa
2401 gtaaaatggc cacagtacac aggaagcaca ggcaagggtta ggttaggatt tgggtgccct
2461 gactctgtgg cctttgtcct tggggcttgc tgtgggcatc ctgctctctc tgcaggttgt
2521 cggttcaatg gggacatggg caggggtggg cactaggagg ggctgggttt gcattccaa
2581 atggcatgtc tccaaatccc tattgggatt tcttccaaat attcctccta tttggagcac
2641 ctttcccga taaggcatga aggctgcatt atattggcca agtccctagc cttctctgcc
2701 agtcggcccc cagagatggt gtaagaagat ctgagtgtgc tgctcttcaa tcctggagtt
2761 gaaagtcac caccagtctt tccaagaggg gttgaagaaa agggaggagg gtgattgatg
2821 atagggagg agaaaaagaa gagcccagga gtaccatgga gaaggagaag agaagatgag
2881 gaaagcctac tctcccctcc aagttctgag gggctgtctc ctcttctctt cctcctcca
2941 tgccctcagc ttgcaggagc agccaatggt atggccttta acaagggggc cctcctcagc
3001 atctgatgct ctctcctcag ggggacctta ccacctccta gagtgtctgt tcacctacac
3061 tacttacaag atcccgcgtc agcggattat ggattactat gagaccaaca gccagtgtct
3121 caagcccga attgtgtagg ttgtacacac acatcacact ggggggagag ggagccagca
3181 gggcctcctg gaggggaagca gggagtgtgt gtggaatggg gaccccccagc gtacctcca
3241 ggtgtgacta catggggaga ggcagctgag gggcaatctg agcgtcttct ggctggagcc
3301 tgcaggagcc atggggaaac tgaccccatg gatggggaga tgacagagaa ggggagaagaa
3361 ggcaagaggg cacttctcca gggggacaca gagactagat ggtcttaggg gtccctagaa
3421 ccgaagagta tgtctcagag aggaagactg cttctgtgga cctctgtgga agaaaggaaa
3481 agcagtata gtcaggtggg gaatttagga gggagggaag atgggctgtc tcttccggcc
3541 actgggcccc tcggtttgtg atccttctcc ctcttgctcc acagcttcat caccaaaagg
3601 ggccattccg tctgtaccaa cccagtgac aagtgggtcc aggaactatg caaggacatg
3661 aaggagaact gagtgaccca gaagggtgg cgaaggcaca gctcagagac ataaagagaa
3721 gatgccaagg cccctctctc caccaccgc taactctcag cccagtcac cctcttggag
3781 cttccctgct ttgaattaaa gaccactcat gctcttccct ggctcattc ctttctacgg
3841 gatctactca ttggccatgc actgaggaca ccagggtgtg gcaccctcg catcaagct
3901 cgctctcag aagttttggt ggagcctggt acaaaaaata ggtcaggcct gcaatgcagg
3961 tagtgagaag cagaaagtga gaaagaaaag cagtgtaaag accgtctcct cctcagcagc
4021 aacagtagca gaccccg

```

```

//
LOCUS       HSCC21             925 bp      mRNA             PRI             30-JUN-1998
DEFINITION  H.sapiens mRNA for chemokine CC-2 and CC-1.
ACCESSION   Z70292
NID         g1296608
KEYWORDS    chemokine CC-1; chemokine CC-2.
SOURCE      human.
  ORGANISM  Homo sapiens
            Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1 (bases 1 to 925)

```

AUTHORS Pardigol, A., Forssmann, U., Zucht, H.D., Loetscher, P.,
Schulz-Knappe, P., Baggiolini, M., Forssmann, W.G. and Magert, H.J.

TITLE HCC-2, a human chemokine: gene structure, expression pattern,
and biological activity

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 95 (11), 6308-6313 (1998)

MEDLINE 98263352

REFERENCE 2 (bases 1 to 925)

AUTHORS Pardigol, A.

TITLE Direct Submission

JOURNAL Submitted (25-MAR-1996) Andreas Pardigol, IV - Molecular
Biology, Lower Saxony Institute for Peptide Research, Feodor-Lynen-
Strasse 31, Hannover, Lower Saxony, 30625, Germany

FEATURES Location/Qualifiers

source 1..925
/organism="Homo sapiens"
/db_xref="taxon:9606"
/dev_stage="adult"
/tissue_type="liver"
/clone_lib="PCR fragments"

5'UTR 1..55

CDS 56..397
/note="putative; first coding region of a bicistronic
mRNA"
/codon_start=1
/product="chemokine CC-2"
/db_xref="PID:e233855"
/db_xref="PID:g1296609"
/db_xref="SWISS-PROT:Q16663"

/translation="MKVSVAAALSCMLVAVLGSQAQFTNDAETELMMSKLPLENPVVL

NSFHFAADCCTSYISQSIPLMKSYFETSSECSKPGVIFLTKKGRQVCAKPSGPGVQ

misc_feature DCMKKLPYSI"
398..498
/note="spacing region between two coding regions of
the bicistronic mRNA"

CDS 499..780
/codon_start=1
/evidence=experimental
/product="chemokine CC-1"
/db_xref="PID:e233856"
/db_xref="PID:g1296610"
/db_xref="SWISS-PROT:Q16627"

/translation="MKISVAAIPFFLLITIALGKTESSSRGPHYHPSECCFTYTTYKI

3'UTR PRQRIMDYETNSQCSKPGIVFITKRGHVCTNP SDKWVQDYIKDMKEN"
781..925

polyA_signal 902..908

BASE COUNT 240 a 296 c 199 g 190 t

ORIGIN

1 ccaggaagca gtagagccag gaggcctcgg ccagccctgc ctgcccacca ggaggatgaa
61 ggtctccgtg gctgcctctt cctgcctcat gcttgttget gtccttggat cccaggccca
121 gttcacaat gatgcagaga cagagttaat gatgtcaaag cttccactgg aaaaatccagt
181 agttctgaac agctttcact ttgctgtgta ctgctgcacc tcctacatct cacaagcat
241 cccgtgttca ctcatgaaaa gttattttga aacgagcagc gagggtctca agccaggtgt
301 catattcctc accaagaagg ggcggcaagt ctgtgcaaaa cccagtgggtc cgggagttca
361 ggattgcatg aaaaagctga agccctactc aatataataa taaagagaca aaagaggcca
421 gccaccacc tccaacacct cctgagcctc tgaagctccc accaggccag ctctcctccc
481 acaacagctt cccacagcat gaagatctcc gtggctgcca ttcccttctt cctcctcatc
541 accatcgccc tagggaccac gactgaatcc tcctcacggg gacettacca cccctcagag
601 tgctgcttca cctacactac ctacaagatc ccgcgtcagc ggattatgga ttactatgag
661 accaacagcc agtgctccaa gcccggaatt gtcttcatca ccaaaagggg ccattccgtc
721 tgtaccaacc ccagtgacaa gtgggtccag gactatatca aggacatgaa ggagaactga
781 gtgaccaga aggggtggcg aaggcacagc tcagagacat aaagagaaga tgccaaggcc
841 ccctcctcca cccaccgcta actctcagcc ccagtcaccc tcttgagact tccctgcttt
901 gaattaaaga ccactcatgc tcttct

//

LOCUS HSCC23 973 bp RNA PRI 03-MAY-1996
 DEFINITION H.sapiens mRNA for chemokine CC-2 and CC-3.
 ACCESSION Z70293
 NID g1296611
 KEYWORDS Human chemokine CC-2; Human chemokine CC-3.
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 973)
 AUTHORS Pardigol, A., Maegert, H.J., Zucht, HD., Forssmann, W.G. and
 Schulz-Knappe, P.
 TITLE Transcription of a Human Tandem Gene results in a Mature
 Bicistronic mRNA encoding two Novel CC-Chemokines
 JOURNAL Unpublished
 REFERENCE 2 (bases 1 to 973)
 AUTHORS Pardigol, A.
 TITLE Direct Submission
 JOURNAL Submitted (25-MAR-1996) Andreas Pardigol, IV - Molecular
 Biology, Lower Saxony Institute for Peptide Research, Feodor-Lynen-
 Strasse 31, Hannover, Lower Saxony, 30625, Germany
 FEATURES
 source Location/Qualifiers
 1..973
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /dev_stage="adult"
 /tissue_type="liver"
 /clone_lib="PCR fragments"
 5'UTR 1..55
 CDS 56..397
 /note="putative; first coding region of a bicistronic
 mRNA"
 /codon_start=1
 /product="chemokine CC-2"
 /db_xref="PID:e233857"
 /db_xref="PID:g1296612"
 /translation="MKVSVAAALSCMLVAVLGSQAQFTNDAETELMMSKLPLENPVVL
 NSFHFAADCCTSYISQSI PCSLMKSYFETSSECSKPGVIFLTKKGRQVCAKPSGPGVQ
 DCMKKLKPYSI"
 misc_feature 398..498
 /note="spacing region between two coding regions of
 the
 bicistronic mRNA"
 CDS 499..828
 /note="putative"
 /codon_start=1
 /product="chemokine CC-3"
 /db_xref="PID:e233858"
 /db_xref="PID:g1296613"
 /translation="MKISVAAIPFFLLITIALGKTTESSQTGGKPKVVKIQLKLVGG
 PYHPSECCFTYTTYKIPRQIRIMDYETNSQCSKPGIVFITKRGHVSVCTNP SDKWVQDY
 IKDMKEN"
 3'UTR 829..973
 polyA_signal 950..956
 BASE COUNT 257 a 301 c 215 g 200 t
 ORIGIN
 1 ccaggaagca gtgagcccg gagtcctcgg ccagccctgc ctgcccacca ggaggatgaa
 61 ggtctccgtg gctgcccctc cctgcctcat gcttggtgct gtccttggat cccaggccca
 121 gttcacaaat gatgcagaga cagagttaat gatgtcaaag cttccactgg aaaatccagt
 181 agttctgaac agctttcact ttgctgctga ctgctgcacc tcctacatct cacaaagcat
 241 cccgtgttca ctcataaaaa gttattttga aacgagcagc gagtgctcca agccagggtg
 301 catattcctc accaagaagg ggcggcaagt ctgtgcaaaa cccagtgggc cgaggattca
 361 ggattgcata aaaaagctga agccctactc aatataataa taaagagaca aaaggagcca
 421 gccacccacc tccaacacct cctgagcctc tgaagctccc accaggccag ctctcctccc

```

481 acaacagctt cccacagcat gaagatctcc gtggtgcca ttcccttctt cctcctcatc
541 accatcgccc tagggaccaa gactgaatcc tctcacaaa ctggggggaa accgaagggtt
601 gttaaaatag agctaaagt ggtgggggga ccttaccacc cctcagagtg ctgcttcacc
661 tacactacct acaagatccc gcgtcagcgg attatggatt actatgagac caacagccag
721 tgctccaagc ccggaattgt cttcatcacc aaaaggggccc attccgtctg tacciaacccc
781 agtgacaagt ggggccagga ctatatcaag gacatgaagg agaactgagt gaccagaag
841 ggggtggcgaa ggcacagctc agagacataa agagaagatg ccaaggcccc ctcctccacc
901 caccgctaac tctcagcccc agtcaccctc ttggagcttc cctgctttga attaaagacc
961 actcatgctc ttc

```

//

```

LOCUS      HSU91746      1430 bp      mRNA      PRI      12-MAR-1998
DEFINITION Homo sapiens IL-10-inducible chemokine (HCC-4) mRNA, complete
            cds.
ACCESSION  U91746
NID        g2581780
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1430)
AUTHORS    Hedrick,J.A., Helms,A., Gorman,D. and Zlotnik,A.
TITLE      Identification of a novel human CC chemokine upregulated by IL-
10
JOURNAL    Blood (1998) In press
REFERENCE  2 (bases 1 to 1430)
AUTHORS    Hedrick,J.A., Helms,A., Gorman,D. and Zlotnik,A.
TITLE      Direct Submission
JOURNAL    Submitted (02-MAR-1997) Immunology, DNAX Research Institute,
901
            California Ave, Palo Alto, CA 94304, USA
FEATURES   Location/Qualifiers
            source          1..1430
                               /organism="Homo sapiens"
                               /db_xref="taxon:9606"
                               /chromosome="17"
            gene           1..1430
                               /gene="HCC-4"
            CDS             1..363
                               /gene="HCC-4"
                               /note="CC or beta chemokine family member"
                               /codon_start=1
                               /product="IL-10-inducible chemokine"
                               /db_xref="PID:g2581781"

```

/translation="MKVSEAALSLVLILIITSASRSQPKVPEWVNTPTCCLKYEK

VLPRRLVVG YRKALNCHLPAIIFVTKRNREVCTNPNDWVQEIYKDPNLP LLLPTRNLS
TVKIITAKNGQPQLLSQ"

BASE COUNT 401 a 351 c 293 g 385 t
ORIGIN

```

1 atgaaggctt ccgaggctgc cctgtctctc cttgtcctca tccttatcat tacttcggct
61 tctcgagccc agccaaaagt tcttgagtgg gtgaacaccc catccacctg ctgcctgaag
121 tattatgaga aagtgttgcc aaggagacta gtgggtggat acagaaaggc cctcaactgt
181 cacctgccag caatcatctt cgtcaccaag aggaaccgag aagtctgcac caacccaat
241 gacgactggg tccaagagta catcaaggat cccaacctac ctttgtgccc taccaggaac
301 ttgtccacgg ttaaaattat tacagcaaag aatggtcaac cccagctcct caactcccag
361 tgatgaccag gcttttagtg aagcccttgt ttacagaaga gaggggtaaa cctatgaaaa
421 caggggaagc cttattagtc tgaaactagc cagtcacatt gagagaagca gaacaatgat
481 caaaataaag gagaagtatt tcgaatatct tctcaatctt aggaggaaat accaaagtta
541 agggacgtgg gcagaggtac gctcttttat ttttatattt atatttttat ttttttgaga
601 taggtcttac tctgtcacc caggctggagt gcagtgggtg gatcttggct cacttgatct
661 tggctcactg taacctccac ctcccaggct caagtgatcc tcccaccca gctcccag
721 tagctgggac tacaggcttg cgccaccaca cctggctaatt ttttgattt ttggtagaga
781 cgggattcta ccatgttgcc caggctgggt tcaaactcgt gtgcccgaag aatccacctg
841 cctcagcctt ccaaaagtgc tgggattaca ggctgagcc accacatccg gccagtgcac
901 tcttaataca cagaaaaata tatttcacat ccttctctg ctctctttca attcctcact
961 tcacaccagt acacaagcca ttctaaatac ttagccagtt tccagccttc cagatgatct
1021 ttgccctctg ggtcttgacc cattaagagc cccatagaac tcttgatttt tctgttccat
1081 ctttatggat ttttctggat ctatatcttc ttcaattatt ctttcatttt ataagtcaac

```



```

1141 tttttcatag gaagtcgga tgggaatatt cacattaatc atttttgag agactttgct
1201 agatcctctc atattttgtc ttctcaggg tggcaggggt acagagagtg cctgattgga
1261 aaaaaaaaaa aaagagagag agagagaaga agaagaagaa gagacacaaa tctctacctc
1321 ccatgttaag ctttgcagga cagggaaaga aagggtatga gacacggcta ggggtaaact
1381 cttagtccaa aaccaagca tgcaataaat aaaactcctt tatttgacaa

```

//

LOCUS AB007454 1503 bp mRNA PRI 09-APR-1998
 DEFINITION Homo sapiens mRNA for chemokine LEC precursor, complete cds.
 ACCESSION AB007454
 NID g2723285
 KEYWORDS chemokine LEC precursor.
 SOURCE Homo sapiens liver cDNA to mRNA.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
 Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (sites)
 AUTHORS Shoudai, K., Hieshima, K., Fukuda, S., Iio, M., Miura, R., Imai, T.,
 Yoshie, O. and Nomiyama, H.
 TITLE Isolation of cDNA encoding a novel human CC chemokine NCC-4/LEC
 JOURNAL Biochim. Biophys. Acta 1396 (3), 273-277 (1998)
 MEDLINE 98207719
 REFERENCE 2 (bases 1 to 1503)
 AUTHORS Nomiyama, H.
 TITLE Direct Submission
 JOURNAL Submitted (19-SEP-1997) to the DDBJ/EMBL/GenBank databases.
 Hisayuki Nomiyama, Kumamoto University Medical School,
 Department of Biochemistry; Honjo 2-2-1, Kumamoto, Kumamoto 860-0811,
 Japan
 (E-mail: nomiyama@gpo.kumamoto-u.ac.jp, Tel: 81-96-373-5063,
 Fax: 81-96-372-6140)

FEATURES
 source Location/Qualifiers
 1..1503
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /tissue_type="liver"
 sig_peptide 77..145
 CDS 77..439
 /codon_start=1
 /product="chemokine LEC precursor"
 /db_xref="PID:d1024963"
 /db_xref="PID:g2723286"

/translation="MKVSEALSLVLILITSAASRQPKVPEWNTPTCCLKYIEK

VLPRRLVVGGRKALNCHLPAIIFVTKRNREVCTNPNDWVQYIKDPNLPLLPTRNLS
 TVKIIITAKNGQPQLLSQ"

mat_peptide 146..436
 polyA_signal 560..565
 polyA_signal 1485..1490

BASE COUNT 417 a 374 c 312 g 400 t

ORIGIN

```

1 gttggcaagc ggaccaccag caacagacaa catcttcatt cggctctccc tgaagctgta
61 ctgcctcgct gagaggatga aggtctccga ggctgccttg tctctccttg tctcctcct
121 tatcattact tcggtctctc gcagccagcc aaaagttcct gagggtgga acaccccatc
181 cacctgctgc ctgaagtatt atgagaaagt gttgccaagg agactagtgg tgggatacag
241 aaaggccctc aactgtcacc tgccagcaat catcttcgtc accaagagga accgagaagt
301 ctgcaccaac cccaatgacg actgggtcca agagtacatc aaggatccca acctaccttt
361 gctgcctacc aggaacttgt ccacgggtta aattattaca gcaaagaatg gtcaacccca
421 gctcctcaac tccagtgat gaccaggctt tagtggaagc cctgtttac agaagagagg
481 ggtaaaccta tgaaaacagg ggaagcctta ttaggctgaa actagccagt cacattgaga
541 gaagcagaac aatgatcaaa ataaaggaga agtatttcga atattttctc aatcttagga
601 ggaataacca aagttaaggg acgtgggcag aggtacgctc ttttattttt atatttata
661 ttttattttt ttgagatagg gtcttactct gtcacccagg ctggagtga gtggtgtgat
721 cttggctcac ttgatcttgg ctactgtaa cctccacctc ccaggctcaa gtgactctcc
781 caccacagcc tcccgagtag ctgggactac aggtctgctc caccacacct ggctaatttt
841 tgtatttttg gtagagacgg gattctacca tgttgcccag gctggtctca aactcgtgtg
901 cccaagcaat ccacctgctc cagccttcca aaagtgtctg gattacagge gtgagccacc
961 acatccggcc agtgcactct taatacacag aaaaaatata ttcacatcct tctcctgctc
1021 tctttcaatt cctcacttca caccagtaca caagccattc taaatactta gccagtttcc

```

```

1081 agccttccag atgatctttg ccctctgggt cttgacccat taagagcccc atagaactct
1141 tgatttttcc tgtccatctt tatggatttt tctggatcta tttttcttcc aattattctt
1201 tcattttata atgcaacttt ttcataaggaa gtccggatgg gaattattcac attaatcatt
1261 tttgcagaga ctttgctaga tcctctcata tttgtcttc ctcaggggtg caggggtaca
1321 gagatgtcct gattggaaaa aaaaaaaaaa gagagagaga gagaagaaga agaagaagag
1381 acacaaatct ctacctcca tgttaagctt tgcaggacag ggaaagaaag ggtatgagac
1441 acggctaggg gtaaactctt agtccaaaac ccaagcatgc aataataaaa actcccttat
1501 ttg

```

//

```

LOCUS      AF001979      800 bp      mRNA      PRI      20-NOV-1997
DEFINITION Homo sapiens beta chemokine mRNA, complete cds.
ACCESSION  AF001979
NID        g2624924
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 800)
AUTHORS    Hedrick, J.A. and Zlotnik, A.
TITLE      Identification and characterization of a novel beta chemokine
            containing six conserved cysteines
JOURNAL    J. Immunol. 159 (4), 1589-1593 (1997)
MEDLINE    97400322
REFERENCE  2 (bases 1 to 800)
AUTHORS    Hedrick, J.A. and Zlotnik, A.
TITLE      Direct Submission
JOURNAL    Submitted (01-MAY-1997) Immunobiology, DNAX Research Institute,
901        California Ave, Palo Alto, CA 94304, USA
FEATURES   Location/Qualifiers
            source          1..800
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
            CDS             1..405
                        /note="6Ckine; CC chemokine"
                        /codon_start=1
                        /product="beta chemokine"
                        /db_xref="PID:g2624925"

```

/translation="MAQSLALSLLILVLAFGIPRTQGS DGAQDCCLKYSQRKIPAKV

```

VRSYRKQEPSLGC SIPAILFLPKRSQAE LCADPKELWVQQLMQHLDKTPSPQKPAQG
CRKDRGASKTGKKGKSGKCKRTERSQTPKGP"

```

```

BASE COUNT      203 a      248 c      210 g      139 t
ORIGIN

```

```

1 atggctcagt cactggctct gaggctcctt atcctgggtc tggcctttgg aatccccagg
61 acccaaggca gtgatggagg ggctcaggac tgttgccctca agtacagcca aaggaagatt
121 cccgccaaagg ttgtccgcag ctaccggaag caggaaccaa gcttaggctg ctccatccca
181 gctatcctgt tcttgcccg caagcgctct caggcagagc tatgtgcaga cccaaaggag
241 ctctgggtgc agcagctgat gcagcatctg gacaagacac catccccaca gaaaccagcc
301 cagggctgca ggaaggacag gggggcctcc aagactggca agaaaggaaa gggctccaaa
361 ggctgcaaga ggactgagcg gtcacagacc cctaaagggc catagcccag tgagcagcct
421 ggagccctgg agacccacc agcttcacca gcgcttgaag cctgaaccca agatgcaaga
481 aggaggctat gctcaggggc cctggagcag ccaccccatg ctggccttgc cacactcttt
541 ctctgctttt aaccacccca tctgcattcc cagctctacc ctgcatggct gagctgcccc
601 cagcaggcca ggtccagaga gaccgaggag ggagagtctc ccagggagca tgagaggagg
661 cagcaggact gtccccttga aggagaatca tcaggaccct ggacctgata cggctcccca
721 gtacacccca cctcttcctt gtaaatatga tttataccta actgaataaa aagctgttct
781 gtcttccac ccaaaaaaaa

```

//

```

LOCUS      HSU64197      821 bp      mRNA      PRI      25-JUN-1997
DEFINITION Homo sapiens chemokine exodus-1 mRNA, complete cds.
ACCESSION  U64197
NID        g1778716
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;

```

Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae;
Homo.
REFERENCE 1 (bases 1 to 821)
AUTHORS Hromas, R., Gray, P.W., Chantry, D., Godiska, R., Krathwohl, M.,
Fife, K., Bell, G.I., Takeda, J., Aronica, S., Gordon, M.,
Cooper, S.,
Broxmeyer, H.E. and Klemsz, M.J.
TITLE Cloning and characterization of exodus, a novel beta-chemokine
JOURNAL Blood 89 (9), 3315-3322 (1997)
MEDLINE 97275143
REFERENCE 2 (bases 1 to 821)
AUTHORS Hromas, R.A.
TITLE Direct Submission
JOURNAL Submitted (17-JUL-1996) Indiana University Medical Center,
Medicine, 975 W. Walnut St., Indianapolis, IN 46202, USA
FEATURES Location/Qualifiers
source 1..821
/organism="Homo sapiens"
/db_xref="taxon:9606"
/clone="Exodus-1"
/cell_type="islet"
/tissue_type="pancreas"
/dev_stage="adult"
CDS 43..330
/function="inhibits proliferation of hematopoietic
progenitors and HIV"
/codon_start=1
/product="chemokine exodus-1"
/db_xref="PID:g1778717"
/translation="MCCTKSLLLAALMSVLLHLHLCGESEASNFDCCLGYTDRILHPKF
IVGFTRQLANEGCDINAIIFHTKKKLSVCANPKQTVWKYIVRLLSKKVKNM"
variation 121^122
/insertion="insertion of an extra codon GCA at nt 121,
encoding for an alanine after the alanine at amino acid
position 26, represents the allelic difference of the
transcript isolated from macrophages"
BASE COUNT 258 a 134 c 156 g 273 t
ORIGIN
1 ggtactcaac actgagcaga tctgttcttt gagctaaaaa ccatgtgctg taccaagagt
61 ttgctcctgg ctgctttgat gtcagtgtcg ctactccacc tctgcggcga atcagaagca
121 agcaactttg actgctgtct tggatacaca gaccgtattc ttcactctaa atttattgtg
181 ggcttcacac ggcagctggc caatgaaggc tgtgacatca atgctatcat ctttcacaca
241 aagaaaaagt tgtctgtgtg cgcaaatcca aaacagactt gggtgaaata tattgtgcgt
301 ctccctcagta aaaaagtcaa gaacatgtaa aaactgtggc ttttctggaa tgggaattgga
361 catagcccaa gaacagaaaag aaccttgcgt ggggtggagg tttcacttgc acatcatgga
421 ggggtttagt cttatctaat ttgtgcctca cctggacttg tccaattaat gaagtgtgatt
481 catattgcat catagtttgc tttgtttaag catcacatta aagtgaact gtattttatg
541 ttatttatag ctgtaggttt tctgtgttta gctatttaac actaattttc cataagctat
601 tttgggttag tgcaaagtat aaaattatat ttggggggga ataagattat atggactttc
661 ttgcaagcaa caagctattt tttaaaaaaa actatttaac attcttttgt ttatattgtt
721 ttgtctccta aattgttgta atgtcattat aaaaataagaa aaatattaat aagacaaata
781 ttgaaaataa agaaacaaaa agtgcttctg ttaaaaaaaaa a
//
LOCUS HSU88320 828 bp mRNA PRI 18-DEC-1997
DEFINITION Human beta chemokine Exodus-2 mRNA, complete cds.
ACCESSION U88320
NID g2196919
KEYWORDS
SOURCE human.
ORGANISM Homo sapiens
Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (bases 1 to 828)
AUTHORS Hromas, R., Kim, C.H., Klemsz, M., Krathwohl, M., Fife, K.,
Cooper, S.,

Schnizlein-Bick, C. and Broxmeyer, H.E.
 TITLE Isolation and characterization of Exodus-2, a novel C-C
 chemokine
 JOURNAL with a unique 37-amino acid carboxyl-terminal extension
 MEDLINE J. Immunol. 159 (6), 2554-2558 (1997)
 REFERENCE 97444139
 2 (bases 1 to 828)
 AUTHORS Hromas, R.A.
 TITLE Direct Submission
 JOURNAL Submitted (04-FEB-1997) Medicine, Indiana University Medical
 Center, 975 West Walnut, Indianapolis, IN 46202, USA
 FEATURES Location/Qualifiers
 source 1..828
 /organism="Homo sapiens"
 /note="PCR amplified from activated THP-1 cells"
 /db_xref="taxon:9606"
 /clone_lib="Soares human placenta cDNA"
 /cell_line="THP-1"
 /cell_type="monoblast"
 CDS 15..419
 /codon_start=1
 /product="beta chemokine Exodus-2"
 /db_xref="PID:g2196920"

/translation="MAQSLALSLILVLAFGIPRTQGSDDGAQDCCLKYSQRKIPAKV

VRSYRKQEPSLGCIPAILFLPRKRSQAEKCADPKELWVQQLMQHLDKTPSPQKPAQG
 CRKDRGASKTGKKGKSGKCKRTERSQTPKGP"

BASE COUNT 218 a 255 c 216 g 139 t
 ORIGIN

```

1  ggacagagc agacatggc cagtcactg ctctgagcct cttatcctg gttctggcct
61  ttggcatccc caggacccaa ggcagtgatg gaggggctca ggactgttgc ctcaagtaca
121 gccaaaggaa gattcccgcc aaggttgctc gcagctaccg gaagcaggaa ccaagcttag
181 gctgctccat ccagctatc ctgttcttgc cccgcaagcg ctctcaggca gagctatgtg
241 cagacccaaa ggagctcttg gtgcagcagc tgatgcagca tctggacaag acaccatccc
301 cacagaaaac agcccagggc tgcaggaagg acaggggggc ctccaagact ggcaagaaag
361 gaaagggctc caaaggctgc aagaggactg agcggtcaca gaccctaaa gggccatagc
421 ccagttagca gcctggagcc ctggagaccc caccagcctc accagcgctt gaagcctgaa
481 cccaagatgc aagaaggagg ctatgctcag gggccctgga gcagccacc catgctggcc
541 ttgccacact ctttctcctg ctttaaccac cccatctgca ttcacagctc tcacctgca
601 tggtcagatc tgcccacagc aggccaggtc cagagagacc gaggaggag agtctccag
661 ggagcatgag aggaggcagc aggactgtcc cttgaagga gaatcatcag gacctggag
721 ctgatacggc tcccagtag accccacctc ttccttgtaa atatgattta tacctaactg
781 aataaaaagc tgttctgtct tcccacccaa aaaaaaaaaa aaaaaaaa

```

//
 LOCUS HSU88321 502 bp mRNA PRI 22-JUN-1998
 DEFINITION Human beta chemokine Exodus-3 mRNA, complete cds.
 ACCESSION U88321
 NID g2196921
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
 Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 502)
 AUTHORS Hromas, R.A., Gray, P., Klemsz, M., Fife, K. and Broxmeyer, H.
 TITLE DCCL chemokines represent a novel beta chemokine subfamily
 JOURNAL Unpublished
 REFERENCE 2 (bases 1 to 502)
 AUTHORS Hromas, R.A.
 TITLE Direct Submission
 JOURNAL Submitted (04-FEB-1997) Medicine, Indiana University Medical
 Center, 975 West Walnut, Indianapolis, IN 46202, USA
 REFERENCE 3 (bases 1 to 502)
 AUTHORS Hromas, R.A.
 TITLE Direct Submission
 JOURNAL Submitted (22-JUN-1998) Medicine, Indiana University Medical
 Center, 975 West Walnut, Indianapolis, IN 46202, USA
 REMARK Amino acid sequence updated by submitter
 FEATURES Location/Qualifiers
 source 1..502

```

/organism="Homo sapiens"
/note="PCR amplified from THP-1 cells"
/db_xref="taxon:9606"
/cell_line="THP-1"
/cell_type="monoblast"
/dev_stage="adult"
CDS
120..416
/note="Mip-3alpha/ELC/CKbeta11"
/codon_start=1
/product="beta chemokine Exodus-3"
/db_xref="PID:g3243080"

/translation="MALLLALLSLVLWTSAPTLSGTNDAEDCCLSVTQKPIPGYIVR

NFHYLLIKDGCVRPAVVFTTLRGRQLCAPDQPWVERIIQRLQRTSAKMKRRSS"
BASE COUNT      113 a      170 c      121 g      98 t
ORIGIN
    1 ctcacacctt gcatttcacc cctgcatccc atgcgccttg cagcctcaca cagatcctgc
   61 acacacccag acagctggcg ctcacacatt caccgttggc ctgcctctgt tcaccctcca
  121 tggccctgct actggccctc agcctgctgg ttctctggac ttccccagcc ccaactctga
  181 gtggcaccaa tgatgctgaa gactgctgcc tgtctgtgac ccagaaaccc atccctgggt
  241 acatcgtgag gaacttcac taccttctca tcaaggatgg ctgcagggtg cctgctgtag
  301 tgttcaccac actgaggggc cgccagctct gtgcaccccc agaccagccc tgggtagaac
  361 gcatcatcca gagactgcag aggacctcag ccaagatgaa gcgccgcagc agttaaccta
  421 tgaccgtgca gagggagccc cgagtcagg tcaagcattg tgaattatta ctaactggga
  481 acgaggacag aaggaaggac ag

//

LOCUS      HSU86358      879 bp      mRNA      PRI      11-SEP-1997
DEFINITION Human chemokine (TECK) mRNA, complete cds.
ACCESSION  U86358
NID        g2388626
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 879)
AUTHORS    Vicari,A.P., Figueroa,D.J., Hedrick,J.A., Foster,J.S.,
Singh,K.P.,
            Menon,S., Copeland,N.G., Gilbert,D.J., Jenkins,N.A., Bacon,K.B.
and
            Zlotnik,A.
TITLE      TECK: a novel cc chemokine specifically expressed by thymic
            dendritic cells and potentially involved in T cell development
JOURNAL    Immunology 7, 291-301 (1997)
REFERENCE  2 (bases 1 to 879)
AUTHORS    Vicari,A.P. and Zlotnik,A.
TITLE      Direct Submission
JOURNAL    Submitted (21-JAN-1997) Immunology, DNAX Research Institute,
901
            California Ave., Palo Alto, CA 94304, USA
FEATURES   Location/Qualifiers
            source      1..879
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /chromosome="4"
                        /tissue_type="thymus"
            gene        1..879
                        /gene="TECK"
            CDS         1..453
                        /gene="TECK"
                        /codon_start=1
                        /product="chemokine"
                        /db_xref="PID:g2388627"

/translation="MNLWLLACLIVAGFLGAWAPAVHTQGVFEDCCLAYHYPIGWAVLR

RAWTYRIQEVSGSCNLPAAIFYLPKRHRKVCNPKSREVQRAMKLLDARNKVFAKLHH
NMQTFQAGPHAVKKLSSGSKLSSSKFSNPISSSKRNVSLLISANSGL"
BASE COUNT      191 a      264 c      218 g      206 t

```

ORIGIN

```

1 atgaacctgt ggctcctggc ctgcctgggt gccggcttcc tgggagcctg ggcccccgct
61 gtccacaccc aagggtgtctt tgaggactgc tgcttggcct accactaccc cattgggtgg
121 gctgtgtctc ggcgcgcttg gacttacccg atccaggagg tgagcgggag ctgcaatctg
181 cctgctgcga tattctacct ccccaagaga cacaggaagg tgtgtgggaa ccccaaaagc
241 agggaggtgc agagagccat gaagctcctg gatgctcgaa ataaggtttt tgcaaaagctc
301 caccacaaca tgcagacctt ccaagcaggc cctcatgctg taaagaagtt gagtctctgga
361 aactccaagt tatcatcatc caagttagc aatcccatca gcagcagcaa gaggaatgtc
421 tccctcctga tatcagctaa ttcaggactg tgagccggct catttctggg ctccatcggc
481 acaggagggg ccggatcttt ctccgataaa accgtcgccc tacagaccca gctgtcccca
541 cgcctctgtc ttttgggtca agtcttaate cctgcacctg agttggctct ccctctgcac
601 cccaccacc tcctgcccgt ctggcaactg gaaagaagga gttggcctga ttttaacctt
661 ttgccgctcc gggaacacgc acaatcctgg gcagccagtg gctctttag agaaaactta
721 ggatacctct ctcaatttct gtttcttgcc gtccaccctg ggccatgcca gtgtgtctc
781 tgggtccctt ccaaaaatct ggtcattcaa ggatccctc ccaaggctat gcttttctat
841 aacttttaaa taaaccttgg ggggtgaatg gaataaaaa

```

//

LOCUS AB002409 852 bp mRNA PRI 15-AUG-1997

DEFINITION Homo sapiens mRNA for SLC, complete cds.

ACCESSION AB002409

NID g2335034

KEYWORDS SLC; mature ELC.

SOURCE Homo sapiens cDNA to mRNA.

ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae;
Homo.

REFERENCE 1 (bases 1 to 852)

AUTHORS Nomiya, H.

TITLE Direct Submission

JOURNAL Submitted (28-MAR-1997) to the DDBJ/EMBL/GenBank databases.
Hisayuki Nomiya, Kumamoto University Medical School,
Department of Biochemistry; Honjo 2-2-1, Kumamoto, Kumamoto 860, Japan
(E-mail:nomiyama@gpo.kumamoto-u.ac.jp, Tel:81-96-373-5063,
Fax:81-96-372-6140)

REFERENCE 2 (bases 1 to 852)

AUTHORS Nagira, M., Imai, T., Hieshima, K., Kusuda, J., Ridanpaa, M.,
Takagi, S., Nishimura, M., Kakizaki, M., Nomiya, H. and Yoshie, O.

TITLE Molecular Cloning of a Novel Human CC Chemokine Secondary
Lymphoid-Tissue Chemokine (SLC) That is an Efficient
Chemoattractant for Lymphocytes and Mapped to Chromosome 9p13

JOURNAL Unpublished (1997)

FEATURES

source Location/Qualifiers

1..852

/organism="Homo sapiens"

/db_xref="taxon:9606"

CDS

59..463

/codon_start=1

/product="SLC"

/db_xref="PID:d1022673"

/db_xref="PID:g2335035"

/translation="MAQSLALSLILVLAFGIPRTQSGDGAQDCCLKYSQRKIPAKV

VRSYRKQEPSLGCIPAILFLPRKRSQAEKADPKELWVQQLMQHLDKTPSPQKPAQG

CRKDRGASKTGKKGKSGKCKRTERSQTPKGP"

mat_peptide

<107..460

/product="mature ELC"

polyA_site

823..828

BASE COUNT 205 a 279 c 217 g 151 t

ORIGIN

```

1 cttgcagctg cccacctcac cctcagctct gccctcttac tcacctcta ccacagacat
61 ggctcagtcg ctggctctga gcctccttat cctggttctg gcctttggca tccccaggac
121 ccaaggcagt gatggagggg ctcaggactg ttgcctcaag tacagccaaa ggaagattcc
181 cgccaagggt gtccgcagct accggaagca ggaaccaagc ttaggctgct ccatccacgc
241 tatctgttct ttgccccgca agcgtcttca ggcagagcta tgtgcagacc caaaggagct
301 ctgggtgcag cagctgatgc agcatctgga caagacacca tccccacaga aaccagccca

```

```

361 gggctgcagg aaggacaggg gggcctccaa gactggcaag aaaggaaaagg gctccaaagg
421 ctgcaagagg actgagcggg cacagacccc taaaggggcca tagcccagtg agcagcctgg
481 agccctggag accccaccag cctcaccaac gcttgaagcc tgaacccaag atgcaagaag
541 gaggctatgc tcagggggccc tggagcagcc accccatgct ggccttgcca cactctttct
601 cctgctttaa ccaccccatc tgcattccca gctctaccct gcatggctga gctgcccaca
661 gcaggccagg tccagagaga ccgaggaggg agagtctccc agggagcatg agaggaggca
721 gcaggactgt ccccttgaag gagaatcatc aggaccctgg acctgatacg gctccccagt
781 acacccacc tcttccttgt aaatatgatt tatacctaac tgaataaaaa gctgttctgt
841 ctccccacc gc

```

//

```

LOCUS      AF055467      1481 bp      mRNA      PRI      06-AUG-1998
DEFINITION Homo sapiens monotactin-1 mRNA, complete cds.
ACCESSION  AF055467
NID        g3395775
KEYWORDS   .
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1481)
AUTHORS    Youn,B.S., Zhang,S., Broxmeyer,H.E., Antol,K., Fraser,M.J. Jr.,
            Hangoc,G. and Kwon,B.S.
TITLE      Isolation and characterization of LMC, a novel lymphocyte and
            monocyte chemoattractant human CC chemokine, with
myelosuppressive
            activity
JOURNAL     Biochem. Biophys. Res. Commun. 247 (2), 217-222 (1998)
MEDLINE     98308096
REFERENCE  2 (bases 1 to 1481)
AUTHORS    Youn,B.S. and Kwon,B.S.
TITLE      Direct Submission
JOURNAL     Submitted (24-MAR-1998) Microbiology and Immunology, Indiana
            University, School of Medicine, 605 Barnhill Dr. Medical
Science
            Bldg., Indianapolis, IN 46202, USA
FEATURES   Location/Qualifiers
            source          1..1481
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /chromosome="17"
            5'UTR          1..34
            CDS            35..397
                        /note="Mtn-1; LMC; lymphocyte and monocyte
chemoattractant
                        CC chemokine"
                        /codon_start=1
                        /product="monotactin-1"
                        /db_xref="PID:g3395776"

/translation="MKVSEAALSLVLILITSASRSQPKVPEWVNTPTSCCLKYYEK
VLPRRLVGYRKALNCHLPAILFVTKRNREVCTNPNDWVQEYIKDPNLPPLPTRNLS
TVKIIITAKNGQPQLLSQ"
            3'UTR          398..1481
BASE COUNT 412 a      362 c      302 g      405 t
ORIGIN      1 gcacgagctg aagctgtact gcctcgetga gaggatgaag gtctccgagg ctgccctgtc
            61 tctccttgtc ctcatcctta tcattacttc ggcttctcgc agccagccaa aagttcctga
            121 gtgggtgaac accccatcca cctgctgcct gaagtattat gagaaagtgt tgccaaggag
            181 actagtgggt ggatacagaa aggccctcaa ctgtcacctg ccagcaatca tcttcgtcac
            241 caagaggaac cgagaagtct gcaccaaccc caatgacgac tgggtccaag agtacatcaa
            301 ggatcccaac ctacctttgc tgcctaccag gaacttgccc acggttaaaa ttattacagc
            361 aaagaatggt caacccccagc tcctcaactc ccagtgatga ccaagcttta gtggaagccc
            421 ttgtttacag aagagagggg taaactatga aaacagggga agccttatta ggctgaaact
            481 agccagtcac attgagagaa gcagaacaat gatcaaaata aaggagaagt atttgaata
            541 ttttctcaat cttaggagga aataccaaag ttaagggacg tgggcagagg tacgctcttt
            601 tatttttata tttatatatt tttttttttg agataggggtc ttactctgtc acccaggtg
            661 gagtgcagtg gtgtgatctt ggtcacttg atcttggctc actgtaacct ccacctccca
            721 ggctcaagtg atcctccac cccaccctcc cgagttagctg ggactacagg cttgcgccac
            781 cacacctggc taatttttgt atttttggtg gagacgggat tctaccatgt tgcccaggct

```

```

841 ggtctcaaac tcgtgtgccc aagcaatcca cctgcctcag ccttccaaaa gtgctgggct
901 tacaggcgtg agccaccaca tccggccagt ccactcttaa tacacagaaa aatatatttc
961 acatccttct cctgctctct ttcaattcct cacttcacac cagtacacaa gccattctaa
1021 atacttagcc agtttccagc cttccagatg atctttgccc tctgggtctt gaccatttaa
1081 gagcccccata gaactcttga tttttcctgt ccactcttat gggatttttc tggatctata
1141 ttttcttcaa ttattctttc attttataat gcaacttttt cataggaagt ccggtaggga
1201 atattcacat taatcatttt tgcagagact ttgctagatc ctctcatatt ttgtcttctc
1261 cagggtggca ggggtacaga agtgcctgat tgggtttttt tttttttgag agagagagag
1321 aagaagaaga agaagagaca caaatctcta cctcccatgt taagctttgc aggacaggga
1381 aagaaagggt atgagacacg gctagggtaa actcttagtc caaaaccaa gcatgcaata
1441 aataaaactc ccttatttga caaaaaaaa aaaaaaaaaa a

```

//

```

LOCUS      HSRNAATAC      557 bp      RNA      PRI      06-JUL-1995
DEFINITION H.sapiens mRNA for ATAC protein.
ACCESSION  X86474
NID        g895846
KEYWORDS   ATAC gene.
SOURCE     human.
  ORGANISM Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 557)
AUTHORS   Muller,S., Dorner,B., Korthauer,U., Mages,H.W., D'Apuzzo,M.,
            Senger,G. and Kroczeck,R.A.
TITLE      Cloning of ATAC, an activation-induced, chemokine-related
molecule  exclusively expressed in CD8+ T lymphocytes
JOURNAL    Eur. J. Immunol. 25 (6), 1744-1748 (1995)
MEDLINE    95339892
REFERENCE  2 (bases 1 to 557)
AUTHORS   Kroczeck,R.A.
TITLE      Direct Submission
JOURNAL    Submitted (20-APR-1995) R.A. Kroczeck, Molecular Immunology,
            Robert-Koch-Institute, Nordufer 20, 13353 Berlin, FRG
FEATURES   Location/Qualifiers
  source    1..557
            /organism="Homo sapiens"
            /db_xref="taxon:9606"
            /tissue_type="peripheral blood"
            /cell_type="lymphocyte"
            /chromosome="1"
            /map="q23"
  gene      25..369
            /gene="ATAC"
  CDS       25..369
            /gene="ATAC"
            /codon_start=1
            /product="CD8+T cell specific protein"
            /db_xref="PID:g895847"
            /db_xref="SWISS-PROT:P47992"

/translation="MRLILALLGICSLTAYIVEGVGSEVSDKRTCVSLLTQRLPVSR

```

IKTYTITEGSLRAVIFITKRGLKVCADPQATWVRDVVRSMDRKSNTNRNMIQT KPTGT

QOSTNTAVTLTG"

polyA_signal 469..474

polyA_signal 534..539

BASE COUNT 157 a 139 c 112 g 149 t

ORIGIN

```

1 gcacagctca gcaggacctc agccatgaga cttctcatcc tggccctcct tggcatctgc
61 tctctcactg catacattgt ggaagggtga gggagtgaag tctcagataa gaggacctgt
121 gtgagcctca ctaccagcgc actgccggtt agcagaatca agacctacac catcacggaa
181 ggctccttga gagcagtaat ttttattacc aaacgtggcc taaaagtctg tgtgatccaa
241 caagccacat ggggtagaga cgtggtcagg agcatggaca ggaaatccaa caccagaaat
301 aacatgatcc agaccaagcc aacaggaacc cagcaatcga ccaatacagc tgtgactctg
361 actggctagt agtctctggc accctgtccg tctccagcca gccagctcat ttcactttac
421 acgctcatgg actgagttaa tactcgcctt ttatgaaagc actgcatgaa taaaattatt
481 cctttgtatt tttactttta aatgtcttct gtattcactt atatgttcta attaataaat
541 tatttattat taagaat

```

//

LOCUS HSU85767 563 bp mRNA PRI 01-APR-1997
 DEFINITION Human myeloid progenitor inhibitory factor-1 MPIF-1 mRNA,
 complete
 cds.
 ACCESSION U85767
 NID g1916249
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 563)
 AUTHORS Patel,V.P., Kreider,B.L., Li,Y., Li,H., Leung,K., Salcedo,T.,
 Nardelli,B., Pippalla,V., Gentz,S., Thotakura,R., Parmelee,D.,
 Gentz,R. and Garotta,G.
 TITLE Molecular and functional characterization of two novel human C-
 C chemokines as inhibitors of two distinct classes of myeloid
 progenitors
 JOURNAL J. Exp. Med. (1997) In press
 REFERENCE 2 (bases 1 to 563)
 AUTHORS Li,H. and Patel,V.P.
 TITLE Direct Submission
 JOURNAL Submitted (17-JAN-1997) Cell Biology, Human Genome Sciences,
 9410 Keywest Ave., Rockville, MD 20850, USA
 FEATURES Location/Qualifiers
 source 1..563
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 CDS 31..393
 /note="myeloid progenitor inhibitory factor-1"
 /codon_start=1
 /product="MPIF-1"
 /db_xref="PID:g1916250"
 /translation="MKVSVAAALSCLMLVTALGSQARVTKDAETEFMMSKLPLENPVLL
 DRFHATSADCCISYTPRSIPCSLLESYFETNSECSKPGVIFLTKKGRRFCANPSDKQV
 QVCMRMLKLDTRIKTRKN"
 BASE COUNT 164 a 143 c 117 g 139 t
 ORIGIN
 1 ctcagccagc cctgcctgcc caccaggagg atgaaggtct ccgtggctgc cctctcctgc
 61 ctcattgctt ttactgccct tggatcccag gcccggttca caaaagatgc agagacagag
 121 ttcatgatgt caaagcttcc attggaaaat ccagtacttc tggacagatt ccattgctact
 181 agtgcctgact gctgcatttc ctacacccca cgaagcatcc cgtgttcact cctggagagt
 241 tactttgaaa cgaacagcga gtgtctcaag ccgggtgtca tcttcctcac caagaagggg
 301 cgacgtttct gtgccaaccc cagtgataag caagttcagg tttgcatgag aatgctgaag
 361 ctggacacac ggatcaagac caggaagaat tgaacttgtc aaggtgaagg gacacaagtt
 421 gccagccacc aactttcttg cctcaactac ctctctgaat tattttttta agaagcattt
 481 attcttgtgt tctggattta gagcaattca tctaataaac agtttctcac ttttaaaaaa
 541 aaaaaaaaaa aaaaaaaaaa aaa
 //

LOCUS HSU85768 360 bp mRNA PRI 01-APR-1997
 DEFINITION Human myeloid progenitor inhibitory factor-1 MPIF-2 mRNA,
 complete
 cds.
 ACCESSION U85768
 NID g1916251
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 360)
 AUTHORS Patel,V.P., Kreider,B.L., Li,Y., Li,H., Leung,K., Salcedo,T.,
 Nardelli,B., Pippalla,V., Gentz,S., Thotakura,R., Parmelee,D.,
 Gentz,R. and Garotta,G.
 TITLE Molecular and functional characterization of two novel human C-

C

chemokines as inhibitors of two distinct classes of myeloid progenitors

J. Exp. Med. (1997) In press

REFERENCE 2 (bases 1 to 360)

AUTHORS Li, H. and Patel, V. P.

TITLE Direct Submission

JOURNAL Submitted (17-JAN-1997) Cell Biology, Human Genome Sciences, 9410 Keywest Ave., Rockville, MD 20850, USA

FEATURES Location/Qualifiers

source 1..360

/organism="Homo sapiens"

/db_xref="taxon:9606"

CDS 1..360

/note="myeloid progenitor inhibitory factor-2"

/codon_start=1

/product="MPIF-2"

/db_xref="PID:g1916252"

/translation="MAGLMTIVTSLFLGVCAHHIIP TGSVVIPSPCCMFFVSKRIPE

NRVVS YQLSSRSTCLKGGVIFTTKKGQQFCGDPKQEWVQRYMKNLDAKQKKASPRARA

VAVKGPVQRYPGNQTTTC"

BASE COUNT 85 a 106 c 96 g 73 t

ORIGIN

1 atggcaggcc tgaatgacct agtaaccagc cttctgttcc ttggtgtctg tgcccaccac

61 atcatcccta cgggctctgt ggatcatacc tctccctgct gcatgttctt tggttccaag

121 agaattcctg agaaccgagt ggatcagctac cagctgtcca gcaggagcac atgcctcaag

181 ggaggagtga tcttcaccac caagaagggc cagcagttct gtggcgaccc caagcaggag

241 tgggtccaga ggtacatgaa gaacctggac gccaaagcaga agaaggcttc ccctagggcc

301 agggcagtggt ctgtcaaggg ccctgtccag agatatcctg gcaaccaaac cacctgctaa

//

LOCUS HUMSDF1A 1847 bp mRNA PRI 26-DEC-1996

DEFINITION Human pre-B cell stimulating factor homologue (SDF1a) mRNA, complete cds.

ACCESSION L36034

NID g1220363

KEYWORDS intercrine; intercrine CXC subfamily; pre-B cell stimulating factor

homologue; alpha-chemokine.

SOURCE human.

ORGANISM Homo sapiens

Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata; Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 1847)

AUTHORS Shirozu, M., Nakano, T., Inazawa, J., Tashiro, K., Tada, H., Shinohara, T. and Honjo, T.

TITLE Structure and chromosomal localization of the human stromal cell-derived factor 1 (SDF1) gene

JOURNAL Genomics 28 (3), 495-500 (1995)

MEDLINE 96039262

FEATURES Location/Qualifiers

source 1..1847

/organism="Homo sapiens"

/db_xref="taxon:9606"

/clone="h5"

/cell_line="FLEB14-14"

sig_peptide 80..142

/gene="SDF1a"

CDS 80..349

/codon_start=1

/product="pre-B cell stimulating factor homologue"

/db_xref="PID:g1220364"

/translation="MNAKVVVVLVLVTALCLSDGKPVLSYRCP RFFESHVARANV

KHLKILNTPNCALQIVARLKNNNRQVCIDPKLKIQYLEKALNK"

gene 80..346

/gene="SDF1a"

mat_peptide 143..346

```

/ gene="SDF1a"
/ product="pre-B cell stimulating factor homologue"
BASE COUNT      459 a      471 c      417 g      500 t
ORIGIN
1  tctccgctcag cgcgattgcc cgctcggcgt cgggcccccg acccgtgctc gtccgccccg
61  cgcgccccgc gcccgcgcca tgaacgcca ggtcgtggtc gtgctgggtc tcgtgctgac
121 cgcgctctgc ctcagcgacg ggaagcccg cagcctgagc tacagatgcc catgccgatt
181 cttcgaaagc catgttgcca gagccaacgt caagcatctc aaaattctca acactccaaa
241 ctgtgccctt cagattgtag cccggctgaa gaacaacaac agacaagtgt gcattgacct
301 gaagctaaag tggattcagg agtacctgga gaaagcttta aacaagtaag cacaacagcc
361 aaaaaggact ttccgctaga cccactcgag gaaaactaaa accttgtgag agatgaaagg
421 gcaaagacgt gggggagggg gccttaacca tgaggaccag gtgtgtgtgt ggggtgggca
481 cattgatctg ggatcgggcc tgagggttgc agcatttaga ccctgcattt atagcatacg
541 gtatgatatt gcagcttata ttcatccatg ccctgtacct gtgcacgttg gaacttttat
601 tactgggggt tttctaagaa agaaattgta ttatcaacag cattttcaag cagttaaagg
661 cttcatgatc atcacatca tcatcattct cattctcatt ttttaaatac acgagtactt
721 caagatctga atttggcttg tttggagcat ctccctctgt cccctgggga gtctgggcac
781 agtcagggtg tggcttaaca gggagctgga aaaagtgtcc tttcttcaga cactgaggct
841 cccgcagcag cgccctccc aagaggaagg cctctgtggc actcagatac cgactggggc
901 tggggcgccg ccactgcctt cactctctct ttcaaacctc agtgattggc tctgtgggct
961 ccattgtaga gccactatta ctgggactgt ctcagagacc cctctcccag ctattcctac
1021 tctctccccg actccgagag catgcttaat cttgcttctg cttctcattt ctgtagcctg
1081 atcagcgccg caccagcccg gaagagggtg attgctgggg ctcgtgccct gcacccctct
1141 cctcccaggg cctgccccac agctcggggc ctctgtgaga tccgtctttg gcctcctcca
1201 gaattggagc tggccctctc tggggatgtg taatggtccc cctgcttacc cgcaaaagac
1261 aagctctttc agaatcaaat gcaattttaa atctgagagc tcgcttgagt gactggggtt
1321 gtgattgcct ctgaagccta tgtatgccat ggaggcacta acaaactctg aggtttccga
1381 aatcagaagc gaaaaaatca gtgaataaac catcatcttg ccactacccc ctctgaagc
1441 cacagcaggg gttcagggtc caatcagaac tgttggcaag gtgacatttc catgcataga
1501 tgcgatccac agaaggctct ggtgggtatt gtaacttttt gcaaggcatt tttttatata
1561 tatttttgtg cacatttttt ttacgattc tttagaaaac aaatgtattt caaaatatat
1621 ttatagtcga acaagtcata tatatgaatg agagccatat gaatgtcagt agttttactt
1681 tctctattat ctcaaactac tggcaatttg taaagaaata tatatgatat ataatgtga
1741 ttgcagcttt tcaatgttag ccacagtgtg ttttttctt tgtactaaaa ttgtatcaaa
1801 tgtgacatta tatgcactag caataaaatg ctaattgttt catggta

```

//

```

LOCUS      HUMSDF1B      3524 bp      mRNA      PRI      26-DEC-1996
DEFINITION Human pre-B cell stimulating factor homologue (SDF1b) mRNA,
            complete cds.
ACCESSION  L36033
NID        g1220365
KEYWORDS   intercrine; intercrine CXC subfamily; pre-B cell stimulating
factor
SOURCE     human.
ORGANISM   Homo sapiens
REFERENCE  1 (bases 1 to 3524)
AUTHORS    Shirozu,M., Nakano,T., Inazawa,J., Tashiro,K., Tada,H.,
            Shinohara,T. and Honjo,T.
TITLE      Structure and chromosomal localization of the human stromal
            cell-derived factor 1 (SDF1) gene
JOURNAL    Genomics 28 (3), 495-500 (1995)
MEDLINE    96039262
FEATURES   Location/Qualifiers
            source          1..3524
                               /organism="Homo sapiens"
                               /db_xref="taxon:9606"
                               /clone="h17"
                               /cell_line="FLEB14-14"
            sig_peptide     80..142
                               /gene="SDF1b"
            CDS              80..361
                               /codon_start=1
                               /product="pre-B cell stimulating factor homologue"
                               /db_xref="PID:g1220366"

```

```

/translation="MNAKVVVVVLVLVLTALCLSDGKPVSLSYRCPCRFFESHVARANV
KHLKILNTPNCALQIVARLKNNNRQVCIDPKLWIIQYLEKALNKRFKM"

```

```

gene      80..358
          /gene="SDF1b"
mat_peptide 143..358
          /gene="SDF1b"
          /product="pre-B cell stimulating factor homologue"
BASE COUNT 903 a 886 c 793 g 942 t
ORIGIN
1  tctccgtcag cgcattgcc cgctggcgt cgggcccccg acccgtgctc gtcggcccg
61  ccgcccgcgc gccgcgcgca tgaacgccaa ggtcgtgggtc gtgctgggtcc tcgtgctgac
121 cgcgctctgc ctcagcgagc ggaagcccgt cagcctgagc tacagatgcc catgccgatt
181 ctcgaaaagc catgttgcca gagccaacgt caagcatctc aaaattctca acactccaaa
241 ctgtgccctt cagattgtag cccggctgaa gaacaacaac agacaagtgt gcattgacct
301 gaagctaaag tggattcagg agtacctgga gaaagcttta aacaagaggt tcaagatgtg
361 agaggggtcag acgcttgagg aacccttaca gtaggagccc agctctgaaa ccagtgttag
421 ggaagggcct gccacagcct cccctgccag ggcaaggccc caggcattgc caagggcttt
481 gttttgcaca ctttgccata ttttcaccat ttgattatgt agcaaaatac atgacattta
541 tttttcattt agtttgatta ttcagtgta ctggcgacac gttagcagctt agactaaggc
601 cattattgta cttgccttat tagagtgtct tcccacggag ccactcctct gactcagggc
661 tcttgggttt tgtattctct gagctgtgca ggtggggaga ctgggctgag ggagcctggc
721 cccatgggtc gccctagggt ggagagccac caagagggac gcctgggggt gccaggacca
781 gtcaacctgg gcaaagccta gtgaaggctt ctctctgtgg gatgggatgg tggagggcca
841 catgggaggg tcacccctct ctccatccac atgggagccg ggtctgcctc tctggggagg
901 gcagcagggc taccctgagc tgaggcagca gtgtgaggcc agggcagagt gagaccagc
961 cctcatcccg agcacctcca catcctccac gttctgctca tcattctctg tctcatccat
1021 catcatgtgt gtccacgact gtctccatgg ccccgcaaaa ggactctcag gaccaagct
1081 ttcattgttaa ctgtgcacca agcaggaaat gaaaatgtct tgtgttacct gaaaacactg
1141 tgcacatctg tgtcttgtgt ggaatattgt ccattgtcca atcctatgtt tttgtcaaaa
1201 gccagcgtcc tcctctgtga ccaatgtctt gatgcatgca ctgttcccc tgtgcagccg
1261 ctgagcgagg agatgctcct tgggcccctt gagtgcagtc ctgatcagag ccgtggctct
1321 ttggggtgaa ctaccttggt tccccactg atcacaaaaa catggtgggt ccatgggcag
1381 agcccaaggg aattcgggtg gcaccagggt tgacccaga ggattgctgc cccatcagtg
1441 ctccctcaca tgtcagtacc ttcaactag ggccaagccc agcactgctt gaggaaaaa
1501 agcattcaca acttgttttt ggttttttaa acccagtcca caaaataacc aatcctggac
1561 atgaagattc tttcccaatt cacatctaac ctcatcttct tcaccatttg gcaatgccat
1621 catctcctgc cttcctcctg ggccctctct gctctgcgtg tcacctgtgc tctgggcct
1681 tcccacagga catttctcta agagaacaat gtgctatgtg aagagtaagt caacctgcct
1741 gacatttgga gtgttcccc cccactgagg gcagtcgata gagctgtatt aagccactta
1801 aaatgttcac ttttgacaaa ggcaagcact tgtgggtttt tgttttgttt ttcattcagt
1861 ctacggaata cttttgcctt ttgattaaag actccagtta aaaaaattt taatgaagaa
1921 agtggaatac aaggaagtca aagcaaggaa actatgtaac atgttaggaag taggaagtaa
1981 attatagtga tgaatcttgg aattgttaact gttcgtgaat ttaataatct gtaggtaata
2041 tagtaacatg tgttaagtat ttccataagt atttcaaat ggagcttcat gccagaaggc
2101 aaacccatca acaaaaattg tcccttaaac aaaaattaaa atcctcaatc cagctatgtt
2161 atattgaaaa aatagagcct gagggatcct tactagttaa aaagatacag aactcttcca
2221 aaaccttttg aaattaacct ctactatac cagtataatt gagttttcag tggggcagtc
2281 attatccagg taatccaaga tatttttaaa tctgtcacgt agaacttggg tgtacctgac
2341 cccaatccat gaaccaagac cattgaattc ttggttgagg aaacaaacat gacctaaat
2401 ctgactaca gtcaggaaa gaaatcatttc tatttctcct ccatgggaga aaatagataa
2461 tagtagaaac tgcaggaaa atattttgca taacaattcc tctactaaca atcagctcct
2521 tcttgagac tgcccagcta aagcaatatg catttaataa cagtcttcca tttgcaaggg
2581 aaaagtctct tgtaatccga atctcttttt gctttcgaac tgctagtcaa gtgcgtccac
2641 gagctgttta ctagggatcc ctcatctgtc cctccgggac ctggtgtgac ctctacctga
2701 cactcccttg ggtccctgtt aacctcttca gaggcctctg ctgccagctc tgtatcagga
2761 cccagaggaa ggggccagag gctcgttgac tggctgtgtg ttgggattga gtctgtgcca
2821 cgtgtatgtg ctgtgtgtgt tccccctctg tccaggcact gagataccag cgaggaggct
2881 ccagagggca cttctgttgt tattagagat tacctcctga gaaaaaagct tccgttggg
2941 gcagaggggc tgaatagcag aaggttgcac ctccccaac cttagatgtt ctaagtcttt
3001 ccattggatc tcattggacc ctccatggtg gtgatcgtct gactgggtgt atcacctggg
3061 gtcctctgac tgggagttga tcgcctttcc cagggtgtac acccttttcc agctggatga
3121 gaatttgagt gctctgatcc ctctacagag cttccctgac tcattctgaa ggagccccat
3181 tcctgggaaa tattccctag aaacttccaa atcccctaag cagaccactg ataaaaaccat
3241 gtagaaaaat tgttattttg caacctcgct ggactctcag tctctgagca gtgaatgatt
3301 cagtgttaaa tgtgatgaat actgtatttt gtattgtttc aagtgcattc cccagataat
3361 gtgaaaaatg tccaggagaa ggccaattcc tatacgagc gtgcttttaa aataaataa
3421 gaaacaactc tttgagaaac aacaatttct actttgaagt cataccaatg aaaaaatgta
3481 tatgcactta taattttcct aataaagttc tgtactcaaa tgta

```

//

```

LOCUS      HSJ002211 663 bp mRNA PRI 11-MAR-1998
DEFINITION Homo sapiens cDNA for a CXC chemokine.
ACCESSION AJ002211

```

NID g2832410
 KEYWORDS CXC chemokine.
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
 Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 663)
 AUTHORS Legler,D.F., Loetscher,M., Roos,R.S., Clark-Lewis,I.,
 Baggiolini,M.
 and Moser,B.
 TITLE B cell-attracting chemokine 1, a human CXC chemokine expressed
 in lymphoid tissues, selectively attracts B lymphocytes via
 BLR1/CXCR5
 JOURNAL J. Exp. Med. 187 (4), 655-660 (1998)
 MEDLINE 98130629
 REFERENCE 2 (bases 1 to 663)
 AUTHORS Moser,B.
 TITLE Direct Submission
 JOURNAL Submitted (05-NOV-1997) Moser B., University of Bern, Theodor
 Kocher Institute, Freiestrasse 1, CH-3012 Bern, SWITZERLAND
 FEATURES Location/Qualifiers
 source 1..663
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /cell_type="PBL"
 sig_peptide 35..100
 /gene="BCA-1"
 CDS 35..364
 /gene="BCA-1"
 /codon_start=1
 /product="CXC chemokine"
 /db_xref="PID:e1249325"
 /db_xref="PID:g2832411"
 /translation="MKFISTSLLLMLLVSSLSPVQGVLEVVYTSRLRCRCVQESSVFIP
 RRFIDRIQILPRGNGCPRKEIIVWKNKSIVCVDPAEWIQRMMEVLRKRSSSTLPVP
 VFKRKIP"
 gene 35..364
 /gene="BCA-1"
 mat_peptide 101..361
 /gene="BCA-1"
 BASE COUNT 176 a 136 c 145 g 198 t 8 others
 ORIGIN
 1 cagagctcaa gtctgaactc tacctccaga cagaatgaag ttcattctga catctctgct
 61 tctcatgctg ctggctcagca gcctctctcc agtccaaggt gttctggagg tctattacac
 121 aagcttgagg tgtagatgtg tccaagagag ctccagcttt atccctagac gcttcattga
 181 tcgaattcaa atcttgcccc gtgggaatgg ttgtccaaga aaagaaatca tagtctggaa
 241 gaagaacaag tcaattgtgt gtgtggacc ccaagctgaa tggatacaaa gaatgatgga
 301 agtattgaga aaaagaagtt cttcaactct accagttcca gtgtttaaga gaaagattcc
 361 ctgatgctga tatttccact aagaacacct gcattcttcc cttatccctg ctctgggatt
 421 ttagttttgt gcttagttaa atcttttcca gggagaaaaga acttcccat acaaataagg
 481 catgaggact atgtaaaaaa aaccttgcag gagctggatg gggggccaaa ctcaagcttc
 541 tttcactcca caggcaccct attntacact tgggggtttt gcnttcttn tttcntcagg
 601 gggggggaaa gtttcttttg gaaantagtt nttccagttt ttaggtatta cagggttntt
 661 ttt
 //
 LOCUS HSHUMIG 2545 bp RNA PRI 16-NOV-1993
 DEFINITION H.sapiens Humig mRNA.
 ACCESSION X72755 S60728
 NID g311375
 KEYWORDS chemokine; cytokine; Humig gene; secreted protein.
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 2545)
 AUTHORS Farber,J.M.
 TITLE Direct Submission
 JOURNAL Submitted (22-MAR-1993) J.M. Farber, Johns Hopkins Univ. School

of
 USA Medicine, Ross 1147, 720 Rutland Avenue, Baltimore, MD 21205,
 REFERENCE 2 (bases 1 to 2545)
 AUTHORS Farber, J.M.
 TITLE HuMig: a new human member of the chemokine family of cytokines
 JOURNAL Biochem. Biophys. Res. Commun. 192 (1), 223-230 (1993)
 MEDLINE 93236577
 FEATURES Location/Qualifiers
 source 1..2545
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /germline
 /dev_stage="child"
 /tissue_type="leukaemia"
 /cell_type="monocyte"
 /cell_line="THP-1"
 /clone_lib="THP-1/IFN-gamma cDNA"
 /clone="H-1-3"
 misc_feature 13..19
 /note="cis-acting element; putative"
 gene 40..417
 /gene="Humig"
 CDS 40..417
 /gene="Humig"
 /codon_start=1
 /db_xref="PID:g311376"
 /db_xref="SWISS-PROT:Q07325"

/translation="MKKSGVLFLLGIILLVLIGVQGPVVRKGRCSISTNQGTIHLQ

SLKDLKQFAPSPSCEKIEIIATLKNVQTCNLNPDADVKEIKKWEKQVSQKKQKNG
 KKHQKKKVLKVRKSQRSRQKKTT"

BASE COUNT 755 a 581 c 457 g 752 t
 ORIGIN

```

1 atccaatata ggagtgaactt ggaactccat tctatcacta tgaagaaaaa tgggtgttctt
61 ttcctcttgg gcatcatctt gctgggtctg attggagtgc aaggaacccc agtagtgaga
121 aaggggtcgct gttcctgcat cagcaccaac caagggacta tccacctaca atccttgaaa
181 gaccttaaac aatttgcccc aagcccttcc tgcgagaaaa ttgaaatcat tgctacactg
241 aagaatggag ttcaaacatg tctaaaccca gattcagcag atgtgaagga actgattaaa
301 aagtgggaga aacaggtcag ccaaaagaaa aagcaaaaga atgggaaaaa acatcaaaaa
361 aagaaagtgc tgaagtgtcg aaaatctcaa cgttctcgtc aaaagaagac tacataagag
421 accacttcac caataagtat tctgtgttaa aaatgttcta ttttaattat accgctatca
481 ttccaaagga ggatggcata taatacaaag gcttattaat ttgactagaa aattttaaac
541 attactctga aattgtaact aaagttagaa agttgatttt aagaatccaa acgttaagaa
601 ttgttaaagg ctatgattgt ctttgttctt ctaccacca ccagttgaat ttcacatgc
661 ttaaggccat gattttagca ataccatgt ctacacagat gttcacccaa ccacatccca
721 ctcaaacag ctgcctggaa gaggcgcct aggcctccac gtactgcagc ctccagagag
781 tatctgaggc acatgtcagc aagtcctaag cctgttagca tgctggtagc ccaagcagtt
841 tgaaaattgag ctggacctca ccaagctgct gtggccatca acctctgtat ttgaatcagc
901 ctacaggcct cacacacaat gtgtctgaga gattcatgct gattgttatt gggtatcacc
961 actggagatc accagtgtgt ggctttcaga gcctccttcc tggcttttga agccatgtga
1021 ttccatcttg cccgctcagg ctgaccactt tatttctttt tgttccccct tgcttcatc
1081 aagtcagctc ttctccatcc taccacaatg cagtgccttt ctctctcca gtgcacctgt
1141 catatgctct gatttatctg agtcaactcc tttctcatct tgtccccaac accccacaga
1201 agtgctttct tctcccaatt catcctcact cagtccagct tagttcaagt cctgcctctt
1261 aaataaacct ttttggacac acaaattatc ttaaaactcc tgtttcactt ggttcagtag
1321 cacatgggtg aacactcaat ggttaactaa ttcttgggtg tttatcctat ctctccaacc
1381 agattgtcag ctcccttgagg gcaagagcca cagtatatct ccctgtttct tccacagtgc
1441 ctaataatac tgtggaacta ggttttaata attttttaat tgatgttgtt atgggcagga
1501 tggcaaccag accattgtct cagagcaggt gctggctctt tccctggctac tccatgtttg
1561 ctagcctctg gtaacctctt acttattatc ttcaggacac tcaactacag gaccagggat
1621 gatgcaacat ccttgtcttt ttatgacagg atgtttgtct agcttctcca acaataagaa
1681 gcacgtggta aaacacttgc ggatattctg gactgttttt aaaaaatata cagtttaccg
1741 aaaaatcatat aatcttacia tgaaaaggac tttatagatc agccagtgc caaccttttc
1801 ccaaccatac aaaaattcct ttcccgaag gaaaagggtc ttctcaataa gcctcagctt
1861 tctaagatct aacaagatag ccaccgagat ccttatcgaa actcatttta ggcaaatatg
1921 agttttattg tccgtttact tgtttcagag tttgtattgt gattatcaat taccacacca
1981 tctcccatga agaaagggaa cgggtgaagta ctaagcgcta gaggaagcag ccaagtcggt
2041 tagtggaagc atgattgggtg ccagtttagc ctctgcagga tgtggaaacc tccttcagg
2101 ggaggttcag tgaattgtgt aggagaggtt gtctgtggcc agaattttaa cctataactca

```

```

2161 ctttcccaaa ttgaatcact gctcacactg ctgatgattt agagtgcgtg ccgggtggaga
2221 tcccccccgga acgtcttatc taatcatgaa actccctagt tccttcatgt aacttccctg
2281 aaaaaatctaa gtgtttcata aatttgagag tctgtgaccc atttaccttg catctcacag
2341 gtagacagta tataactaac aaccaaagac tacatattgt cactgacaca caggttataa
2401 tcatttatca tatatatata tacatgcata cactctcaaa gcaaataatt ttctacttca
2461 aaacagtatt gacttgtata ccttgtaatt tgaaatattt tctttgttaa aatagaatgg
2521 tatcaataaa tagaccatta atcag

```

//

```

LOCUS      HSHUMIG      2545 bp      RNA      PRI      16-NOV-1993
DEFINITION H.sapiens Humig mRNA.
ACCESSION  X72755 S60728
NID        g311375
KEYWORDS   chemokine; cytokine; Humig gene; secreted protein.
SOURCE     human.
  ORGANISM Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 2545)
AUTHORS    Farber, J.M.
TITLE      Direct Submission
JOURNAL    Submitted (22-MAR-1993) J.M. Farber, Johns Hopkins Univ. School
of
            Medicine, Ross 1147, 720 Rutland Avenue, Baltimore, MD 21205,
USA
REFERENCE  2 (bases 1 to 2545)
AUTHORS    Farber, J.M.
TITLE      HuMig: a new human member of the chemokine family of cytokines
JOURNAL    Biochem. Biophys. Res. Commun. 192 (1), 223-230 (1993)
MEDLINE    93236577
FEATURES   Location/Qualifiers
            source          1..2545
                           /organism="Homo sapiens"
                           /db_xref="taxon:9606"
                           /germline
                           /dev_stage="child"
                           /tissue_type="leukaemia"
                           /cell_type="monocyte"
                           /cell_line="THP-1"
                           /clone_lib="THP-1/IFN-gamma cDNA"
                           /clone="H-1-3"
            misc_feature    13..19
                           /note="cis-acting element; putative"
            gene            40..417
                           /gene="Humig"
            CDS             40..417
                           /gene="Humig"
                           /codon_start=1
                           /db_xref="PID:g311376"
                           /db_xref="SWISS-PROT:Q07325"

```

/translation="MKKSGVLFLLGIILLVLIGVQGTPVVRKGRCSISTNQGTIHLQ

SLKDLKQFAPSPSCEKIEIIATLKNQVQTCNLNPDADVKELIKKEQVVSQKKKQKNG
KKHQKKKVLKVRKSQRSQKKT"

BASE COUNT 755 a 581 c 457 g 752 t

ORIGIN

```

1 atccaataca ggagtgactt ggaactccat tctatcacta tgaagaaaag tgggtgttctt
61 ttcctcttgg gcatcatctt gctggttctg attggagtgc aaggaacccc agtagtgaga
121 aagggtegct gttcctgcat cagcaccaac caagggacta tccacctaca atccttgaaa
181 gaccttaaac aatttgcccc aagcccttcc tgcgagaaaa ttgaaatcat tgctacactg
241 aagaatggag ttcaaacatg tctaaaccca gattcagcag atgtgaagga actgattaaa
301 aagtgggaga aacagggtcag ccaaaagaaa aagcaaaaaga atgggaaaaa acatcaaaaa
361 aagaaagtgc tgaaagtctg aaaatctcaa cgttctcgtc aaaagaagac tacataagag
421 accacttcac caataagtat tctgtgttaa aaatgttcta ttttaattat accgctatca
481 ttccaaagga ggatggcata taatacaaaag gtttattaat tgactagaa aatttaaaac
541 attactctga aattgttaact aaagttagaa agttgatttt aagaatccaa acgttaagaa
601 ttgttaaagg ctatgattgt ctttgttctt ctaccaccca ccagttgaat ttcacatgc
661 ttaagggcat gatttttagca ataccatgt ctacacagat gttcacccaa ccacatccca
721 ctcaaacag ctgcctggaa gagcagccct aggcctccac gtactgcagc ctccagagag
781 tatctgagga acatgtcagc aagtcctaag cctgttagca tgctggtgag ccaagcagtt

```

```

841 tgaattgag ctggacctca ccaagctgct gtggccatca acctctgtat ttgaatcagc
901 ctacaggcct cacacacaat gtgtctgaga gattcatgct gattgttatt ggggtaccac
961 actggagatc accagtgtgt ggctttcaga gcctcctttc tggcttttga agccatgtga
1021 ttccatcttg cccgctcagg ctgaccactt tatttctttt tgttccctt tgcttcattc
1081 aagtcagctc ttctccatcc taccacaatg cagtgccttt cttctctcca gtgcacctgt
1141 catatgctct gatttatctg agtcaactcc tttctcatct tgtccccaac acccacacaga
1201 agtgctttct tctcccaatt catcctcact cagtccagct tagttcaagt cctgcctctt
1261 aaataaacct ttttggacac acaaattatc ttaaaactcc tgtttcactt gggtcagtac
1321 cacatgggtg aacactcaat ggtaactaa ttcttgggtg tttatcctat ctctccaacc
1381 agattgtcag ctcttgagg gcaagagcca cagtatatct cctgtttct tccacagtgc
1441 ctaataatac tgtggaacta gggtttaata attttttaat tgatgttgtt atgggcagga
1501 tggcaaccag accattgtct cagagcaggt gctggctctt tcttggtac tccatgttgg
1561 ctgacctctg gtaacctctt acttattatc ttcaggacac tcaactacag gaccagggat
1621 gatgaacat ccttgtcttt ttatgacagg atgtttgtct agcttctcca acaataagaa
1681 gcacgtggta aaacacttgc ggatattctg gactgttttt aaaaaatata cagtttaccg
1741 aaaatcatat aatcttataa tgaaaaggac tttatagatc agccagtgc caaccttttc
1801 ccaaccatac aaaaattcct tttcccgaag gaaaagggtc ttctcaataa gcctcagctt
1861 tctaagatct aacaagatag ccaccgagat ccttatcgaa actcatttta ggcaaatatg
1921 agttttattg tccgtttact tgtttcagag tttgtattgt gattatcaat taccacacca
1981 tctcccatga agaaagggaa cgggtgaagta ctaagcgcta gaggaagcag ccaagtcggt
2041 tagtggaagc atgattgggtg cccagttagc ctctgcagga tgtggaaacc tccctccagg
2101 ggaggttcag tgaattgtgt aggagaggtt gctctgtggc agaattttaa cctatactca
2161 ctttcccaaa ttgaatcact gctcacactg ctgatgattt agagtgtgt ccggtggaga
2221 tcccacccga acgtcttctc taatcatgaa actccctagt tcttctatgt aacttcctg
2281 aaaaatctaa gtgtttcata aatttgagag tctgtgaccc acttaccttg catctcacag
2341 gtagacagta tataactaac aaccaaagac tacatattgt cactgacaca caggtataa
2401 tcatttatca tataataaca tacatgcata cactctcaaa gcaataaatt tttcacttca
2461 aaacagtatt gacttgata ccttgtaatt tgaaatatct tctttgttaa aatagaatgg
2521 tatcaataaa tagaccatta atcag

```

//

```

LOCUS      AF002985      995 bp      mRNA      PRI      01-NOV-1997
DEFINITION Homo sapiens putative alpha chemokine (H174) mRNA, complete
cds.
ACCESSION  AF002985
NID        g2580585
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 995)
AUTHORS    Jacobs,K.A., Collins-Racie,L.A., Colbert,M., Duckett,M.,
            Golden-Fleet,M., Kelleher,K., Kriz,R., LaVallie,E.R.,
            Merberg,D.,
            Spaulding,V., Stover,J., Williamson,M.J. and McCoy,J.M.
TITLE      A genetic selection for isolating cDNAs encoding secreted
proteins
JOURNAL    Gene 198 (1-2), 289-296 (1997)
MEDLINE    98036061
REFERENCE  2 (bases 1 to 995)
AUTHORS    Jacobs,K.A., Collins-Racie,L.A., Colbert,M., Duckett,M.,
            Golden-Fleet,M., Kelleher,K., Kriz,R., LaVallie,E.R.,
            Merberg,D.,
            Spaulding,V., Stover,J., Williamson,M.J. and McCoy,J.M.
TITLE      Direct Submission
JOURNAL    Submitted (07-MAY-1997) Genetics Institute, 87 Cambridge Park
            Drive, Cambridge, MA 02140, USA
FEATURES   Location/Qualifiers
            source      1..995
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /cell_type="PHA and PMA activated human peripheral
blood
                        mononuclear cells"
            gene        1..995
                        /gene="H174"
            CDS          88..372
                        /gene="H174"
                        /codon_start=1
                        /product="putative alpha chemokine"

```


/db_xref="PID:g2580586"

/translation="MSVKGMAIALAVILCATVVQGFPMFKRGRCLCIGPGVKAVKQAD
IEKASIMYPSNNCDKIEVIIITLKENKGQRCLNPKSKQARLIKKVERKNF"

BASE COUNT 382 a 170 c 194 g 249 t

ORIGIN

```

1 gaattcggcc aaagaggcct acttccaaga agagcagcaa agctgaagta gcagcaacag
61 caccagcagc aacagcaaaa aacaaacatg agtgtgaagg gcatggctat agccttggct
121 gtgatattgt gtgctacagt tgttcaaggc ttcccatgt tcaaaagagg acgctgtctt
181 tgcataggcc ctggggtaaa agcagtgaag gtggcagata ttgagaaagc ctccataatg
241 tacccaagta acaactgtga caaaatagaa gtgattatta cctgaaaaga aaataaagga
301 caacgatgcc taaatcccaa atcgaagcaa gcaaggctta taatcaaaaa agttgaaaga
361 aagaattttt aaaaatatca aaacatatga agtcctggaa aagggcatct gaaaaaccta
421 gaacaagttt aactgtgact actgaaatga caagaattct acagtaggaa actgagactt
481 ttctatgggt ttgtgacttt caacttttgt acagttatgt gaaggatgaa aggtgggtga
541 aaggaccaaa aacagaaata cagtcttctt gaatgaatga caatcagaat tccactgccc
601 aaaggagtcc aacaattaaa tggatttcta ggaaaagcta ccttaagaaa ggctggttac
661 catcggagtt tacaaagtgc ttacacgttc ttacttggtg tattatacat tcatgcattt
721 ctaggctaga gaaccttcta gatttgatgc ttacaactat tctgttgatg ctatgagAAC
781 atttctgtct ctagaagtta tctgtctgta ttgatcttta tgctatatta ctatctgtgg
841 ttacagtggg gacattgaca ttattactgg agtcaagccc ttataagta aaagcaccta
901 tgtgtcgtaa agcattcctc aaacatttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
961 aaaaaaaaaa aaaaaaaaaa aaaaaaagcg gccgc

```

//

LOCUS AF030514 1371 bp mRNA PRI 17-JUN-1998
DEFINITION Homo sapiens interferon stimulated T-cell alpha chemoattractant precursor, mRNA, complete cds.
ACCESSION AF030514
NID g3219692
KEYWORDS .
SOURCE human.
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (bases 1 to 1371)
AUTHORS Cole, K.E., Strick, C.A., Paradis, T.J., Ogborne, K.T., Loetscher, M., Gladue, R.P., Lin, W., Boyd, J.G., Moser, B., Wood, D.E., Sahagan, B.G., and Neote, K.
TITLE Interferon-inducible T cell alpha chemoattractant (I-TAC): a novel non-ELR CXC chemokine with potent activity on activated T cells through selective high affinity binding to CXCR3
JOURNAL J. Exp. Med. 187 (12), 2009-2021 (1998)
MEDLINE 98290735
REFERENCE 2 (bases 1 to 1371)
AUTHORS Cole, K.E., Strick, C.A. and Sahagan, B.G.
TITLE Direct Submission
JOURNAL Submitted (20-OCT-1997) Molecular Sciences, Pfizer, Inc., Eastern Point Road, Groton, CT 06340, USA
FEATURES
source Location/Qualifiers
1..1371
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="4"
/cell_type="astrocytes"
sig_peptide 70..132
CDS 70..354
/note="chemokine; I-TAC"
/codon_start=1
/product="interferon stimulated T-cell alpha chemoattractant precursor"
/db_xref="PID:g3219693"
/translation="MSVKGMAIALAVILCATVVQGFPMFKRGRCLCIGPGVKAVKQAD
IEKASIMYPSNNCDKIEVIIITLKENKGQRCLNPKSKQARLIKKVERKNF"
mat_peptide 133..351
/evidence=not_experimental

```

                /product="interferon stimulated T-cell alpha
                chemoattractant"
BASE COUNT      487 a      228 c      244 g      411 t      1 others
ORIGIN
    1 ctccttccaa gaagagcagc aaagctgaag tagcagcaac agcaccagca gcaacagcaa
    61 aaaacaaaca tgagtgtgaa gggcatggct atagccttgg ctgtgatatt gtgtgtctaca
   121 gttgttcaag gcttccccat gttcaaaaga ggacgctgtc tttgcatagg ccctggggta
   181 aaagcagtga aagtggcaga tattgagaaa gcctccataa tgtacccaag taacaactgt
   241 gacaaaatag aagtgattat taccctgaaa gaaaataaag gacaacgatg cctaaatccc
   301 aaatcgaagc aagcaaggct tataatcaaa aaagtgtgaaa gaaagaattt ttaaaaatat
   361 caaaacatat gaagtcctgg aaaagggcat ctgaaaaacc tagaacaagt ttaactgtga
   421 ctactgaaat gacaagaatt ctacagtagg aaactgagac ttttctatgg ttttgtgact
   481 ttcaactttt gtacagttat gtgaaggatg aaagtggggt gaaaggacca aaaacagaaa
   541 tacagtcttc ctgaatgaat gacaatcaga attccactgc ccaaaggagt ccagcaatta
   601 aatggatttc taggaaaagc taccttaaga aaggctgggt accatcggag tttacaaagt
   661 gctttcacgt tcttacttgt tgtattatac attcatgcat ttctaggcta gagaaccttc
   721 tagatttgat gcttacaact attctgttgt gactatgaga acatttctgt ctctagaagt
   781 tatctgtctg tattgatctt tatgctatat tactatctgt ggttacagtg gagacattga
   841 cattattact ggagtcgaagc ccttataagt caaaagcatc tatgtgtcgt aaagcattcc
   901 tcaaacattt tttcatgcaa atacacaytt ctttcccaa atatcatgta gcacatcaat
   961 atgtagggaa acattcttat gcatcatttg gtttgtttta taaccaattc attaaatgta
  1021 attcataaaa tgtactatga aaaaaattat acgctatggg atactggcaa cagtgcacat
  1081 atttcataac caaattagca gcaccggctc taatttgatg tttttcaact tttattcatt
  1141 gagatgtttt gaagcaatta ggatatgtgt gtttactgta ctttttggtt tgatccggtt
  1201 gtataaatga tagcaatatc ttggacacat ttgaaataca aaatgtttt gtctaccaa
  1261 gaaaaatggt gaaaaataag caaatgtata ctagcaatc acttttactt tttgtaattc
  1321 tgtctcttag aaaaatacat aatctaata caaaaaaaa aaaaaaaaaa a

```

//

```

LOCUS      AF030514      1371 bp      mRNA      PRI      17-JUN-1998
DEFINITION Homo sapiens interferon stimulated T-cell alpha chemoattractant
            precursor, mRNA, complete cds.
ACCESSION  AF030514
NID        g3219692
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1371)
AUTHORS    Cole,K.E., Strick,C.A., Paradis,T.J., Ogborne,K.T.,
Loetscher,M., Gladue,R.P., Lin,W., Boyd,J.G., Moser,B., Wood,D.E.,
Sahagan,B.G. and Neote,K.
TITLE      Interferon-inducible T cell alpha chemoattractant (I-TAC): a
            novel
            non-ELR CXC chemokine with potent activity on activated T cells
            through selective high affinity binding to CXCR3
JOURNAL     J. Exp. Med. 187 (12), 2009-2021 (1998)
MEDLINE     98290735
REFERENCE  2 (bases 1 to 1371)
AUTHORS     Cole,K.E., Strick,C.A. and Sahagan,B.G.
TITLE       Direct Submission
JOURNAL     Submitted (20-OCT-1997) Molecular Sciences, Pfizer, Inc.,
            Eastern
            Point Road, Groton, CT 06340, USA
FEATURES
            Location/Qualifiers
            source      1..1371
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /chromosome="4"
                        /cell_type="astrocytes"
            sig_peptide 70..132
            CDS          70..354
                        /note="chemokine; I-TAC"
                        /codon_start=1
                        /product="interferon stimulated T-cell alpha
                        chemoattractant precursor"
                        /db_xref="PID:g3219693"

```

```

/translation="MSVKGMAIALAVILCATVVQGFPMFKRGRCLCIGPGVKAVKVAD
IEKASIMYPSNNCDKIEVIITLKENKGQRCLNPKSKQARLIKKVERKNF"
mat_peptide      133..351
                  /evidence=not_experimental
                  /product="interferon stimulated T-cell alpha
                  chemoattractant"
BASE COUNT      487 a    228 c    244 g    411 t    1 others
ORIGIN
1   ctccctccaa gaagagcagc aaagctgaag tagcagcaac agcaccagca gcaacagcaa
61  aaaacaaaca tgagtgtgaa gggcatggct atagccttgg ctgtgatatt gtgtgctaca
121 gttgttcaag gcttccccc atgtcaaaa ggacgctgtc tttgcatagg ccctggggta
181 aaagcagtga aagtggcaga tattgagaaa gcctccataa tgtaccaag taacaactgt
241 gacaaaatag aagtgattat taccctgaaa gaaaataaag gacaacgatg cctaaatccc
301 aaatcgaagc aagcaaggct tataatcaaa aaagtgtgaa gaaagaattt ttaaaaatat
361 caaacatat gaagtcctgg aaaagggcat ctgaaaaacc tagaacaagt ttaactgtga
421 ctactgaaat gacaagaatt ctacagtagg aaactgagac ttttctatgg tttgtgact
481 ttcaactttt gtacagttat gtgaaggatg aaaggtgggt gaaaggacca aaaacagaaa
541 tacagtcttc ctgaatgaat gacaatcaga attccactgc ccaaaggagt ccagcaatta
601 aatggatttc taggaaaagc taccttaaga aaggctgggt accatcggag tttacaaagt
661 gctttcacgt tcttacttgt tgtattatc attcatgcgt ttctaggcta gagaaccttc
721 tagatttgat gcttacaact attctgtgtg gactatgaga acatttctgt ctctagaagt
781 tatctgtctg tattgatctt tatgtatata tactatctgt ggttacagtg gagacattga
841 cattattact ggagtcaagc ccttataagt caaaagcatc tatgtgtcgt aaagcattcc
901 tcaaacattt tttcatgcaa atacacaytt ctttcccaa ataatcatgta gcacatcaat
961 atgtagggaa acattcttat gcatcatttg gtttgtttta taaccaattc attaaatgta
1021 attcataaaa tgtactatga aaaaaattat acgctatggg atactggcaa cagtgcacat
1081 atttcataac caaattagca gcaccggtct taatttgatg tttttcaact tttattcatt
1141 gagatgtttt gaagcaatta ggatatgtgt gtttactgta cttttgtgtt tgatccggtt
1201 gtataaatga tagcaatatc ttggacacat ttgaaataca aaatgttttt gtctaccaa
1261 gaaaaatggt gaaaaataag caaatgtata cctagcaatc acttttactt tttgtaattc
1321 tgtctcttag aaaaatacat aatctaatac aaaaaaaaaa aaaaaaaaaa a

//

```

```

LOCUS      AF030514      1371 bp      mRNA      PRI      17-JUN-1998
DEFINITION Homo sapiens interferon stimulated T-cell alpha chemoattractant
precursor, mRNA, complete cds.
ACCESSION  AF030514
NID        g3219692
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
            Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1371)
AUTHORS    Cole,K.E., Strick,C.A., Paradis,T.J., Ogborne,K.T.,
Loetscher,M., Gladue,R.P., Lin,W., Boyd,J.G., Moser,B., Wood,D.E.,
Sahagan,B.G. and Neote,K.
TITLE      Interferon-inducible T cell alpha chemoattractant (I-TAC): a
novel      non-ELR CXC chemokine with potent activity on activated T cells
            through selective high affinity binding to CXCR3
JOURNAL    J. Exp. Med. 187 (12), 2009-2021 (1998)
MEDLINE    98290735
REFERENCE  2 (bases 1 to 1371)
AUTHORS    Cole,K.E., Strick,C.A. and Sahagan,B.G.
TITLE      Direct Submission
JOURNAL    Submitted (20-OCT-1997) Molecular Sciences, Pfizer, Inc.,
Eastern    Point Road, Groton, CT 06340, USA
FEATURES   Location/Qualifiers
            source      1..1371
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /chromosome="4"
                        /cell_type="astrocytes"
            sig_peptide  70..132
            CDS          70..354
                        /note="chemokine; I-TAC"
                        /codon_start=1

```

/product="interferon stimulated T-cell alpha
chemoattractant precursor"
/db_xref="PID:g3219693"

/translation="MSVKGMAIALAVILCATVVQGFPMFKRGRCLCIGPGVKAVKVAD
IEKASIMYPSNNCDKIEVIITLKENKGQRCLNPKSKQARLIKKVERKNF"
mat_peptide 133..351
/evidence=not_experimental
/product="interferon stimulated T-cell alpha
chemoattractant"

BASE COUNT 487 a 228 c 244 g 411 t 1 others
ORIGIN
1 ctcccttccaa gaagagcagc aaagctgaag tagcagcaac agcaccagca gcaacagcaa
61 aaaacaaaaca tgagtgtgaa gggcatggct atagccttgg ctgtgatatt gtgtgtctaca
121 gttgttcaag gcttcccat gttcaaaaga ggacgtgtgc ttgtcatagg ccctggggta
181 aaagcagtga aagtggcaga tattgagaaa gcctccataa tgtaccaag taacaactgt
241 gacaaaatag aagtgttat taccctgaaa gaaaataaag gacaacgatg cctaaatccc
301 aaatcgaagc aagcaaggct tataatcaaa aaagttgaaa gaaagaattt ttaaaaatat
361 caaacatat gaagtcttgg aaaagggcat ctgaaaaacc tagaacaagt ttaactgtga
421 ctactgaaat gacaagaatt ctacagtagg aaactgagac ttttctatgg tttgtgact
481 ttcaactttt gtacagttaa gtgaaggatg aaagggtgggt gaaaggacca aaaacagaaa
541 tacagtcttc ctgaatgaat gacaatcaga attccactgc ccaaaggagt ccagcaatta
601 aatggatttc taggaaaagc taccttaaga aaggctgggt accatcggag tttacaaagt
661 gctttcacgt tcttacttgt tgtattatac attcatgcat ttctaggcta gagaaccttc
721 tagatttgat gcttacaact attctgttgt gactatgaga acatttctgt ctctagaagt
781 tatctgtctg tattgatctt tatgtctatat tactatctgt ggttacagtg gagacattga
841 cattattact ggagtcaagc cctataagt caaaagcatc tatgtgtcgt aaagcattcc
901 tcaaacattt tttcatgcaa atacacaytt ctttcccaa atatcatgta gcacatcaat
961 atgtagggaa acattcttat gcatcatttg gtttgtttta taaccaattc attaaatgta
1021 attcataaaa tgtactatga aaaaaattat acgctatggg atactggcaa cagtgcacat
1081 atttcataac caaattagca gcaccgtctc taatttgatg tttttcaact ttatttcatt
1141 gagatgtttt gaagcaatta ggatagtgtg gtttactgta cttttgtgtt tgatcgtttt
1201 gtataaatga tagcaatata ttggacacat ttgaaataca aaatgttttt gtctacaaa
1261 gaaaaatgtt gaaaaataag caaatgtata cctagcaatc acttttactt tttgtaattc
1321 tgtctcttag aaaaatacat aatctaatac aaaaaaaaaa aaaaaaaaaa a

//

LOCUS HSMDNCF 1560 bp RNA PRI 31-MAR-1995
DEFINITION Human mRNA for MDNCF (monocyte-derived neutrophil chemotactic factor).
ACCESSION Y00787
NID g34518
KEYWORDS cytokine.
SOURCE human.
ORGANISM Homo sapiens
Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata; Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (bases 1 to 1560)
AUTHORS Matsushima,K.
TITLE Direct Submission
JOURNAL Submitted (03-MAY-1988) Matsushima K., National Cancer Institute,, Bldg 560, Rm 31-19, Frederick, MD 21701
REFERENCE 2 (bases 1 to 1560)
AUTHORS Matsushima,K., Morishita,K., Yoshimura,T., Lavu,S., Kobayashi,Y., Lew,W., Appella,E., Kung,H.F., Leonard,E.J. and Oppenheim,J.J.
TITLE Molecular cloning of a human monocyte-derived neutrophil chemotactic factor (MDNCF) and the induction of MDNCF mRNA by interleukin 1 and tumor necrosis factor
JOURNAL J. Exp. Med. 167 (6), 1883-1893 (1988)
MEDLINE 88258376
COMMENT for overlapping sequence see M17016 - M17017.
FEATURES
Location/Qualifiers
source 1..1560
/organism="Homo sapiens"
/db_xref="taxon:9606"
/cell_type="monocyte"
/clone_lib="lambda gt10"
sig_peptide 102..182
/note="signal peptide (AA -27 to -1)"

```

CDS                102..401
                   /codon_start=1
                   /product="MDNCF precursor (AA -27 to 72)"
                   /db_xref="PID:g34519"
                   /db_xref="SWISS-PROT:P10145"

/translation="MTSKLAVALLAAFLISAALCEGAVLPRSAKELRCQCIKTYSKPF"

HPKFIKELRVIESGPHCANTEIIVKLSDGRELCLDPKENWVQRVVEKFLKRAENS"
mat_peptide        183..398
                   /note="mat. MDNCF (AA 1 - 72)"
BASE COUNT         526 a    247 c    281 g    506 t
ORIGIN
    1 ctccataagg cacaaacttt cagagacagc agagcacaca agcttctagg acaagagcca
   61 ggaagaaacc accggaagga accatctcac tgtgtgtaaa catgacttcc aagctggccg
  121 tggctctctt ggcagccttc ctgatttctg cagctctgtg tgaaggtgca gttttgccaa
  181 ggagtgtctaa agaacttaga tgtcagtgca taaagacata ctccaaacct ttccacccca
  241 aatttatcaa agaactgaga gtgattgaga gtggaccaca ctgcgccaac acagaaatta
  301 ttgtaaagct ttctgatgga agagagctct gtctggaccc caaggaaaaac tgggtgcaga
  361 gggttgtgga gaagtttttg aagagggctg agaattcata aaaaaattca ttctctgtgg
  421 tatccaagaa tcagtgaaga tgccagttaa acttcaagca aatctacttc aacacttcat
  481 gtattgtgtg ggtctgttgt agggttgcca gatgcaatac aagattcctg gttaaatttg
  541 aatttcagta aacaatgaat agtttttcat tgtaccatga aatatccaga acatacttat
  601 atgtaaagta ttattttatt gaatctacaa aaaacaacaa ataattttta aatataagga
  661 ttttcctaga tattgcacgg gagaatatac aaatagcaaa attggggcaa gggccaagag
  721 aatatccgaa ctttaatttc aggaattgaa tgggtttgct agaattgtat atttgaagca
  781 tcacataaaa atgatgggac aataaatttt gccataaagt caaatttagc tggaaatcct
  841 ggattttttt ctgttaaatc tggcaaccct agtctgctag ccaggatcca caagtccttg
  901 ttccactgtg ccttggtttc tcctttattt ctaagtggaa aaagtattag ccaccatctt
  961 acctcacagt gatgttgtga ggacatgtgg aagcacttta agttttttca tcataacata
 1021 aattattttc aagtgttaact tattaacccta tttattattt atgtatttat ttaagcatca
 1081 aatattttgt caagaatttg gaaaaataga agatgaatca ttgattgaat agttataaag
 1141 atgttatagt aaattttatt tattttagat attaaatgat gttttattag ataaatttca
 1201 atcagggttt ttagattaaa caaacaacaa attgggtacc cagttaaatt ttcatttcag
 1261 atatacaaca aataattttt tagtataagt acattattgt ttatctgaaa ttttaattga
 1321 actaacaatc ctagtttgat actcccagtc ttgtcattgc cagctgtgtt ggtagtgtg
 1381 tgttgaatta cggaataatg agttgaactc attaaaacag ccaaaaactcc acagtcaata
 1441 ttagtaattt cttgctggtt gaaacttgtt tattatgtac aaatagattc ttataatatt
 1501 atttaaataa ctgcattttt aaatacaagg ctttatattt ttaactttta aaaaaaccgg

//

LOCUS              HSINFGFR      1172 bp      RNA              PRI              21-MAR-1995
DEFINITION         Human mRNA for gamma-interferon inducible early response gene
(with
homology to platelet proteins).
ACCESSION          X02530 M17752
NID                g33917
KEYWORDS            interferon response; signal peptide.
SOURCE             human.
ORGANISM            Homo sapiens
                    Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
                    Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE           1 (bases 1 to 1172)
AUTHORS            Luster,A.D., Unkeless,J.C. and Ravetch,J.V.
TITLE              Gamma-interferon transcriptionally regulates an early-response
gene
                    containing homology to platelet proteins
JOURNAL            Nature 315 (6021), 672-676 (1985)
MEDLINE            85240552
REFERENCE           2 (bases 1 to 1172)
AUTHORS            Luster,A.D.
TITLE              Direct Submission
JOURNAL            Submitted (29-JUL-1986) to the EMBL/GenBank/DDBJ databases
COMMENT            Data kindly reviewed (29-JUL-1986) by Luster A.D.
FEATURES
    source          Location/Qualifiers
                    1..1172
                    /organism="Homo sapiens"
                    /strain="(U 937 histiocytic lymphoma cell line)"
                    /db_xref="taxon:9606"
    misc_RNA        1
                    /note="cap site"

```

sig_peptide 67..129
 /note="pot. signal peptide (aa-21 to -1)"
 CDS 67..363
 /note="early response precursor polypeptide (aa-21 to 77)"
 /codon_start=1
 /db_xref="PID:g33918"
 /db_xref="SWISS-PROT:P02778"

/translation="MNQTAILICCLIFLTLSGIQGVPLSRTVRCTCISISNQPVNPRS

LEKLEIIPASQFCPRVEIIATMKKKGEKRCINPESKAIKNLLKAVSKEMSKRSP"

mat_peptide 130..360
 /note="mature early response polypeptide (aa 1-77)"
 old_sequence 1138..1141
 /note="ugaa was uga in [1]"
 /citation=[1]
 old_sequence 1146..1148
 /note="caa was ca in [1]"
 /citation=[1]
 misc_feature 1155..1160
 /note="pot. polyA signal"
 polyA_site 1172
 /note="polyA site"

BASE COUNT 384 a 231 c 208 g 349 t
 ORIGIN

```

1 gagacattcc tcaattgctt agacatatcc tgagcctaca gcagaggaac ctccagtctc
61 agcaccatga atcaaaactgc gattcttgatt tgctgcctta tctttctgac tctaagtggc
121 attcaaggag tacctctctc tagaaccgta cgctgtacct gcatacagcat tagtaatcaa
181 cctgttaatc caaggtcttt agaaaaactt gaaattattc ctgcaagcca attttgtcca
241 cgtgttgaga tcattgctac aatgaaaaag aagggtgaga agagatgtct gaatccagaa
301 tcgaaggcca tcaagaattt actgaaagca gttagcaagg aaatgtctaa aagatctcct
361 taaaaccaga ggggagcaaa atcgatgcag tgcttccaag gatggaccac acagaggctg
421 cctctcccat cacttcccta catggagtat atgtcaagcc ataattgttc ttagtttgca
481 gttacactaa aaggtgacca atgatgggtca ccaaactcagc tgctactact cctgtaggaa
541 gggttaattt catcatccta agctattcag taataactct accctggcac tataatgtaa
601 gctctactga ggtgctatgt tcttagtgga tgttctgacc ctgcttcaaa tatttccttc
661 acctttccca tcttccaagg gtactaagga atctttctgc tttgggggtt atcagaattc
721 tcagaatctc aaataactaa aaggatatgca atcaaatctg ctttttaaag aatgctcttt
781 acttcatgga ctccactgc catcttccca aggggcccac attctttcag tggctacctc
841 catacaattc caaacacata caggaaggta gaaatatctg aaaatgtatg tgtaagtatt
901 cttattttaat gaaagactgt acaaaagtata agtcttagat gtatatattt cctatatgtt
961 tttcagtgta catggaataa catgtaatta agtactatgt atcaatgagt aacaggaaaa
1021 ttttaaaaat acagatagat atatgctctg catgtttacat aagataaatg tgctgaatgg
1081 ttttcaataa aaaatgaggt actctcctgg aaatattaag aaagactatc taaatgttga
1141 aagatcaaaa ggttaataaa gtaattataa ct

```

//

LOCUS SYNRP4A 225 bp DNA SYN 15-JUN-1989
 DEFINITION Human recombinant platelet factor 4 (PF4) gene, complete cds.
 ACCESSION M20901
 NID g209285
 KEYWORDS platelet factor; platelet factor 4.
 SOURCE Synthetic oligonucleotide DNA, clone pIN-III-ompA-2.
 ORGANISM artificial sequence
 artificial sequence.
 REFERENCE 1 (bases 1 to 225)
 AUTHORS Barone,A.D., Ghrayeb,J., Hammerling,U., Zucker,M.B. and Thorbecke,G.J.
 TITLE The expression in Escherichia coli of recombinant human platelet factor 4, a protein with immunoregulatory activity
 JOURNAL J. Biol. Chem. 263, 8710-8715 (1988)
 MEDLINE 88243725
 FEATURES
 source Location/Qualifiers
 1..225
 /organism="artificial sequence"
 /db_xref="taxon:29278"
 CDS <1..>225
 /note="recombinant platelet factor 4"
 /codon_start=2

```

/transl_table=11
/db_xref="PID:g209286"

/translation="ASMEAEDGDLQCLCVKTTTSQVRPRHITSLEVIKAGPHCPTAQL
IATLKDGRKICLDLQAPLYKKI IKKLLSEGS"
BASE COUNT      59 a      59 c      51 g      56 t
ORIGIN          HindIII site.
      1 agcttctatg gaagctgaag aagacgggtga cctgcagtgc ctgtgcgtta aaactacttc
     61 tcagggttcgt ccgcgtcata tcacttctct ggaagtatc aaagctggtc cgcattgccc
    121 gactgctcag ctgatcgcta ctctcaaaga cggtcgtaaa atctgcctgg acctgcaggc
    181 tccgctgtac aaaaaaatca tcaaaaaact gctggaatct ggatc
//

LOCUS          HUMGRO          1050 bp      mRNA          PRI          11-JUN-1993
DEFINITION      Human gro (growth regulated) gene.
ACCESSION       J03561
NID             g183622
KEYWORDS        gro gene; tumor cell.
SOURCE          Human bladder tumor cell (T24) cDNA to mRNA.
  ORGANISM      Homo sapiens
    Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
    Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE       1 (bases 1 to 1050)
AUTHORS         Anisowicz,A., Bardwell,L. and Sager,R.
TITLE           Constitutive overexpression of a growth-regulated gene in
                transformed Chinese hamster and human cells
JOURNAL         Proc. Natl. Acad. Sci. U.S.A. 84, 7188-7192 (1987)
MEDLINE         88041072
COMMENT         Draft entry and computer-readable sequence kindly submitted by
                R.Sager (20-NOV-1987).
FEATURES        Location/Qualifiers
  source         1..1050
                /organism="Homo sapiens"
                /db_xref="taxon:9606"
  sig_peptide    54..140
                /note="signal peptide (put.); putative"
  CDS            54..377
                /note="gro protein"
                /codon_start=1
                /db_xref="PID:g306806"

/translation="MARAALSAAPSNPRLLRVALLLLLLVAAGRRAAGASVATELRCQ
CLQTLQGIHPKNIQSVNVKSPGPHCAQTEVIATLKNGRKACLNPPASPIVKKIIEKMLN
SDKSN"
  mat_peptide    141..374
                /note="gro mature protein (put.); putative"
BASE COUNT      270 a      246 c      239 g      295 t
ORIGIN          52 bp upstream of NcoI site.
      1 ctgcgccagt cttcgcgtcc tctcacagcc gccagaccgc cctgctgagc cccatggccc
     61 gcgctgctct ctcgcgcgcc ccagcaatc cccggctcct gcgagtggca ctgctgctcc
    121 tgctcctggt agccgctggc cggcgcgcag caggagcgtc cgtggccact gaactgcgct
    181 gccagtgcct gcagaccctg cagggaattc accccaagaa catccaaagt gtgaacgtga
    241 agtcccccg accccactgc gcccaaacgc aagtcatagc cactcaag aatgggcgga
    301 aagcttgcc caatcctgca tcccccatag ttaagaaaat catcgaaaag atgctgaaca
    361 gtgacaaatc caactgacca gaagggagga ggaagctcac tggtagctgt tctgaagga
    421 ggccctgccc ttataggaac agaagaggaa agagagacac agctgcagag gccacctgga
    481 ttgtgcctaa tgtgtttgag catcgcttag gagaagtctt ctatttattt atttattcat
    541 tagttttgaa gattctatgt taatatatta ggtgtaaaat aattaagggt atgattaact
    601 ctacctgcac actgtcctat tatattcatt ctttttgaaa tgtcaacccc aagttagttc
    661 aatctggatt catatttaat ttgaaggtag aatgttttca aatgttctcc agtcattatg
    721 ttaatatattc tgaggagcct gcaacatgcc agccactgtg atagaggctg gcggatccaa
    781 gcaaatggcc aatgagatca ttgtgaagcc aggggaatgt atgtgcacat ctgttttgta
    841 actgttttaga tgaatgtcag ttgttattta ttgaaatgat ttcacagtgt gtggtcaaca
    901 tttctcatgt tgaaacttta agaactaaaa tgttctaaat atcccttgga cattttatgt
    961 ctttcttgta aggcatactg ccttggttaa tggtagtttt acagtgtttc tggcttagaa
   1021 caaaggggct taattattga tgttttcgga
//

LOCUS          HUMGROB5        1110 bp      mRNA          PRI          07-MAR-1995
DEFINITION      Human cytokine (GRO-beta) mRNA, complete cds.

```

ACCESSION M36820
 NID g183628
 KEYWORDS cytokine.
 SOURCE Human lymphocyte, cDNA to mRNA, clone GRO-beta.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 1110)
 AUTHORS Haskill, S., Peace, A., Morris, J., Sporn, S.A., Anisowicz, A.,
 Lee, S.W., Smith, T., Martin, G., Ralph, P. and Sager, R.
 TITLE Identification of three related human GRO genes encoding
 cytokine
 functions
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 87 (19), 7732-7736 (1990)
 MEDLINE 91017578
 COMMENT Draft entry and computer-readable sequence for [Proc. Natl.
 Acad. Sci. U.S.A. (1990) In press] kindly submitted
 by S.Haskill, 20-JUL-1990.

FEATURES Location/Qualifiers
 source 1..1110
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /clone="GRO-beta"
 /tissue_type="monocyte and lymphocyte"
 gene 75..398
 /gene="GRO-beta"
 CDS 75..398
 /gene="GRO-beta"
 /codon_start=1
 /product="cytokine gro-beta"
 /db_xref="PID:g183629"

/translation="MARATLSAAPSNPRLRLVALLLLLLVAASRRAGAPLATELRQC

CLQTLOGIHLKNIQSVKVKSPGPHCAQTEVIATLKNGQKACLNPA SPMVKKIIEKMLK
 NGKSN"

BASE COUNT 300 a 247 c 247 g 316 t
 ORIGIN

```

1  gacagagccc gggccacgga gtcctctgcc agctctcctc ctgcacagc cgctcgaacc
61  gcctgctgag ccccatggcc cgcgccacgc tctccgcgcg cccagcaat cccgggctcc
121  tgcgggtggc gctgctgctc ctgctcctgg tggccgccag ccggcgcgca gcaggagcgc
181  ccctggccac tgaactgcgc tgccagtgc tgcagaccct gcagggaatt cacctcaaga
241  acatccaaag tgtgaagggt aagtcctccg gacccactg cgcccaaacc gaagtcatag
301  ccacactcaa gaatgggcag aaagcttgct tcaacccgcg atcgcccatg gttaaagaaa
361  tcactgaaaa gatgctgaaa aatggcaaat ccaactgacc agaaggaagg aggaagctta
421  ttggtggctg ttcctgaagg aggccttgcc ttacaggaac agaagaggaa agagagacac
481  agctgcagag gccacctggc ttgcgcctaa tgtgtttgag catacttagg agaagtcctc
541  tatttattta tttatttatt tattgtttg ttttagaaga ttctatgta atattttatg
601  tgtaaaaataa ggttatgatt gaactactt gcacactctc ccattatatt tattgtttat
661  tttagggtcaa acccaagtta gttcaatcct gattcatatt taatttgaag atagaagggtt
721  tgcagatatt ctctagtcac ttgttaatat ttcttcgtga tgacatatca catgtcagcc
781  actgtgatag aggctgagga atccaagaaa atggccagta agatcaatgt gacggcaggg
841  aaatgtatgt gtgtctatgt tgtaactgta aagatgaatg tcagttgta tttattgaaa
901  tgatttcaca gtgtgtgggc aacatttctc atgttgaagc tttagaact aaaatgttct
961  aaatatccct tggcatttta tgtcttctt gtaagatact gccttggtta atgttaatta
1021  tgcagtgttt cctctgtgt tagagcagag aggtttcgat atttattgat gttttcacia
1081  agaacaggaa aataaaatat ttaaaatat
  
```

//

LOCUS HUMGROG5 1064 bp mRNA PRI 07-MAR-1995
 DEFINITION Human cytokine (GRO-gamma) mRNA, complete cds.
 ACCESSION M36821
 NID g183632
 KEYWORDS cytokine.
 SOURCE Human lymphocyte, cDNA to mRNA, clone GRO-gamma.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 1064)
 AUTHORS Haskill, S., Peace, A., Morris, J., Sporn, S.A., Anisowicz, A.,

Lee, S.W., Smith, T., Martin, G., Ralph, P. and Sager, R.
 TITLE Identification of three related human GRO genes encoding
 cytokine functions
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 87 (19), 7732-7736 (1990)
 MEDLINE 91017578
 COMMENT Draft entry and computer-readable sequence for [Proc. Natl.
 Acad. Sci. U.S.A. (1990) In press] kindly submitted
 by S.Haskill, 20-JUL-1990.
 FEATURES Location/Qualifiers
 source 1..1064
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /clone="GRO-gamma"
 /tissue_type="lymphocyte and monocyte"
 gene 78..398
 /gene="GRO-gamma"
 CDS 78..398
 /gene="GRO-gamma"
 /codon_start=1
 /product="cytokine GRO-gamma"
 /db_xref="PID:g183633"

/translation="MAHATLSAAPSNPRLLRVALLLLLLVGSRRRAAGASVVTELRCQC

LQTLQGIHLKNIQSVNVRSPGPHCAQTEVIATLKNKGKACLNPASPMVQKIIIEKILNK
 GSTN"

| BASE COUNT | 281 a | 237 c | 239 g | 305 t | 2 others |
|------------|-------------|------------|-------------|------------|------------------------|
| ORIGIN | | | | | |
| 1 | cacagccggg | tgcaggcac | ctccccngcc | agctctcccg | cattctgcac agcttcccga |
| 61 | cgctctctgct | gagccccatg | gcccacgcca | cgctctccgc | cgccccagc aatccccggc |
| 121 | tcctgcccgt | ggcgctgctg | ctcctgctcc | tggtgggcag | ccggcgcgca gcaggagcgt |
| 181 | ccgtgggtcac | tgaactgcgc | tgccagtgcct | tcagacact | gcagggaatt cacctcaaga |
| 241 | acatccaaag | tgtgaatgta | aggtcccccg | gacccactg | cgcccaaacc gaagtcatag |
| 301 | ccacactcaa | gaatgggaag | aaagcttgtc | tcaacccgc | atcccccatg gtccagaaaa |
| 361 | tcacgaaaa | gatactgaac | aaggggagca | ccaactgaca | ggagagaagt aagaagctta |
| 421 | tcagcgatc | attgacactt | cctgcagggt | ggccctgcc | ctaccagag ctgaaaatga |
| 481 | aaaagagaac | agcagcttcc | tagggacagc | tggaaggga | ctaatgtgt ttgactatct |
| 541 | cttacgagg | ttctacttat | ttatgtatct | atcttgaaa | gcttgatctt taatatttta |
| 601 | catgctgtta | tttaaagatg | tgagtgtgtt | tcacaaaca | tagctcagtc ctgattatct |
| 661 | aattggaata | tgatgggttt | taaatgtgtc | attaaactaa | tatttagtgg gagaccataa |
| 721 | tgtgtcagcc | accttgataa | atgacagggt | ggggaactgg | agggtngggg gattgaaatg |
| 781 | caagcaatta | gtggatcact | gttagggtaa | gggaatgtat | gtacacatct attttttata |
| 841 | cttttttttt | taaaaaagaa | tgtcagttgt | tatttattca | aattatctca cattatgtgt |
| 901 | tcaacatttt | tatgctgaag | tttcccttag | acattttatg | tcttgcttgt agggcataat |
| 961 | gccttgttta | atgtccattc | tgacagcttt | ctcttccct | tggaagaagag aatttatcat |
| 1021 | tactgttaca | tttgtacaaa | tgacatgata | ataaaagttt | tatg |

//

LOCUS HUMCTAP3 673 bp mRNA PRI 06-MAR-1995
 DEFINITION Human connective tissue activation peptide III mRNA, complete
 cds.
 ACCESSION M54995 M38441
 NID g181175
 KEYWORDS connective tissue activating peptide-III; platelet basic
 protein;
 thromboglobulin.
 SOURCE Human platelet, cDNA to mRNA, clone lambda-c[1,2].
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 673)
 AUTHORS Wenger, R.H., Wicki, A.N., Walz, A., Kieffer, N. and Clemetson, K.J.
 TITLE Cloning of cDNA coding for connective tissue activating peptide
 III
 from a human platelet-derived lambda gt11 expression library
 JOURNAL Blood 73 (6), 1498-1503 (1989)
 MEDLINE 89229374
 FEATURES Location/Qualifiers
 source 1..673

```

/organism="Homo sapiens"
/db_xref="taxon:9606"
/tissue_type="platelet"
/clone="lambda-cl"
/cell_type="platelet"
/tissue_type="blood"
/tissue_lib="lambda-gt11"
/map="4p13-q21"
gene      67..453
          /gene="PPBP"
sig_peptide 67..168
          /gene="PPBP"
          /note="G00-127-391"
CDS       67..453
          /gene="PPBP"
          /codon_start=1
          /db_xref="GDB:G00-127-391"
          /product="connective tissue activating peptide III"
          /db_xref="PID:g181176"

/translation="MSLRLDTPSCNSARPLHALQVLLLLLLLTALASSTKGQTKRN

LAKGKEESLSDLYAELRCMCIKTTSGIHPKNIQSLEVIGKGTGHCNQVEVIATLKDGR
KICLDPDAPRIKKIVQKKLAGDESAD"
mat_peptide 196..450
          /gene="PPBP"
          /note="G00-127-391"
          /product="connective tissue activating peptide III"
mat_peptide 208..450
          /gene="PPBP"
          /note="G00-127-391"
          /product="beta-thromboglobulin"
polyA_site 673
          /gene="PPBP"
          /note="G00-127-391"
BASE COUNT      202 a      149 c      139 g      183 t
ORIGIN
    1 gggcaactca ccctcactca gaggtcttct ggttctggaa acaactctag ctcagccttc
   61 tccaccatga gcctcagact tgataccacc ccttcctgta acagtgcgag accacttcat
  121 gccttgcagg tgctgctgct tctgtcattg ctgctgactg ctctggcttc ctcaccacaa
  181 ggacaaacta agagaaactt ggcgaaaggc aaagaggaaa gtctagacag tgacttgtat
  241 gctgaactcc gctgcatgtg tataaagaca acctctggaa ttcattccaa aaacatccaa
  301 agtttggaag tgatcgggaa aggaacccat tgcaaccaag tcgaagtgat agccacactg
  361 aaggatggga ggaaaatctg cctggaccga gatgctccca gaatcaagaa aattgtacag
  421 aaaaaattgg caggtgatga atctgctgat taattgttc tgtttctgcc aaacttcttt
  481 aactcccagg aagggtagaa ttttgaaacc ttgattttct agagttctca tttattcagg
  541 atacctattc ttactgtatt aaaatttgga tatgtgtttc attctgtctc aaaaatcaca
  601 ttttattctg agaaggttgg ttaaaagatg gcagaaagaa gatgaaaata aataagcctg
  661 gtttcaaccc tct

//

LOCUS      HUMENA78A      2177 bp      DNA      PRI      31-JAN-1996
DEFINITION Homo sapiens neutrophil-activating peptide 78 (ENA-78) gene,
            complete cds.
ACCESSION  L37036 246254
NID        g607030
KEYWORDS   ENA-78 gene; homologue; neutrophil-activating factor;
            neutrophil-activating peptide 78.
SOURCE     Homo sapiens DNA.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 2177)
AUTHORS    Walz,A., Burgener,R., Car,B., Baggiolini,M., Kunkel,S.L. and
            Strieter,R.M.
TITLE      Structure and neutrophil-activating properties of a novel
            inflammatory peptide (ENA-78) with homology to interleukin 8
JOURNAL     J. Exp. Med. 174 (6), 1355-1362 (1991)
MEDLINE     92078844
REFERENCE  2 (bases 1 to 2177)
AUTHORS     Walz,A.

```

TITLE Direct Submission
 JOURNAL Submitted (14-OCT-1994) A. Walz, University of Bern, Theodor
 Kocher
 Institute, Freiestr. 1, Bern, Switzerland 3012
 REFERENCE 3 (bases 1 to 2177)
 AUTHORS Corbett,M.S., Schmitt,I., Riess,O. and Walz,A.
 TITLE Characterization of the gene for human neutrophil-activating
 peptide 78 (ENA-78)
 JOURNAL Biochem. Biophys. Res. Commun. 205 (1), 612-617 (1994)
 MEDLINE 95091791
 FEATURES
 source Location/Qualifiers
 1..2177
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /cell_type="lymphoblastoid cells"
 /clone="4H2, 178C11, 106C1"
 /chromosome="4"
 /clone_lib="Chromosome 4 cosmid library of Riess et
 al."
 gene 539..1747
 /gene="ENA-78"
 CAAT_signal 539..547
 /gene="ENA-78"
 TATA_signal 675..681
 /gene="ENA-78"
 exon <803..911
 /gene="ENA-78"
 /number=1
 CDS join(803..911,1046..1178,1289..1372,1729..1747)
 /gene="ENA-78"
 /note="homologue to interleukin-8"
 /codon_start=1
 /product="neutrophil-activating peptide 78"
 /db_xref="PID:g607031"
 /translation="MSLLSSRAARVPGPSSSLCALLVLLLLLTQPGPIASAGPAAAVL
 RELRCVCLQTTQGVHPKMISNLQVFAIGPQCSKVEVVASLKNGKEICLDPEAPFLKKV
 IQKILDGGNKEN"
 intron 912..1045
 /gene="ENA-78"
 /number=1
 exon 1046..1178
 /gene="ENA-78"
 /number=2
 intron 1179..1288
 /gene="ENA-78"
 /number=2
 exon 1289..1372
 /gene="ENA-78"
 /number=3
 intron 1373..1728
 /gene="ENA-78"
 /number=3
 exon 1729..>1747
 /gene="ENA-78"
 /number=4
 BASE COUNT 539 a 512 c 496 g 630 t
 ORIGIN
 1 gaattctcag taagcggact taccaaagta ggtgatctgt aggggagtta acaaaattca
 61 gtggctcctt caggccactg acttcaagtg gcaagagaca aggggtctctt gttatcatgt
 121 tatcttggct tccaaagctg gttgaagtcc agagattcat aaagtcattc aagaaacctc
 181 gaatgacctg cctgcaagaa gacaggaagg actttcagtt tatagcaatt caaacatgaa
 241 taacatttcc tgattaatag taataataat tagaaaggat tgactttcag aaatttttct
 301 caaatcaagg ctctgtttac tttggttcca ccttttctct ctagaaggag aggaggagca
 361 tctcccagat gctgcgtgct ccagaaaagc cggcatccct agcccgtctt ggcacaggcc
 421 atgaggcgct gctgaatcct gctgaatagc tactcccttc tagctggagc cacagctccc
 481 tccaccgcgg aacagggtta caacgtccct ctggtagag gtgcacgcag ctctctctgg
 541 ccaccctccc caccagttcc cattgtcttg cccccctccc ccaacctctt ctttccacac
 601 tgcccatga gttcaggga tttcccagc atcccaaagc ttgagtttcc tgtcagtggg
 661 gagagatgag tgtagataaa aggagtgcag aaggaacgag gaagccacag tgctccgag

```

721 cctccaatct tgcctcctcc aatctccgct cctccacca gttcaggaac ccgcgaccgc
781 tcgcagcgct ctcttgacca ctatgagcct cctgtccagc cgcgcgcccc gtgtccccgg
841 tccttcgagc tccttgtgcg cgctgttggt gctgtgctg ctgctgacgc agccagggcc
901 catcgccagc ggtgagagcg catggcgcg cgacgcact cgcactcggg cacagaggtg
961 catcccagcc tctgcggggt cgctgcgttc cagggaaact tcccagcaac ctgccctata
1021 aagggtgtct ctctttcttc cccagctggt cctgccgctg ctgtgttgag agagctgcgt
1081 tgcgtttggt tacagaccac gcaaggagtt catcccaaaa tgatcagtaa tctgcaagtg
1141 ttcgccatag gccacagtg ctccaaggtg gaagtgtgtg aagtctctgt ctgctgtgtc
1201 cgctgtgacc ttggcaagag agaaatcccg cagcctgggt cttcaacctt ggtatctcat
1261 gagtgtatct tctttttctt tccttcagag cctccctgaa gaacgggaag gaaatttgtc
1321 ttgatccaga agcccccttt ctaaagaaag tcatccagaa aattttggac gggacttgt
1381 cactttgatc tttgtggttt ctaaatctga tctagggaga ccatagactt cacaaggtct
1441 ttattctctg tacgatttaa gtaacacttt tcatgtttag aattaaaagg ttgttgaatt
1501 gggaaagtgt ttttggtattg tcctgggaaa atataccaat cttacatgta attacttgag
1561 caattacaca cagcttgta ctaagttatg tttttgttt acccattgct tttattgatt
1621 tttgtattct ccttttttac caaacatcat aaacgctgag ttttgacaag ggtggagtag
1681 aaaggagtgt gaaaaatggt taaactaata taacattttt ctcaacagtg gaaacaagga
1741 aaactgatta agagaaatga gcacgcattg aaaagtttcc cagtcttcag cagagaagtt
1801 ttttgagggt ctctgaaccc aggggaagaca agaaggaaag attttgttgt tgtttgttta
1861 tttgtttttc cagtagttag cttttctctt ggattcctca ctttgaagag tgtgaggaaa
1921 acctatgttt gccgcttaag ctttcagctc agctaagaa gtgttttagca tagtacctct
1981 gctatttgcg gttattttat ctgctatgct attgaagttt tggcaattga ctatagtgtg
2041 agccaggaat cactggctgt taatctttca aagtgtcttg aattgtaggt gactattata
2101 tttccaagaa atattcctta agatattaac tgagaaggct gtggatttaa tgtggaaatg
2161 atgtttcata agaatttc

```

//

```

LOCUS       HSGCP2             254 bp      RNA          PRI          04-MAR-1997
DEFINITION  H.sapiens mRNA for granulocyte chemotactic protein.
ACCESSION   Y08770
NID         g1769436
KEYWORDS    cell surface receptor; CXC chemokine; GCP-2 gene; granulocyte
            chemotactic protein.
SOURCE      human.
ORGANISM    Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1 (bases 1 to 254)
AUTHORS     Froyen,G., Proost,P., Ronsse,I., Mitera,T., Haelens,A.,
            Wuyts,A.,
            Opdenakker,G., Van Damme,J. and Billiau,A.
TITLE       Cloning, bacterial expression and biological characterization
            of
            recombinant human granulocyte chemotactic protein-2 and
            differential expression of granulocyte chemotactic protein-2
            and
            epithelial cell-derived neutrophil activating peptide-78 mRNAs
JOURNAL     Eur. J. Biochem. 243 (3), 762-769 (1997)
MEDLINE     97210779
REFERENCE   2 (bases 1 to 254)
AUTHORS     Froyen,G.F.V.
TITLE       Direct Submission
JOURNAL     Submitted (10-OCT-1996) G.F.V. Froyen, Rega Institute,
            University
            of Leuven, Minderbroedersstraat 10, B-3000 Leuven, BELGIUM
FEATURES    Location/Qualifiers
            source          1..254
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /haplotype="diploid"
                        /tissue_type="embryonic"
                        /rearranged
                        /cell_type="fibroblast"
                        /cell_line="E6SM (embryonic strain - skin and muscle)"
            gene            1..254
                        /gene="GCP-2"
            exon            <1..131
                        /gene="GCP-2"
                        /number=2
            CDS              <1..234
                        /gene="GCP-2"

```

```

/codon_start=1
/product="granulocyte chemotactic protein"
/db_xref="PID:e283124"
/db_xref="PID:g1769437"

/translation="GPVSAVLTELRCTCLRVTLRVNPKTIGKLQVFPAGPQCSKVEVV
              ASLKNGKQVCLDPEAPFLKKVIQKILDSGNKKN"
      exon      132..215
                /gene="GCP-2"
                /number=3
      exon      216..254
                /gene="GCP-2"
                /number=4
      3'UTR      235..254
                /gene="GCP-2"
BASE COUNT      66 a      64 c      70 g      54 t
ORIGIN
      1 ggtcctgtct ctgctgtgct cacggagctg cgttgcactt gtttacgcgt tacgctgaga
     61 gtaaacccca aaacgattgg taaactgcag gtgttccccg caggcccgcg gtgctccaag
    121 gtggaagtgg tagcctccct gaagaacggg aagcaagttt gtctggaccc ggaagcccct
    181 tttctaaaga aagtcatcca gaaaattttg gacagtggaa acaagaaaaa ctgagtaaca
    241 gtcgacgcgg ccgc
//

LOCUS      D63789      5669 bp      DNA      PRI      27-DEC-1996
DEFINITION Human DNA for SCM-1beta precursor, complete cds.
ACCESSION  D63789
NID        g1754608
KEYWORDS   SCM-1beta; SCM-1beta precursor.
SOURCE      Homo sapiens placenta DNA, clone:hg44.
ORGANISM    Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE  1 (sites)
AUTHORS    Yoshida,T., Imai,T., Kakizaki,M., Nishimura,M. and Yoshie,O.
TITLE      Molecular cloning of a novel C or gamma type chemokine, SCM-1
JOURNAL    FEBS Lett. 360 (2), 155-159 (1995)
MEDLINE    95180438
REFERENCE  2 (sites)
AUTHORS    Yoshida,T., Imai,T., Takagi,S., Nishimura,M., Ishikawa,I.,
Yaoi,T.
            and Yoshie,O.
TITLE      Structure and expression of two highly related genes encoding
            SCM-1/human lymphotactin
JOURNAL    FEBS Lett. 395 (1), 82-88 (1996)
MEDLINE    97002294
REFERENCE  3 (bases 1 to 5669)
AUTHORS    Yoshida,T.
JOURNAL    Unpublished (1995)
REFERENCE  4 (bases 1 to 5669)
AUTHORS    Yoshida,T.
TITLE      Direct Submission
JOURNAL    Submitted (07-AUG-1995) to the DDBJ/EMBL/GenBank databases.
Tetsuya
            Yoshida, Shionogi Institute for Medical Science; 2-5-1,
Mishima,
            Settsu, Osaka 566, Japan (E-
            mail:teyoshid@fl.lab.shionogi.co.jp,
            Tel:06-382-2612, Fax:06-382-2598)
FEATURES
      source      Location/Qualifiers
                  1..5669
                  /organism="Homo sapiens"
                  /db_xref="taxon:9606"
                  /chromosome="1"
                  /clone="hg44"
                  /map="1q23"
                  /tissue_type="placenta"
      TATA_signal  2154..2158
      exon        2197..2278

```

```

        /number=1
prim_transcript 2197..5349
gene            2218..5230
               /gene="SCM-1beta"
CDS             join(2218..2278,4075..4189,5062..5230)
               /gene="SCM-1beta"
               /codon_start=1
               /product="SCM-1beta precursor"
               /db_xref="PID:d1010504"
               /db_xref="PID:g1754609"

/translation="MRLILALLGICSLTAYIVEGVGSEVSHRRTCVSLLTQRLPVS
IKTYTITEGSLRAVIFITKRLKVCADPQATWVRDVVRSMRKSNTNRNMIQTKPTGT
               QQSTNTAVTLTG"
intron          2279..4074
               /gene="SCM-1beta"
               /number=1
exon            4075..4189
               /gene="SCM-1beta"
               /number=2
mat_peptide     join(4077..4189,5062..5227)
               /gene="SCM-1beta"
intron          4190..5061
               /gene="SCM-1beta"
               /number=2
exon            5062..5349
               /number=3
BASE COUNT     1702 a   1058 c   1248 g   1661 t
ORIGIN
1  ggatccagga ggataacaag ggaatctcct actctcaaag agtctgccat ctagtgggag
61  acgcaggaat gtaattgagt aggagaacac aatgagattc gttgcagaac agccatgaga
121 acagaacaaa gttctaaagag agcataaagg ggtggcaciaa cttaatTTta tcaaaaaaat
181 tcaggaacaa ttatacagag aggaggagtt tacaagtaac tatgtaggga gctgtcatgg
241 gtattccagt taaaggaaac atgtgaggag cataaaagag gctggcccat tgggttggct
301 gcacatgtat gtgtttgtta aggtttggga gtgtgtgagt gaatggtgga aggtgagTct
361 aaagggaag cagtactaga tcttgagcat tcttatatat cacaatgaaa gatttgaaat
421 aacatcctga ggcatctgaa gttagcaaaa gaggttctca gtagggaaat ggcattgatta
481 gattgaggct ttacagtgat taccctggca aagctgcaga gaacagactg agaggaggcc
541 ctggtctggg taaccagtta gtccactgta atttgcctaa catttgagca gtgtggggag
601 aaaggaggac acatctcaaa ggactaccta gaaggtatac ttagtccagt ttggttgaga
661 atgacatgta ggtggatagc aatgagctca agatgatccc tatatatcag tatttgaaa
721 ttagatggaa gagaacacat tgctccatgc taaggactaa tatgaggaga agcagtttga
781 atagaagatg tgtccagtgt tcaaggagtg attgtgcagt aaggtagaga tcattaaaga
841 gccagtttga agtttagaat gaagtctggg tgaaaaatca aatgcgatta gtgggaagtc
901 tcttagaggT taacctatac tttttatgaa aacataggaa tttatTTtca tattccctaa
961 tagcaaaagg cctttaacta cacatatatt taataaatac attttataga tacctatgta
1021 aaataaaaga caaccaccac acaacaaact cagtggcaga aaagttccag tgcaatcagt
1081 atatttaaat tctactgggg gtattgcaaa tcaaatttac attttgggag actatatggc
1141 aatatataat aagagctata aagatgatca tatcattttt cccagtaatc ctactcctgg
1201 ttattaaagg gaagtaattc agtgtatatT aatgaaatcg accttatagt tctgatcttt
1261 ataataagtt ataagaaaat ggtttaactt gtatgtgtat atatttactc gaagtagaaa
1321 tatgaaggct aaaaaaatgg gaagatatTT aaattggTgt taaaacagca tttaaaatta
1381 ccacaattat gagaacacat gtatgccaat gcagattcac tggaaaaata ttgaaaaatga
1441 aaactgtcag atggttaagat tataattTTta tttctTTTTt aatttgaaat aaattggtaa
1501 cagcacagct tttcaaaagc ttctataaat gtgtatgtta agttgtaata aagcaaacac
1561 atgcatgtaa gacatgctta aacagttatt taattgtTtc ttgggtacct ggggagatgg
1621 ggtgaagaaa ggggggtgac ttgaatgaag gtggaggaga aaaaTgagaa ccaagaagac
1681 aaaggatcga gaagctcagt gtggcagcag cctctctTcc cctcctgaga gagtcaaagg
1741 gtggcatcag ggactcatga tccatggttg tggaaagcctc atgtcacact ggatgtcaca
1801 tgaggtggga tggaaacacag tgaccacccc acctcatttc ctttacagct tccgtggggg
1861 ccatggcagt gaacagcctt caggcatgtc tacggtggaa gatctgaatt caggctggTg
1921 gcaggagaca acacaaccac gttttctTTt atgcatgcat ttggtTTaat tgacacatta
1981 accacagaca aaggggtaaa ggccacaagg cgataggTTa gtatgaacag ggaaagggac
2041 attttttttT ttaagaaaaa ataaaagcat cagtattTga aagactTTtc atgatcctat
2101 acccacctcg aaagcccccT ctcaccacag gaagtgcact gaccattTga ggcataaaag
2161 agatcctcaa agagcccgat cctcactctc cctgcacagc tcagcgggac ctcagccatg
2221 agactctTca tcttgccctc ccttggcacTc tgctctctca ctgcatacat tgtggaaaggT
2281 aagtggagaa gctgtctgtg agataaagaa tagggaggca aggcaggtgg gcacacattt
2341 tgggtttgac tcgggttttg actggactaa actgctgtct ccaggggagc cttaaacttc
2401 ccatgtgcaa gaaaggaatg atgattttga ctgtagaggg cttcgtaaac ttccaaaaca

```

```

2461 gggagaattt gatttagtgc tgggctccta cttttcctaä ttgggtaatt tcaggtaaat
2521 tccttaacca ctcagggcct gtgcttattt atgtataaac tgaatagaat aagagacatg
2581 atcacctgag attaaagatta aataaaatatt atggtttatt taataacatc agatttcctt
2641 acaagcagta attttttgat taatgttagc tatggattag aggtgatgat tataaatgca
2701 tttgtaggtt ttgccattt aatatatagt ttgataaatt atcaaatctc tagagagttc
2761 agttacaata tggggatgca ccagaggatg tatgttctgg agcaaatcaa tgttttcaat
2821 acaaaaacctg tgtgaaggcg acagtagtgc ttgctgtgga ctggatgtcc cagtcttgcc
2881 ttccctcccc ttgataatgc aataagggac cccattttta ggacgcagga caggcagaaa
2941 gataaccagg ttgatggggt ccacaccatg tgcaatcact accagctgag acttcttgtt
3001 ttccagcaag gtggtgatga tgttaacccc tgctcaaaga acaggtgatt tcctagtggg
3061 gacaaccctt ttgctagcag ctttcttctc agcctgggccc aacagtctct gcttcttctc
3121 ttgctttgtc tctggtcagt acttgtggat cagcttaagt ggctgagtag ctgtttgggg
3181 gtctaaggct tgggtgaact ggttaatggc agaaggcatt ttcagctgct tatagaggat
3241 agctctttgc agctggaacc agatatagcg gggccatttc acaaagcagt ggaggtctct
3301 tttgggctgg atgtcctgtc caatgcctgc ctaagaaaac tcttaggcct tttctcacac
3361 agcgggttca tcactttctt agcctcctgc ttcctcacga cggcagggac tgggcccact
3421 tctttccttt ggccttcttt ctttccagca tcttaggcag ctgacagaga gggaaatttg
3481 accattttaa aaggggaaaca cttttattta ctcagtcaaa agcatgtctc cttccctcac
3541 tgaatgttgc cttgcttaga gtactcttca cgcattactc tgtcatctca cttatggtac
3601 tgtaacatgt tgcactattt gaaatgatct tttctgtttg cctgtctgct gcctggctcc
3661 ctcagtagaa gatagctct atgaaaacag ggataatgtc tgtcttaata aaacatgttg
3721 gacacaacag gcaccattgt ataaatgaat gaatgcgtgt cactggggca tttgctagcc
3781 gtcccaaatg tctaagtga aataacacag gagacgggat aacatcttgt tatttctct
3841 cagcatgaaa ttctgaaac aattctgttg attgagtttt taaatttagc aaatatttac
3901 taagaatctg tgacgggcaa gagattcggg atgcttatca gtccctctct ccccaaaaa
3961 gcaaatggcc ttatattctc acaacattct cagagtaatt taacagacga ttgttctgt
4021 gatctgggta attgctttat ttttaattgt ctgtgttttt tttttcctca tcaggtgtag
4081 ggagtgaagt ctcacatagg aggacctgtg tgagcctcac taccagcga ctgccagtta
4141 gcagaatcaa gacctacacc atcacggaag gctccttgag agcagtaatg tgagtctgcc
4201 tcctcagaag ttgggctggg tgggtaccta gaggtataga aatacactct atagaaatgc
4261 tgccatcctc aggaaaagta ggtcagcata gaggaacacc tcaacttaac caaaaacctc
4321 tttagttttc cttatcaacc atgtctttct gcagcccaac cgaatagcga ttattgcaga
4381 aattgggctg ccaaagaaag aatagaagtc ctctctatt tgtcttagtg gaagagtctg
4441 ttgaatactg tgcacagctc tgagatctgg gtttagagat ggctgggcta tgtcagggtt
4501 tccctgcaag cctcactgga gttgggggat cttagggttg agttaggcag agtcccatac
4561 tttatcagtt gccatatttc aagaaaatga gtcaatgcac aacctacatg gtccctttct
4621 tctaccagaa tctcattttt agaagtaata actcttccca atacatattg caagctttgc
4681 tctaaagaat gaaaatgtaa aaatcacctt tttaaaaaaa ataagatgag tattttcaaa
4741 tttgaaaagg aagaggttat ataataatgg aactagatgg cctcaaattg ctttttggtta
4801 caacatttgg tgacatggat gagaaaagga gcctgtgaat tatggtgaac aaaggggctg
4861 gatactactt gcagatattt ctcttttatg ttaaaaataga tggcagaaga aggggtgctc
4921 tttatgatct catggctctg aaagactatt tcttgacgta atttctgcac aagatctctt
4981 catgtctgcc ctgatcttaa ctcctgaccc tgaggctttg agaacgtggc taacttcac
5041 tgtcttttcc ttgcgttaca gttttattac caaacgtggc ctaaaagtct gtgctgatcc
5101 acaagccacg tgggtgagag acgtgggtcag gagcatggac aggaaatcca acaccagaaa
5161 taacatgac cagaccaagc caacaggaac ccagcaatcg accaatacag ctgtgacctt
5221 gactggctag tagtctctgg caccctgtcc gtctccagcc agccagctca tttacttta
5281 caccctcatg gactgagatt atactcacct tttatgaaag cactgcatga ataaaattat
5341 tcctttgtat ttttactttt aaatgtcttc tgtattcact tatatgttct aattaataaa
5401 ttattttatta ttaagaatag ttccctagtc tattcattat atttagggaa aggtagtgtg
5461 tcattgttgt ttgatttctg acctgtgacc tctctttgat ggtaaccata atggaagaga
5521 ttctggctag tgtctatcag aggtgaaagc tatatcgatc actcttagag tccagcttgt
5581 aatggttctt tacacatcag tcacaagtta cagctgtgac aatggcaaca atttgagatc
5641 tatttcaact tgtctctata atagaattc

```

//

LOCUS D63790 5660 bp DNA PRI 27-DEC-1996
 DEFINITION Human DNA for SCM-lalpha precursor, complete cds.
 ACCESSION D63790
 NID g1754610
 KEYWORDS SCM-lalpha precursor; SCM-1 alpha.
 SOURCE Homo sapiens placenta DNA, clone:hg40.
 ORGANISM Homo sapiens
 Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
 Vertebrata; Mammalia; Eutheria; Primates; Catarrhini;
 Hominidae;
 Homo.
 REFERENCE 1 (sites)
 AUTHORS Yoshida,T., Imai,T., Kakizaki,M., Nishimura,M. and Yoshie,O.
 TITLE Molecular cloning of a novel C or gamma type chemokine, SCM-1
 JOURNAL FEBS Lett. 360 (2), 155-159 (1995)

MEDLINE 95180438
REFERENCE 2 (sites)
AUTHORS Yoshida,T., Imai,T., Takagi,S., Nishimura,M., Ishikawa,I.,
Yaoi,T.
and Yoshie,O.
TITLE Structure and expression of two highly related genes encoding
SCM-1/human lymphotactin
JOURNAL FEBS Lett. 395 (1), 82-88 (1996)
MEDLINE 97002294
REFERENCE 3 (bases 1 to 5660)
AUTHORS Yoshida,T.
JOURNAL Unpublished (1995)
REFERENCE 4 (bases 1 to 5660)
AUTHORS Yoshida,T.
TITLE Direct Submission
JOURNAL Submitted (07-AUG-1995) to the DDBJ/EMBL/GenBank databases.
Tetsuya Yoshida, Shionogi Institute for Medical Science; 2-5-1,
Mishima, Settsu, Osaka 566, Japan (E-
mail:teyoshid@fl.lab.shionogi.co.jp,
Tel:06-382-2612, Fax:06-382-2598)
FEATURES Location/Qualifiers
source 1..5660
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="1"
/clone="hg40"
/map="1q23"
/tissue_type="placenta"
TATA_signal 640..644
exon 683..764
/number=1
prim_transcript 683..5340
CDS join(704..764,4064..4178,5053..5221)
/codon_start=1
/product="SCM-1alpha precursor"
/db_xref="PID:d1010505"
/db_xref="PID:g1754611"
/translation="MRLILALLGICSLTAYIVEGVGSEVSDKRTC VSLTTQRLPVSR
IKTYTITEGSLRAVIFITKRGLKVCADPQATWVRDVVRSMRKSNTNRNMIQTKPTGT
QQSTNTAVTLTG"
intron 765..4063
/number=1
exon 4064..4178
/number=2
mat_peptide join(4066..4178,5053..5218)
/note="SCM-1alpha mature peptide"
intron 4179..5052
/number=2
exon 5053..5340
/number=3
BASE COUNT 1623 a 1139 c 1175 g 1723 t
ORIGIN
1 aagcttctat aaatgtgtat gttaagttgt aataaagcaa acacatgcat gtagacatgc
61 ttaaacagtt atttaattgt ttcttgggta cctggggaga tgggggtgaag aaaggggggt
121 gacttgaatg aaggtggagg agaaaaatga gaaccaagaa agcaaaggat cgagaagctc
181 agtgtggcag cagctctctt cccctcctga gagagtcaaa ggggtggcatc agggactcat
241 gatccatggt tgtggaagcc tcatgtcaca ctggatgtca catgagggtgg gatggaacac
301 agtgaccacc ccacctcatt tcctttacag cttccgtggg gggccatggc agtgaaacacc
361 ttcaggcatg tctacggcgg aatatctgaa ttcaggctgg tggcaggaga caacacaacc
421 acgttttctt ttatgcatgc atttggttta attgacacat taaccacaga caaaggggta
481 aaggccacaa ggcgttaggt tagtatgaac agggaaaagg gacttttttt ttttttttta
541 agaaaaataa aagcatcagt attgcaaaga ctttccatga tcctacaccc acctcgaaag
601 cccctcttca ccacaggaag tgcactgacc actggaggca taaaagagggt cctcaaagag
661 cccgatcctc actctccttg cacagctcag caggacctca gccatgagac ttctcatcct
721 ggccctcctt ggcatctgct ctctcactgc atacattgtg gaaggtaagt ggagaagctg
781 tctgtgagat aaagaacagg gaggcaaggc aggtgggcac acattttggg tttgactcag
841 gttatgactg gactaatctg ctttcccag gggagcctta aacttcccat gtgcaagaaa


```

901 ggaatgatga ttttgactgt agagggcttc gtaaaacttc aaaacagggga gaatttgatt
961 agtatcttgg ctcctacttt tcctaattgg gtaatttcag gtaaatccct taaccactca
1021 gggcctgtgc ttatttatgt ataactgaat agtataacag acttgatcac ctgagattaa
1081 gattaaataa atattatggg ttatttaata acatcagatt tccttacaag cagtaatttt
1141 ttgattaatg ttagctatgg attagagggt atgattataa atgcatttga aggttttggc
1201 catttaatat atagtttgat aaattatcaa aatcttagag agttcagtta cagtggtggg
1261 atgcaccatt ggatgtatgt tctggagtaa atcaatgttt tcaatacaaa actaagcccc
1321 aaatgactgg aagttcaaac ctcatgtccc agaaaatcaa tattaccttc aagtacgtgg
1381 gggactctgt tagtaatgcc atgactatta ctatttatga gaaattttct gttttgttaa
1441 gagaacatac aataataact actaccaaag agatcagcac cttatacaca gtttcaataa
1501 cctgcaagac acatccaggt aagattcaga tataccgagc ccttacctga gcattcagta
1561 ggtatttttt aaggattgat ttttccatag actggagggt aatctgtcga cttattttgt
1621 ttctagttgg taggcttatt acttagacta tgataattata acttaataat ggttcccaa
1681 ggggttccat gaataaagggt ggctaagtct ggaagtcctt gaaattatgg ataaaaaaa
1741 aaaatactga tgaacaaaaa gagtttgatt actacattag gccacatgtt gctacctggc
1801 tggcattttg ctgagacaaat gggcatacca tttgagggag actcagatct gagtagggga
1861 aaggagctct ataagtccca ctggtgctta gcttcttaca tacaaaaatg agggaaaacg
1921 gtctctgctt tgactcaatt ttgcaacctg agtgaagggt atatttttaa aaataacaca
1981 gacactcaaa cattgtctgac aataaggaaa aggtcttgtg gtttcaagca taacaggatt
2041 ccttgagtct taggagtcca cttcagatac ttcacagaga gaaatattgt ttcttaataa
2101 tgagagaaac agagaaaaaa cccagatttt tcctctttca ttggctacag aaacaattca
2161 ccactaaaaa taaattggca aaggtagagg atagcaatgt gcagactggc attgagagtg
2221 aagaaatgat gaagaaaagc acacaatgaa cactctttgt ttagtccctt gctttaaaaa
2281 atgccttctg atattagcaa cactacagac caatgttggc cattatcagt gggttactta
2341 gatgcttttt agctgcctat ttccctggga agcaaagacc agtgtctaca gcttaaggga
2401 aaatcagcac ttagaaactt ggattagatt tcacccaacc cttaacagta ttaattctcc
2461 caagttattt ttccctcatg aatgtttttt tgattctcta cacttaatat ttaattctct
2521 ttgggcattt actattgggg atgcataatt aagggtctgac ttccctttat atatatctta
2581 ctttttacca tttaattaatt ttttgagag tttttattat ttttatgtac agaaaactca
2641 acagtgtaca tttaaccagc tttagtggca agttcctctg cctttgtctat tccagcttg
2701 gcattgtgag ccacagattt tggactcggg acattgcaga tctcatcata tccgtcattg
2761 taatttgtcc tgatagcttc caccagctta gccaaagctc ctttgtcttc ctggttaact
2821 tgtgtgaagg ccacagtggg gcttctctgt gactggatgt cccagtcttg ccttcttacc
2881 ccttgataat gcattaaagg acccccat ttaggacaca ggacagacag aaagttaacc
2941 agcttgatgg ggtccacacc atgtgcaata ccagctgagc cttctcttct tccagcaagg
3001 tgggtatagt gttaaacctt gctcaaagaa caggtgattt cctagtgggg acaacctctt
3061 tgctagcagc tttcttctca gctggggcca acagtctctg cttcttctct tgctttgtct
3121 ctggtcagta cttgtggatc agcttaagtg gctgagtagc tgtttggggg tctaaggctt
3181 ggggtgaactg gttaatggca gaaggcactt tcagctgctt atagaggata gctctttgca
3241 gctggaacca gatatagcgg ggccatttca caaagcagcg gaggtccctt ttctcacaca tggggttcat
3301 tgtcctgtcc aatgcctgcc taagaaaact cttaggcctt ttctcacaca ggggttctat
3361 cactttctta gccctctgct tcttcacgac ggcagggtct gggggcactt tcttctctt
3421 ggccatcttt cctttcagca tcttaggcag ctgacagaga gagacatttg accatttaaa
3481 aaggagaaca cttttattta gtctgtcaaa agcatgcttc cttccctcac tgaatgttgc
3541 cttgcttaga gtactcttca cgcattactc tgtcttctca ctatggtact gtaacatgtt
3601 gcactatttg aaatgatctt ttctgtttgc ctgtctgctg cctggctcct tcatgagaga
3661 gatatgctct atgaaaacag gagtaatgtc tgcttagtaa aacatgtggg acacaacagg
3721 caccatgtga taaatgaatg aatgcgtgtc actggggcat ttgctagccg tcccaaatgt
3781 ctaagtgaat atatacacag agacgggata acatcttgtt attttctctc agcatgaaat
3841 tcttgaaaca attctgttga ttgagttttt aaattagtca aatatttact aagaatctgt
3901 gacgggcaag agattcggga tgccatcagc tcctctcttc ccccaaaaag caaatggcct
3961 taaattctca caacattctc agagtaattt aacagatgat tgttctctgt atctggataa
4021 ttgctttatt tttaattgtc tgttgttttt ttttctctac cagggttagg gagtgaagtc
4081 tcagataaga ggacctgtgt gagcctcact acccagcgac tgccggttag cagaatcaag
4141 acctacacca tcacggaagg ctcttgaga gcagtaatgt gagtctgcct cctcagaagt
4201 tgtgtgggtt gggatatctg aagtatagaa atacactctg tagaaatgct gccgtcctca
4261 ggaaaagtag gtcagcatag aggaacactt caacttaacc aaaaacctct ttagttttcc
4321 ttatcaatca tgtctttctg cagcccaacc gaatagcgat tattgcagaa attgggctgc
4381 caaagaaaga atagaagtcc tctctatatt agcttagtgg aagagtctgt tgaatactgt
4441 gcacagctct gagacctggg tttagagatg actggcccat gtcagggttt ccttgcaagc
4501 ctactggagg ttgggggatc tttaggttga gtcaggcaga gtcccatatt tttcaggt
4561 gccataattc aagaaaatga gtcaatgcac aacctacatg gtccctctct ctaccagaat
4621 ctcattttta gaagttaata actcttctca acatgtaatt gcaagcttta ctctaaaaaa
4681 tgaaaaatga aaaatcactt tttattttta aaataagatg aatattttta aatttgaaaa
4741 ggaagagggt atgttaataa ggaactagt tttattgtga aaactcaggt acaacatttg
4801 gtgacatgga tgagaaaagg accctgtgaa ttattgtgaa caaaggggct ggatactact
4861 tgcagatatt actcctttat gttaaaaatg atggcagaag aaggggtact atttatgatc
4921 tcatggctct gaaagactat ttcttgcaat aatttctgca caagatctct tcatgtctgc
4981 cctgactcta actcctgacc ctgaggcttt gagaatgtgg ctaactctgt ctgtcttttc
5041 cttgcgttac agttttatta ccaaactgtg cctaaaagtc tgtgtctgat cacaagccac
5101 atgggtgaga gacgtgggtc ggagcatgga caggaaatcc aacaccagaa ataacatgat

```

```

5161 ccagaccaag ccaacaggaa cccagcaatc gaccaataca gctgtgactc tgactggcta
5221 gtagtctctg gcaccctgtc cgtctccagc cagccagctc atttcacttt acacgctcat
5281 ggactgagtt tatactcacc ttttatgaaa gcactgcatg aataaaatta ttcttttgta
5341 tttttacttt taaatgtctt ctgtattcac ttatatgttc taattaataa attattttatt
5401 attaagaata gttccctagt ctattcatta tatttaggga aaggtagtgt atcattgttg
5461 tttgatttct gacctgttac ctctcttga tggtaaccaat aatggaagag attctggcta
5521 gtgtctatca gaggtgaaag ctatatcaat ctctcttaga gtccagcttg taatggttct
5581 ttacacatca gtcacaagtt acagctgtga caatggcaac aatttgagat gtatttcaac
5641 ttgtctctat aatagaattc

```

//

```

LOCUS      HSU91835      1635 bp      mRNA      PRI      21-MAR-1997
DEFINITION Human CX3C chemokine precursor, mRNA, alternatively spliced,
            complete cds.
ACCESSION  U91835
NID        g1899258
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1635)
AUTHORS    Bazan,J.F., Bacon,K.B., Hardiman,G., Wang,W., Soo,K., Rossi,D.,
            Greaves,D.R., Zlotnik,A. and Schall,T.J.
TITLE      A new class of membrane-bound chemokine with a CX3C motif
JOURNAL    Nature 385 (6617), 640-644 (1997)
MEDLINE    97177111
REFERENCE  2 (bases 1 to 1635)
AUTHORS    Bazan,J.F., Bacon,K.B., Hardiman,G., Wang,W., Rossi,D.,
            Greaves,D.R., Zlotnik,A. and Schall,T.J.
TITLE      Direct Submission
JOURNAL    Submitted (03-MAR-1997) Molecular Biology, DNAX Research
Institute, 901 California Ave., Palo Alto, CA 94304-1104, USA
FEATURES
  source    Location/Qualifiers
            1..1635
            /organism="Homo sapiens"
            /db_xref="taxon:9606"
  CDS       80..1273
            /note="membrane-tethered chemokine module"
            /codon_start=1
            /product="CX3C chemokine precursor"
            /db_xref="PID:g1899259"

/translation="MAPISLSWLLRLATFCHLTVLLAGQHGVTKCNITCSKMTSKIP
VALLIHYQQNQASC GKRAIIETRQHRLFCADPKEQWVKDAMQHLDRQAAALTRNGGT
FEKQIGEVKPRTPPAAGGMDES VVLEPEATGESSSLEPTPSSQEAQRALGTSPELPTG
VTGSSGTRLPPTPKAQDGGPVGTFLFRVPPVSTAATWQSSAPHQPGPSLWAEAKTSEA
PSTQDPSTQASTASSPAPEENAPSEGRVWGQGQSPRPENSLEREEMGPVPAHTDAFQ
DWGPGSMAHVSVPVSSEGTSPREPVASGSWTPKAEPIHATMDPQRLGVLITPVPDA
QAATRRQAVGLLAFLGLLFC LGVAMFTYQSLQGCPRKMAGEMA EGLRYIPRSCGSNSY
VLVPV"
  sig_peptide      80..151
  mat_peptide      152..1270
                  /product="CX3C chemokine"
  misc_feature     152..379
                  /note="encodes chemokine module"
  misc_feature     380..1102
                  /note="encodes glycosylation stalk"
  misc_feature     1103..1159
                  /note="encodes transmembrane helix"
  misc_feature     1160..1270
                  /note="encodes intracellular domain"
  3'UTR            1274..1635
                  /note="alternatively spliced; long transcript can be

```

found

in GenBank Accession Number U84487"

BASE COUNT

338 a 544 c 464 g 289 t

ORIGIN

```

1  ggcacgaggg cactgagctc tgccgcctgg ctctagccgc ctgcctggcc cccgcccggga
61  ctcttgccca ccctcagcca tggctccgat atctctgtcg tggctgtccc gcttggccac
121  cttctgccat ctgactgtcc tgctggctgg acagcaccac ggtgtgacga aatgcaacat
181  cactgtcagc aagatgacat caaagatacc tgtagctttg ctcatccact atcaacagaa
241  ccaggcatca tgccggcaaac gcgcaatcat cttggagacg agacagcaca ggctgttctg
301  tgccgacccg aaggagcaat gggtaagga cgcgatgcag catctggacc gccaggctgc
361  tgccctaact cgaaatggcg gcaccttcga gaagcagatc ggcgaggtga agcccaggac
421  cacccttgcc gccgggggaa tggacgagtc tgtggtcctg gagcccgaag ccacaggcga
481  aagcagtagc ctggagccga ctcttctctc ccaggaagca cagagggccc tggggacctc
541  cccagagctg ccgacgggcg tgactggttc ctcagggacc aggtctcccc cgacgcaaaa
601  ggctcaggat ggagggcctg tgggcacgga gcttttccga gtgcctcccc tctccactgc
661  cgccacgtgg cagagttctg ctccccacca acctggggcc agcctctggg ctgaggcaaa
721  gacctctgag gccccgtcca cccaggacce ctccaccag gcctccactg cgtctcccc
781  agccccagag gagaatgtc cgtctgaagg ccagcgtgtg tggggtcagg gacagagccc
841  caggccagag aactctctg agcgggagga gatgggtccc gtgccagcgc acacggatgc
901  ctccaggac tgggggcctg gcagcatggc ccacgtctct gtggtccctg tctctcaga
961  agggaccccc agcaggagc cagtggcttc aggcagctgg acccctaagg ctgaggaacc
1021  catccatgcc accatggacc cccagaggct gggcgtcctt atcactcctg tccttgacgc
1081  ccaggctgcc acccggaggc aggcgggtgg gctgctggcc ttccttggcc tctcttctg
1141  cctgggggtg gccatgttca cctaccagag cctccagggc tgccctcgaa agatggcagg
1201  agagatggcg gagggccttc gctacatccc ccggagctgt ggtagtaatt catatgtcct
1261  ggtgcccgtg tgaactctc tggcctgtgt ctagtgtttt gattcagaca gctgcctggg
1321  atccctcatc ctcataccca cccccacca agggcctggc ctgagctggg atgattggag
1381  gggggaggtg ggatcctcca ggtgcacaag ctccaagctc ccaggcattc cccaggaggc
1441  cagccttgac cattctccac ctccagggga cagagggggt ggctcccaa ctcacccag
1501  ccccaaaact ctctctgct gctggctggt tagaggttcc ctttgacgcc atcccagccc
1561  caatgaacaa ttatttatta aatgcccagc cccttctgaa aaaaaaaaaa aaaaaaaaaa
1621  aaaaaaaaaa aaaaa

```

//

```

LOCUS      HSU84487      3310 bp      mRNA      PRI      15-MAR-1997
DEFINITION Human CX3C chemokine precursor, mRNA, alternatively spliced,
            complete cds.
ACCESSION  U84487
NID        g1888522
KEYWORDS   .
SOURCE     human.
  ORGANISM Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 3310)
AUTHORS    Bazan,J.F., Bacon,K.B., Hardiman,G., Wang,W., Soo,K., Rossi,D.,
            Greaves,D.R., Zlotnik,A. and Schall,T.J.
TITLE      A new class of membrane-bound chemokine with a CX3C motif
JOURNAL    Nature 385 (6617), 640-644 (1997)
MEDLINE    97177111
REFERENCE  2 (bases 1 to 3310)
AUTHORS    Bazan,J.F., Bacon,K.B., Hardiman,G., Wang,W., Rossi,D.,
            Greaves,D.R., Zlotnik,A. and Schall,T.J.
TITLE      Direct Submission
JOURNAL    Submitted (07-JAN-1997) Molecular Biology, DNAX Research
Institute, 901 California Ave., Palo Alto, CA 94304-1104, USA
FEATURES   Location/Qualifiers
            source      1..3310
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
            CDS          80..1273
                        /note="membrane-tethered chemokine module"
                        /codon_start=1
                        /product="CX3C chemokine precursor"
                        /db_xref="PID:g1888523"

```

/translation="MAPISLSWLLRLATFCHLTVLLAGQHGVTKCNITCSKMTSKIP

VALLIHYQQNQASCGKRAIILETRQHRLFCADPKEQWVKDAMQHLDRQAAALTRNGGT

FEKQIGEVKPRRTTPAAGGMDESIVLEPEATGESSSSLEPTPSSQEAQRALGTSPELPTG
 VTGSSGTRLPPTPKAQDGGPVGTELFRVPPVSTAATWQSSAPHQPGPSLWAEAKTSEA
 PSTQDPSTQASTASSPAPEENAPSEGQRVWVGQGSPPRPNLSLEREEMGPVPAHTDAFQ
 DWGPGSMAHVSVPVVSSEGTSPREPVASGSWTPKAEPIHATMDPQRLGVLITPVPDA

QAATRRQAVGLLAFLLFCLGVAMFTYQSLQGCPRKMAGEMAEBGLRYIPRSCGSNSY

VLVPV"
 sig_peptide 80..151
 mat_peptide 152..1270
 /product="CX3C chemokine"
 misc_feature 152..379
 /note="encodes chemokine module"
 misc_feature 380..1102
 /note="encodes glycosylation stalk"
 misc_feature 1103..1159
 /note="encodes transmembrane helix"
 misc_feature 1160..1270
 /note="encodes intracellular domain"
 3'UTR 1274..3310
 /note="alternatively spliced; short transcript

deposited

as GenBank Accession Number U91835"

BASE COUNT 659 a 1051 c 916 g 682 t 2 others
 ORIGIN

```

1  ggcacgaggg cactgagctc tgccgcctgg ctctagccgc ctgcctggcc cccgccggga
61 ctcttgccca ccctcagcca tggctccgat atctctgtcg tggctgctcc gcttggccac
121 cttctgccat ctgactgtcc tgctggctgg acagcaccac ggtgtgacga aatgcaacat
181 cactgacgac aagatgacat caaagatacc tgtagctttg ctcattccat atcaacagaa
241 ccaggcatca tgccggcaaac gcgcaatcat cttggagacg agacagcaca ggctgttctg
301 tgccgacccg aaggagcaat gggatcaagg cgcgatgcag catctggacc gccaggctgc
361 tgccctaact cgaaatggcg gcaccttcca gaagcagatc ggcgagggtga agcccaggac
421 caccctcgcc gccgggggaa tggacgagtc tgtgttctct gagcccgaa ccacaggcga
481 aagcagtagc ctggagccga ctcttctctc ccaggaagca cagagggccc tggggacctc
541 cccagagctg ccgacgggag tgactggttc ctccaggacc aggtctcccc cgacgccaaa
601 ggctcaggat ggagggcctg tgggcacgga gcttttccga gtgcctcccc tctccactgc
661 cgccacgtgg cagagtcttg ctccccacca acctgggccc agcctctggg ctgaggcaaa
721 gacctctgag gccccgtcca cccaggaccc ctccaccagc gcctccactg cgtcctcccc
781 agccccagag gagaatgttc cgtctgaagg ccagcgtgtg tggggtcagg gacagagccc
841 caggccagag aactctctgg agcgggagga gatgggtccc gtgccagcgc acacggatgc
901 cttccaggac tgggggcctg gcagcatggc ccacgtctct gtggtccctg tctcctcaga
961 agggaccccc agcaggagag cagtggcttc aggcagctgg acccctaagg ctgaggaaac
1021 catccatgcc accatggacc cccagaggtt gggcgctcct atcactcctg tccctgacgc
1081 ccaggctgcc acccggaagg aggcggtggg gctgctggcc ttccttggcc tctcttctg
1141 cctgggggtg gccatgttca cctaccagag cctccagggc tgccctcgaa agatggcagg
1201 agagatggcg gagggccttc gctacatccc ccggagctgt ggtagtaatt catatgtcct
1261 ggtgccccgt tgaactcctc tggcctgtgt ctagtgtgtt gattcagaca gctgcctggg
1321 atccctcctc ctcatacca cccccacca agggcctggc ctgagctggg atgattggag
1381 gggggagggt ggatcctcca ggtgcacaag ctccaagctc ccaggcatte cccaggaggc
1441 cagccttgac cattctccac ctccaggga cagagggggg ggcctcccaa ctcacccagc
1501 ccccaaaact ctctctgtgt gctggctggg tagagggttc ctttgacgcc atcccagccc
1561 caatgaacaa ttatttatta aatgccagc cccttctgac ccatgctgcc ctgtgagtac
1621 tacagtcttc ccatctcaca catgagcacc agggccaggc ctctgccacc tccctgcaac
1681 ctgatttgtt ctcttggttc tgctgcagtt gccagtcacc ccggccacct ggggtgctat
1741 ctccccagc cccatcctct gtacagagcc cagcctccca ctggtgacat gtcttttctt
1801 gcatgaggct agtgtgtgtt ttctgggcca ctgcttccag tgaggctctg cccttggtta
1861 ggsatttgtg gaaggggaga taaggggtat tgggtgacttt cctctttggt ctacactgtg
1921 ctgagtctga aggcgtgggt ctgacccatg ttccaccatc aagccaccaa catactccca
1981 tctgtgaaag gaaagaggga ggttaaggaa acctgtcccc ctgacaacac tcattgacct
2041 gaggcccttc tctccagccc ctggatgcag cctcacagtc cttaccagca gagcacctta
2101 gacagtccct gccaatggag taacttgtct ttggaccctg agggccagag ggcctgcarg
2161 gatagtagtt gatagcacag acctgcccct gtgggcccc aaatggaaat gggcagagca
2221 gagaccatcc ctgaaggccc cgcccaggct tagtcaactg gacagcccgg gctctgcttc
2281 ccatcaccgc ctaagaggga gggagggtct cagacacatg tccaagaagc ccaggaaagg
2341 ctccaggagc agccacattc ctgatgtctc ttccagagat cctgcaggca gccaggccac
2401 aagacccttg tggctccacc ccacacagc cagattcttt cctgaggctg ggctcccttc
2461 ccacctctct cactccttga aaacactgtt ctctgcccct caagaccttc tcttccacct
2521 ttgtccccac cgcagacagg accaggggat ttccatgatg ttttccatga gtccctgtt
2581 tgtttctgaa agggacgcta cccgggaagg gggctgggac atgggaaagg ggaagtgtga

```

```

2641 ggcataaagt caggggttcc cttttttggc tgctgaaggc tcgagcatgc ctggatgggg
2701 ctgcaccggc tggcctggcc cctcagggtc cctgggtggc gctcacctct cccttgatt
2761 gtecccgacc cttgccgtct acctgagggg cctcttatgg gctgggttct acccaggtgc
2821 taggaacact ctttcacaga tgggtgcttg gaggaaggaa acccagctct ggtccataga
2881 gagcaaaacg ctgtgctgcc ctgcccaccc tggcctctgc actcccctgc tgggtgtggc
2941 gcagcatatt caggaagctc agggccctgg ctcagggtgg gtactctgg cagctcagag
3001 aggggtggag tgggtccaat gcactttgtt ctggctcttc caggctggga gagcctttca
3061 ggggtgggac accctgtgat ggggccctgc ctcttttggt aggaagccgc tggggccagt
3121 tgggtccctt tccatggact ttgttagttt ctccaagcag gacatggaca aggatgatct
3181 aggaagactt tggaaagagt aggaagactt tggaaagact ttccaaccc tcatcaccaa
3241 cgtctgtgcc attttgtatt ttactaataa aatttaaaag tcttgtgaaa aaaaaaaaaa
3301 aaaaaaaaaa

```

//

```

LOCUS      HSU91746      1430 bp      mRNA                      PRI      12-MAR-1998
DEFINITION Homo sapiens IL-10-inducible chemokine (HCC-4) mRNA, complete
cds.
ACCESSION  U91746
NID        g2581780
KEYWORDS
SOURCE     human.
ORGANISM   Homo sapiens
            Eukaryotae; mitochondrial eukaryotes; Metazoa; Chordata;
            Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 1430)
AUTHORS    Hedrick,J.A., Helms,A., Gorman,D. and Zlotnik,A.
TITLE      Identification of a novel human CC chemokine upregulated by IL-
10
JOURNAL     Blood (1998) In press
REFERENCE  2 (bases 1 to 1430)
AUTHORS    Hedrick,J.A., Helms,A., Gorman,D. and Zlotnik,A.
TITLE      Direct Submission
JOURNAL     Submitted (02-MAR-1997) Immunology, DNAX Research Institute,
901
            California Ave, Palo Alto, CA 94304, USA
FEATURES
  source    Location/Qualifiers
            1..1430
            /organism="Homo sapiens"
            /db_xref="taxon:9606"
            /chromosome="17"
  gene      1..1430
            /gene="HCC-4"
  CDS       1..363
            /gene="HCC-4"
            /note="CC or beta chemokine family member"
            /codon_start=1
            /product="IL-10-inducible chemokine"
            /db_xref="PID:g2581781"

/translation="MKVSEAALSLVLILIIITSASRSQPKVPEWVNTPTCCLKYYEK
VLPRLVVGYRKALNCHLPAIIFVTKRNREVCTNPNDWVQEIYKDPNLPLLPTRNLS
TVKIITAKNGQPQLLSQ"
BASE COUNT      401 a      351 c      293 g      385 t
ORIGIN
  1 atgaaggtct ccgaggtctgc cctgtctctc cttgtctctca tccttatcat tacttcgggt
  61 tctcgcagcc agccaaaagt tcctgagtgg gtgaacaccc catccacctg ctgectgaag
  121 tattatgaga aagtgttgcc aaggagacta gtggtgggat acagaaaagg cctcaactgt
  181 cacctgccag caatcatctt cgtcaccaag aggaaccgag aagtctgcac caaccccaat
  241 gacgactggg tccaagagta catcaaggat cccaacctac ctttgctgcc taccaggaac
  301 ttgtccacgg ttaaaattat tacagcaaag aatggtcaac cccagctcct caactcccag
  361 ttatgaccag gctttagtgg aagcccttgt ttacagaaga gaggggtaaa cctatgaaaa
  421 caggggaagc cttattaggg tgaaactagc cagtcacatt gagagaagca gaacaatgat
  481 caaaataaag gagaagtatt tcgaatatct tctcaatctt aggaggaaat accaaagtta
  541 agggacgtgg gcagaggtac gctcttttat ttttatattt atatttttat tttttgaga
  601 taggtcttac tctgtcacc caggctggag gcagtggtgt gatcttggt cacttgatct
  661 tggctcactg taacctccac ctcccaggct caagtgatcc tcccaccca gccctccgag
  721 tagctgggac tacaggcttg cgccaccaca cctggctaat ttttgattt ttggtagaga
  781 cgggattcta ccatgttgcc caggctggtc tcaaactcgt gtgcccaagc aatccacctg
  841 cctcagcctt ccaaaagtgc tgggattaca ggcgtgagcc accacatccg gccagtcac
  901 tcttaataca cagaaaaata tatttcacat ccttctcctg ctctctttca attcctcact

```

```
961 tcacaccagt acacaagcca ttctaaatac ttagccagtt tccagccttc cagatgatct
1021 ttgccctctg ggtcttgacc cattaagagc cccatagaac tcttgatttt tcctgtccat
1081 ctttatggat ttttctggat ctatattttc ttcaattatt ctttcatttt ataatgcaac
1141 tttttcatag gaagtcgga tgggaatatt cacattaatc atttttgcag agactttgct
1201 agatcctctc atattttgtc ttctctcaggg tggcaggggt acagagagtg cctgattgga
1261 aaaaaaaaaa aaagagagag agagagaaga agaagaagaa gagacacaaa tctctacctc
1321 ccatgttaag ctttgcagga cagggaaaga aagggtatga gacacggcta ggggtaaact
1381 cttagtccaa aaccaagca tgcaataaat aaaactccct tatttgacaa
```

//

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26291

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/84, 85.1, 184.1, 186.1, 188.1, 278.1; 514/2, 8, 12, 44; 530/300, 324

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | US 5,141, 867 A (IVANOFF et al.) 25 August 1992, see entire document. | 22-32, 45-55 |
| A | ENG et al. The Stimulatory Effects of Interleukin (IL)-12 On Hematopoiesis Are Antagonized by IL-12-induced Interferon γ In Vivo. J. Exp. Med. May 1995, Vol.181, pages 1893-1898, see entire document. | 1-21, 33-44 |
| A | ORANGE et al. Mechanism of Interleukin 12-mediated Toxicities during Experimental Viral Infections: Role of Tumor Necrosis Factor and Glucocorticoids. J. Exp. Med. March 1995, Vol.181, pages 901-914, see entire document. | 1-21, 33-44 |



Further documents are listed in the continuation of Box C.



See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *E* earlier document published on or after the international filing date | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *A* document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

25 MARCH 1999

Date of mailing of the international search report

15 APR 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/26291

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | WU et al. Receptor-mediated in Vitro Gene Transformation by a Soluble DNA Carrier System. The Journal of Biological Chemistry. 05 April 1987, Vol.252, No. 10, pages 4429-4432, see entire document. | 22-32, 45-55 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/26291

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/26291

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C07K 14/47, 14/52; C12N 15/12, 15/19, 15/63; A61K 38/16, 38/19, 48/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/84, 85.1, 184.1, 186.1, 188.1, 278.1; 514/2, 8, 12, 44; 530/300, 324

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, CAN ONLINE, MEDLINE, CAPLUS

search terms: chemokine, vaccination, immunogenic, antigen, HIV, efficacy, macrophage-derived chemokine, stromal cell-derived factor, monocyte chemotactic protein, composition, administration

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1.

Group I, claims 1-21, 33-44, drawn to a method to enhance the efficacy of a vaccine in a subject comprising administering an antigen and one or more chemokines and a composition thereto.

Group II, claims 22-32, 45-55, drawn to a method to enhance the efficacy of a vaccine in a subject comprising administering nucleic acid sequences encoding one or more antigens and nucleic acid sequences encoding one or more chemokines.

The inventions listed as Groups I-II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The inventions listed as Groups I-II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first-recited product and method, a method to enhance the efficacy of a vaccine in a subject comprising administering an antigen and one or more chemokines and a composition thereto. Further pursuant to 37

C.F.R. § 1.475 (d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.